

OPEN PETRI NETS

John C. Baez

Department of Mathematics
University of California
Riverside CA, USA 92521
and

Centre for Quantum Technologies
National University of Singapore
Singapore 117543

Jade Master

Department of Mathematics
University of California
Riverside CA, USA 92521

email: baez@math.ucr.edu, jmast003@ucr.edu

November 30, 2018

ABSTRACT. The reachability semantics for Petri nets can be studied using open Petri nets. For us an ‘open’ Petri net is one with certain places designated as inputs and outputs via a cospan of sets. We can compose open Petri nets by gluing the outputs of one to the inputs of another. Open Petri nets can be treated as morphisms of a category $\mathbf{Open}(\mathbf{Petri})$, which becomes symmetric monoidal under disjoint union. However, since the composite of open Petri nets is defined only up to isomorphism, it is better to treat them as morphisms of a symmetric monoidal *double* category $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$. Various choices of semantics for open Petri nets can be described using symmetric monoidal double functors out of $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$. Here we describe the reachability semantics, which assigns to each open Petri net the relation saying which markings of the outputs can be obtained from a given marking of the inputs via a sequence of transitions. We show this semantics gives a symmetric monoidal lax double functor from $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$ to the double category of relations. A key step in the proof is to treat Petri nets as presentations of symmetric monoidal categories; for this we use the work of Meseguer, Montanari, Sassone and others.


1. INTRODUCTION

Petri nets are a simple and widely studied model of computation [14, 23], with generalizations applicable to many forms of modeling [17]. Recently more attention has been paid to a compositional treatment in which Petri nets can be assembled from smaller ‘open’ Petri nets [3, 4, 5, 6, 7]. In particular, the reachability problem for Petri nets, which asks whether one marking of a Petri net can be obtained from another via a sequence of transitions, can be studied compositionally [24, 28, 30].

Here we seek to give this line of work a firmer footing in category theory. Petri nets are closely tied to symmetric monoidal categories in two ways. First, a Petri net P can be seen as a presentation


of a free symmetric monoidal category FP , with the places and transitions of P serving to freely generate the objects and morphisms of FP . We show how to construct this in Section 2, after reviewing a line of previous work going back to Meseguer and Montanari [22]. In these terms, the reachability problem asks whether there is a morphism from one object of FP to another.

Second, there is a symmetric monoidal category where the objects are sets and the morphisms are equivalence classes of open Petri nets. We construct this in Section 3, but the basic idea is very simple. Here is an open Petri net P from a set X to a set Y :




The yellow circles are places and the blue rectangle is a transition. The bold arrows from places to transitions and from transitions to places complete the structure of a Petri net. There are also arbitrary functions from X and Y into the set of places. These indicate points at which tokens could flow in or out, making our Petri net ‘open’. We write this open Petri net as $P: X \rightarrow Y$ for short.


Given another open Petri net $Q: Y \rightarrow Z$:



the first step in composing P and Q is to put the pictures together:



At this point, if we ignore the sets X, Y, Z , we have a new Petri net whose set of places is the disjoint union of those for P and Q . The second step is to identify a place of P with a place of Q whenever both are images of the same point in Y . We can then stop drawing everything involving Y , and get an open Petri net $QP: X \rightarrow Z$:



Formalizing this simple construction leads us into a bit of higher category theory. The process of taking the disjoint union of two sets of places and then quotienting by an equivalence relation is a pushout. Pushouts are defined only up to canonical isomorphism: for example, the place labeled C in the last diagram above could equally well have been labeled D or E . This is why to get a category, with composition strictly associative, we need to use *isomorphism classes* of open Petri nets as morphisms. There are advantages to avoiding this and working with open Petri nets themselves. If we do this, we obtain not a category but a bicategory [28].

However, this bicategory is equipped with more structure. Besides composing open Petri nets, we can also ‘tensor’ them via disjoint union: this describes Petri nets being run in parallel rather than in series. The result is a symmetric monoidal bicategory. Unfortunately, the axioms for a symmetric monoidal bicategory are cumbersome to check directly. Double categories turn out to be more convenient.

Double categories were introduced in the 1960s by Ehresmann [12, 13]. More recently they have been used to study open dynamical systems [18], open electrical circuits and chemical reaction networks [9], open discrete-time Markov chains [8], and coarse-graining for open continuous-time Markov chains [1].

A 2-morphism in a double category can be drawn as a square:


$$\begin{array}{ccc} X_1 & \xrightarrow{M} & Y_1 \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ X_2 & \xrightarrow{N} & Y_2. \end{array}$$

We call X_1, X_2, Y_1 and Y_2 ‘objects’, f and g ‘vertical 1-morphisms’, M and N ‘horizontal 1-cells’, and α a ‘2-morphism’. We can compose vertical 1-morphisms to get new vertical 1-morphisms and compose horizontal 1-cells to get new horizontal 1-cells. We can compose the 2-morphisms in two ways: horizontally and vertically. This is just a quick sketch of the ideas; for full definitions see Appendix A.


In Thm. 13 we construct a symmetric monoidal double category $\mathbb{O}pen(\mathbf{Petri})$ with:

- sets X, Y, Z, \dots as objects,
- functions $f: X \rightarrow Y$ as vertical 1-morphisms,
- open Petri nets $P: X \rightarrow Y$ as horizontal 1-cells,
- morphisms between open Petri nets as 2-morphisms.

To get a feeling for morphisms between open Petri nets, some examples may be helpful. There is a morphism from this open Petri net:



to this one:



mapping both primed and unprimed symbols to unprimed ones. This describes a process of ‘simplifying’ an open Petri net. There are also morphisms that include simple open Petri nets more complicated ones. For example, the above morphism of open Petri nets has a right inverse.

The main goal of this paper is to describe the reachability semantics for open Petri nets as a map from $\mathbf{Open}(\mathbf{Petri})$ to the double category of relations, \mathbf{Rel} , which has:

- sets X, Y, Z, \dots as objects,
- functions $f: X \rightarrow Y$ as vertical 1-morphisms,
- relations $R \subseteq X \times Y$ as horizontal 1-cells,
- squares

$$\begin{array}{ccc}
 X_1 & \xrightarrow{R \subseteq X_1 \times Y_1} & Y_1 \\
 f \downarrow & & \downarrow g \\
 X_2 & \xrightarrow{S \subseteq X_2 \times Y_2} & Y_2
 \end{array}$$

obeying $(f \times g)R \subseteq S$ as 2-morphisms.

In Petri net theory, a ‘marking’ of a set X is a finite multisubset of X : we can think of this as a way of placing finitely many tokens on the points of X . Let $\mathbb{N}[X]$ denote the set of markings of X . Given an open Petri net $P: X \rightarrow Y$, there is a ‘reachability relation’ saying when a given marking of X can be carried by a sequence of transitions in P to a given marking of Y , leaving no tokens behind. We write the reachability relation of P as

$$\blacksquare P \subseteq \mathbb{N}[X] \times \mathbb{N}[Y].$$

In Theorem 23 we show that the map sending P to $\blacksquare P$ extends to a lax double functor

$$\blacksquare: \mathbf{Open}(\mathbf{Petri}) \rightarrow \mathbf{Rel}.$$

In Theorem 24 we go further and show that this double functor is symmetric monoidal.

If the reader prefers bicategories to double categories, they may be relieved to learn that any double category \mathbb{D} gives rise to a bicategory $H(\mathbb{D})$ whose 2-morphisms are those 2-morphisms of \mathbb{D} of the form

$$\begin{array}{ccc}
 X & \xrightarrow{M} & Y \\
 1_X \downarrow & \Downarrow \alpha & \downarrow 1_Y \\
 X & \xrightarrow{N} & Y.
 \end{array}$$

Shulman has described conditions under which symmetric monoidal double categories give rise to symmetric monoidal bicategories [29], and using his work one can show that the reachability

semantics gives rise to a map between symmetric monoidal bicategories [2]. However, only the double category framework presents the reachability semantics in its full glory.

2. FROM PETRI NETS TO COMMUTATIVE MONOIDAL CATEGORIES

In this section we treat Petri nets as presentations of symmetric monoidal categories. As we shall explain, this has already been done by various authors. Unfortunately there are different notions of symmetric monoidal category, and also different notions of morphism between Petri nets, which combine to yield a confusing variety of possible approaches.

Here we take the maximally strict approach, and work with ‘commutative’ monoidal categories. These are just commutative monoid objects in \mathbf{Cat} , so their associator:

$$\alpha_{a,b,c}: (a \otimes b) \otimes c \xrightarrow{\sim} a \otimes (b \otimes c),$$

their left and right unitor:

$$\lambda_a: I \otimes a \xrightarrow{\sim} a, \quad \rho_a: a \otimes I \xrightarrow{\sim} a,$$

and even—disturbingly—their symmetry:

$$\sigma_{a,b}: a \otimes b \xrightarrow{\sim} b \otimes a$$

are all identity morphisms. The last would ordinarily be seen as ‘going too far’, since while every symmetric monoidal category is equivalent to one with trivial associator and unitors, this ceases to be true if we also require the symmetry to be trivial. However, it seems that Petri nets most naturally serve to present symmetric monoidal categories of this very strict sort. Thus, we construct a functor from the category of Petri nets to the category of commutative monoidal categories, which we call CMC:

$$\hat{F}: \mathbf{Petri} \rightarrow \mathbf{CMC}.$$

Intuitively speaking, this functor sends any Petri net P to the free commutative monoidal category on P . However, this functor appears not to be a left adjoint. We need a left adjoint to construct the double functor sending open Petri nets to ‘open Petri categories’ in Lemma 17. Thus, we corestrict \hat{F} to its image, which we call $\mathbf{PetriCat}$. We call a commutative monoidal category in this image a ‘Petri category’. This gives a functor sending Petri nets to Petri categories,

$$F: \mathbf{Petri} \rightarrow \mathbf{PetriCat}.$$

In fact F is an equivalence! It is thus the left adjoint of a functor

$$U: \mathbf{PetriCat} \rightarrow \mathbf{Petri}.$$

It seems Montanari and Meseguer were the first to treat Petri nets as presentations of commutative monoidal categories [22]. They constructed a closely related but different left adjoint functor from a category of Petri nets to a category of Petri categories. Our category \mathbf{Petri} is a subcategory of their category of Petri nets: our morphisms of Petri nets send places to places, while they allow more general maps that send a place to a formal linear combination of places. Our $\mathbf{PetriCat}$ is also a subcategory of their category of Petri categories, which they call $\mathbf{CatPetri}$. $\mathbf{CatPetri}$ is the full subcategory of CMC containing only commutative monoidal categories whose objects form a *free* commutative monoid. $\mathbf{PetriCat}$ is further restricted, as explained in Def. 9.

Since our adjunction

$$\mathbf{Petri} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} \mathbf{PetriCat}$$

is a restriction of Montanari and Meseguer’s, why do we not use theirs? The reason is that `PetriCat` has pushouts, while their category `CatPetri` does not. We need pushouts to construct a double category of open Petri categories in Thm. 16.

In short, the situation is surprisingly subtle given the elementary nature of the concepts involved. The paper by Montanari and Meseguer actually discusses over half a dozen categories of Petri nets and commutative monoidal categories, none of which provides an adjunction with the properties we need. Further work by Degano, Meseguer, Montanari [11], Sassone [25, 26, 27], and Sassone and Sobociński [28] explores other variations on these themes. Resisting the temptation to dwell on the subtleties of this topic, we present our approach with no further ado.

Definition 1. Let `CommMon` be the category of commutative monoids and monoid homomorphisms.

Definition 2. Let $J: \mathbf{Set} \rightarrow \mathbf{CommMon}$ be the free commutative monoid functor, that is, the left adjoint of the functor $K: \mathbf{CommMon} \rightarrow \mathbf{Set}$ that sends commutative monoids to their underlying sets and monoid homomorphisms to their underlying functions. Let

$$\mathbb{N}: \mathbf{Set} \rightarrow \mathbf{Set}$$

be the free commutative monoid monad given by the composite KJ .

For any set X , $\mathbb{N}[X]$ is the set of formal finite linear combinations of elements of X with natural number coefficients. The set X naturally includes in $\mathbb{N}[X]$, and for any function $f: X \rightarrow Y$, $\mathbb{N}[f]: \mathbb{N}[X] \rightarrow \mathbb{N}[Y]$ is given by the unique monoid homomorphism that extends f .

Definition 3. We define a **Petri net** to be a pair of functions of the following form:

$$T \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} \mathbb{N}[S].$$

We call T the set of **transitions**, S the set of **places**, s the **source** function and t the **target** function.

Definition 4. A **Petri net morphism** from the Petri net $s, t: T \rightarrow \mathbb{N}[S]$ to the Petri net $s', t': T' \rightarrow \mathbb{N}[S']$ is a pair of functions $(f: T \rightarrow T', g: S \rightarrow S')$ such that the following diagrams commute:

$$\begin{array}{ccc} T & \xrightarrow{s} & \mathbb{N}[S] \\ f \downarrow & & \downarrow \mathbb{N}[g] \\ T' & \xrightarrow{s'} & \mathbb{N}[S'] \end{array} \quad \begin{array}{ccc} T & \xrightarrow{t} & \mathbb{N}[S] \\ f \downarrow & & \downarrow \mathbb{N}[g] \\ T' & \xrightarrow{t'} & \mathbb{N}[S'] \end{array}$$

Definition 5. Let `Petri` be the category of Petri nets and Petri net morphisms, with composition defined by

$$(f, g) \circ (f', g') = (f \circ f', g \circ g').$$

Our definition of Petri net morphism differs from the earlier definition used by Sassone [25, 26, 27] and Degano–Meseguer–Montanari [11]. The difference is that our definition requires that the homomorphism between free commutative monoids come from a function between the sets of places.

Definition 6. A **commutative monoidal category** is a commutative monoid object internal to `Cat`. Explicitly, a commutative monoidal category is a strict monoidal category (C, \otimes, I) such that for all objects a and b and morphisms f and g in C

$$a \otimes b = b \otimes a \text{ and } f \otimes g = g \otimes f.$$

Note that a commutative monoidal category is the same as a strict symmetric monoidal category where the symmetry isomorphisms $\sigma_{a,b}: a \otimes b \xrightarrow{\sim} b \otimes a$ are all identity morphisms. Every strict monoidal functor between commutative monoidal categories is automatically a strict symmetric monoidal functor. This motivates the following definition:

Definition 7. Let **CMC** be the category whose objects are commutative monoidal categories and whose morphisms are strict monoidal functors.

We can turn a Petri net $P = (s, t: T \rightarrow \mathbb{N}[S])$ into a commutative monoidal category $\hat{F}P$. We take the commutative monoid of objects $\text{Ob}(\hat{F}P)$ to be the free commutative monoid on S . We construct the commutative monoid of morphisms $\text{Mor}(\hat{F}P)$ as follows. First we generate morphisms recursively:

- for every transition $\tau \in T$ we include a morphism $\tau: s(\tau) \rightarrow t(\tau)$;
- for any object a we include a morphism $1_a: a \rightarrow a$;
- for any morphisms $f: a \rightarrow b$ and $g: a' \rightarrow b'$ we include a morphism denoted $f + g: a + a' \rightarrow b + b'$ to serve as their tensor product;
- for any morphisms $f: a \rightarrow b$ and $g: b \rightarrow c$ we include a morphism $g \circ f: a \rightarrow c$ to serve as their composite.

Then we mod out by an equivalence relation on morphisms that imposes the laws of a commutative monoidal category, obtaining the commutative monoid $\text{Mor}(\hat{F}P)$.

Definition 8. Let $\hat{F}: \text{Petri} \rightarrow \text{CMC}$ be the functor that makes the following assignments on Petri nets and morphisms:

$$\begin{array}{ccc} T & \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{t} \end{array} & \mathbb{N}[S] \\ \downarrow f & & \downarrow \mathbb{N}[g] \\ T' & \begin{array}{c} \xrightarrow{s'} \\ \xrightarrow{t'} \end{array} & \mathbb{N}[S'] \end{array} \mapsto \begin{array}{ccc} \hat{F}P & & \\ \downarrow \hat{F}(f,g) & & \\ \hat{F}P' & & \end{array}$$

Here $\hat{F}(f, g): \hat{F}P \rightarrow \hat{F}P'$ is defined on objects by $\mathbb{N}[g]$. On morphisms, $\hat{F}(f, g)$ is the unique map extending f that preserves identities, composition, and the tensor product.

As far as we can tell, \hat{F} is not a left adjoint. To get an adjunction we corestrict \hat{F} to its image.

Definition 9. Let **PetriCat** be the image of \hat{F} . We call an object of **PetriCat** a **Petri category**.

Explicitly, an object of **PetriCat** is a commutative monoidal category of the form $\hat{F}P$ for some Petri net P , and a morphism in **PetriCat** from $\hat{F}P$ to $\hat{F}P'$ is a strict monoidal functor of the form $\hat{F}(f, g): \hat{F}P \rightarrow \hat{F}P'$ for some morphism of Petri nets $(f, g): P \rightarrow P'$.

Lemma 10. *Let*

$$F: \text{Petri} \rightarrow \text{PetriCat}$$

*be the corestriction of \hat{F} to **PetriCat**. Then F is an equivalence of categories, and thus a left adjoint.*

Proof. Since \hat{F} is essentially surjective and full by construction, it suffices to show that \hat{F} is faithful. Indeed, if $\hat{F}((f, g)): \hat{F}(P) \rightarrow \hat{F}(Q)$ and $F((f', g')): \hat{F}(P) \rightarrow \hat{F}(Q)$ are the same commutative monoidal functor, then we have $\mathbb{N}[f] = \mathbb{N}[f']$ and $\mathbb{N}[g] = \mathbb{N}[g']$. This implies that $f = f'$ and $g = g'$, so (f, g) and (f', g') are the same morphism in **Petri**. \square

3. OPEN PETRI NETS

Our goal in this paper is to use the language of double categories to develop a theory of Petri nets with inputs and outputs that can be glued together. The first step is to construct a double category $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$ whose horizontal 1-morphisms are open Petri nets. For this we need a functor $L: \mathbf{Set} \rightarrow \mathbf{Petri}$ that maps any set S to a Petri net with S as its set of places, and we need L to be a left adjoint.

Definition 11. Let $L: \mathbf{Set} \rightarrow \mathbf{Petri}$ be the functor defined on sets and functions as follows:

$$\begin{array}{ccc} X & & \emptyset \rightrightarrows \mathbb{N}[X] \\ f \downarrow & \mapsto & \downarrow \qquad \downarrow \mathbb{N}[f] \\ Y & & \emptyset \rightrightarrows \mathbb{N}[Y] \end{array}$$

where the unlabeled maps are the unique maps of that type.

Lemma 12. *The functor L has a right adjoint $R: \mathbf{Petri} \rightarrow \mathbf{Set}$ that acts as follows on Petri nets and Petri net morphisms:*

$$\begin{array}{ccc} T \xrightarrow{s} \mathbb{N}[S] & & S \\ f \downarrow & \mapsto & \downarrow g \\ T' \xrightarrow{s'} \mathbb{N}[S] & & S' \\ & & \downarrow t' \end{array}$$

Proof. For any set X and Petri net $P = (s, t: T \rightarrow \mathbb{N}[S])$ we have natural isomorphisms

$$\begin{aligned} \mathrm{hom}_{\mathbf{Petri}}(L(X), T \xrightarrow{s} \mathbb{N}[S]) &\cong \mathrm{hom}_{\mathbf{Petri}}(\emptyset \rightrightarrows \mathbb{N}[X], T \xrightarrow{s} \mathbb{N}[S]) \\ &\cong \mathrm{hom}_{\mathbf{Set}}(X, S) \\ &\cong \mathrm{hom}_{\mathbf{Set}}(X, R(T \xrightarrow{s} \mathbb{N}[S])). \quad \square \end{aligned}$$

We now introduce the main object of study: the double category $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$. Since this is a symmetric monoidal double category, it involves quite a lot of structure. The definition of symmetric monoidal double category can be found in Appendix A.

Theorem 13. *There is a symmetric monoidal double category $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$ for which:*

- objects are sets
- vertical 1-morphisms are functions
- horizontal 1-cells from a set X to a set Y are **open Petri nets** $P: X \rightarrow Y$, that is, cospans in \mathbf{Petri} of the form

$$\begin{array}{ccc} & P & \\ i \nearrow & & \nwarrow o \\ LX & & LY \end{array}$$

- 2-morphisms $\alpha: P \Rightarrow P'$ are commutative diagrams

$$\begin{array}{ccccc} LX & \xrightarrow{i} & P & \xleftarrow{o} & LY \\ Lf \downarrow & & \alpha \downarrow & & \downarrow Lg \\ LX' & \xrightarrow{i'} & P' & \xleftarrow{o'} & LY' \end{array}$$

in Petri.

Composition of vertical 1-morphisms is the usual composition of functions. Composition of horizontal 1-cells is composition of cospans via pushout: given two horizontal 1-cells

$$\begin{array}{ccc} & P & \\ i_1 \nearrow & & \nwarrow o_1 \\ LX & & LY \end{array} \quad \begin{array}{ccc} & Q & \\ i_2 \nearrow & & \nwarrow o_2 \\ LY & & LZ \end{array}$$

their composite is given by this cospan from LX to LZ :

$$\begin{array}{ccccc} & & P +_{LY} Q & & \\ & j_P \nearrow & & \nwarrow j_Q & \\ & P & & Q & \\ i_1 \nearrow & & & & \nwarrow o_2 \\ LX & & LY & & LZ \end{array}$$

where the diamond is a pushout square. The horizontal composite of 2-morphisms

$$\begin{array}{ccccc} LX & \xrightarrow{i_1} & P & \xleftarrow{o_1} & LY \\ Lf \downarrow & & \alpha \downarrow & & \downarrow Lg \\ LX' & \xrightarrow{i'_1} & P' & \xleftarrow{o'_1} & LY' \end{array} \quad \begin{array}{ccccc} LY & \xrightarrow{i_2} & Q & \xleftarrow{o_2} & LZ \\ Lg \downarrow & & \beta \downarrow & & \downarrow Lh \\ LY' & \xrightarrow{i'_2} & Q' & \xleftarrow{o'_2} & LZ' \end{array}$$

is given by

$$\begin{array}{ccccc} LX & \xrightarrow{j_P i_1} & P +_{LY} Q & \xleftarrow{j_Q o_1} & LZ \\ Lf \downarrow & & \alpha +_{Lg} \beta \downarrow & & \downarrow Lh \\ LX' & \xrightarrow{j_{P'} i'_1} & P' +_{LY'} Q' & \xleftarrow{j_{Q'} o'_2} & LZ' \end{array}$$

Vertical composition of 2-morphisms is done using composition of functions. The symmetric monoidal structure comes from coproducts in \mathbf{Set} and \mathbf{Petri} .

Proof. We construct this symmetric monoidal double category using the machinery of ‘structured cospans’ [2]. The main tool is the following lemma, which explains the symmetric monoidal structure in more detail:

Lemma 14. *Let \mathbf{A} be a category with finite coproducts and \mathbf{X} be a category with finite colimits. Given a left adjoint $L: \mathbf{A} \rightarrow \mathbf{X}$, there exists a unique symmetric monoidal double category ${}_L\mathbf{Csp}(\mathbf{X})$, such that:*

- objects are objects of \mathbf{A} ,

- vertical 1-morphisms are morphisms of \mathbf{A} ,
- a horizontal 1-cell from $a \in \mathbf{A}$ to $b \in \mathbf{A}$ is a cospan in \mathbf{X} of this form:

$$La \longrightarrow x \longleftarrow Lb$$

- a 2-morphism is a commutative diagram in \mathbf{X} of this form:

$$\begin{array}{ccccc} La & \longrightarrow & x & \longleftarrow & Lb \\ Lf \downarrow & & h \downarrow & & \downarrow Lg \\ Lc & \longrightarrow & y & \longleftarrow & Ld. \end{array}$$

Composition of vertical 1-morphisms is composition in \mathbf{A} . Composition of horizontal 1-cells is composition of cospans in \mathbf{X} via pushout: given horizontal 1-cells

$$\begin{array}{ccc} & x & \\ i_1 \nearrow & & \nwarrow o_1 \\ La & & Lb \end{array} \quad \begin{array}{ccc} & y & \\ i_2 \nearrow & & \nwarrow o_2 \\ Lb & & Lc \end{array}$$

their composite is this cospan from La to Lc :

$$\begin{array}{ccc} & x +_{Lb} y & \\ j_x \nearrow & & \nwarrow j_y \\ & x & y \\ i_1 \nearrow & & \nwarrow o_1 \quad i_2 \nearrow & & \nwarrow o_2 \\ La & & Lb & & Lc \end{array}$$

where the diamond is a pushout square. The horizontal composite of 2-morphisms

$$\begin{array}{ccc} La \xrightarrow{i_1} x \xleftarrow{o_1} Lb & & Lb \xrightarrow{i_2} y \xleftarrow{o_2} Lc \\ Lf \downarrow & \alpha \downarrow & \downarrow Lg \\ La' \xrightarrow{i'_1} x' \xleftarrow{o'_1} Lb' & & Lb' \xrightarrow{i'_2} y' \xleftarrow{o'_2} Lc' \\ Lg \downarrow & \beta \downarrow & \downarrow Lh \end{array}$$

is given by

$$\begin{array}{ccc} La \xrightarrow{j_x i_1} x +_{Lb} y \xleftarrow{j_y o_2} Lc & & \\ Lf \downarrow & \alpha +_{Lg} \beta \downarrow & \downarrow Lh \\ La' \xrightarrow{j_{x'} i'_1} x' +_{Lb'} y' \xleftarrow{j_{y'} o'_2} Lc' & & \end{array}$$

The vertical composite of 2-morphisms

$$\begin{array}{ccc} La \xrightarrow{i_1} x \xleftarrow{o_1} Lb & & \\ Lf \downarrow & \alpha \downarrow & \downarrow Lg \\ La' \xrightarrow{i'_1} x' \xleftarrow{o'_1} Lb' & & \end{array}$$

$$\begin{array}{ccccc}
 La' & \xrightarrow{i'_1} & x' & \xleftarrow{o'_1} & Lb' \\
 Lf' \downarrow & & \alpha' \downarrow & & \downarrow Lg' \\
 La'' & \xrightarrow{i''_1} & x'' & \xleftarrow{o''_1} & Lb''
 \end{array}$$

is given by

$$\begin{array}{ccccc}
 La & \xrightarrow{i_1} & x & \xleftarrow{o_1} & Lb \\
 L(f'f) \downarrow & & \alpha'\alpha \downarrow & & \downarrow L(g'g) \\
 La'' & \xrightarrow{i''_1} & x'' & \xleftarrow{o''_1} & Lb''.
 \end{array}$$

The tensor product is defined using chosen coproducts in \mathbf{A} and \mathbf{X} . Thus, the tensor product of two objects a_1 and a_2 is $a_1 + a_2$, the tensor product of two vertical 1-morphisms

$$\begin{array}{ccc}
 a_1 & & a_2 \\
 f_1 \downarrow & & f_2 \downarrow \\
 b_1 & & b_2
 \end{array}$$

is

$$\begin{array}{c}
 a_1 + a_2 \\
 f_1 + f_2 \downarrow \\
 b_1 + b_2,
 \end{array}$$

the tensor product of two horizontal 1-cells

$$La_1 \xrightarrow{i_1} x_1 \xleftarrow{o_1} Lb_1 \qquad La_2 \xrightarrow{i_2} x_2 \xleftarrow{o_2} Lb_2$$

is

$$L(a_1 + a_2) \xrightarrow{i_1+i_2} x_1 + x_2 \xleftarrow{o_1+o_2} L(b_1 + b_2),$$

and the tensor product of two 2-morphisms

$$\begin{array}{ccc}
 La_1 \xrightarrow{i_1} x_1 \xleftarrow{o_1} Lb_1 & & La_2 \xrightarrow{i_2} x_2 \xleftarrow{o_2} Lb_2 \\
 Lf_1 \downarrow & \alpha_1 \downarrow & \downarrow Lg_1 \\
 La'_1 \xrightarrow{i'_1} x'_1 \xleftarrow{o'_1} Lb'_1 & & La'_2 \xrightarrow{i'_2} x'_2 \xleftarrow{o'_2} Lb'_2 \\
 Lf_2 \downarrow & \alpha_2 \downarrow & \downarrow Lg_2
 \end{array}$$

is

$$\begin{array}{ccc}
 L(a_1 + a_2) \xrightarrow{i_1+i_2} x_1 + x_2 \xleftarrow{o_1+o_2} L(b_1 + b_2) & & \\
 L(f_1+f_2) \downarrow & \alpha_1+\alpha_2 \downarrow & \downarrow L(g_1+g_2) \\
 L(a'_1 + a'_2) \xrightarrow{i'_1+i'_2} x'_1 + x'_2 \xleftarrow{o'_1+o'_2} L(b'_1 + b'_2). & &
 \end{array}$$

The units for these tensor products are taken to be initial objects, and the symmetry is defined using the canonical isomorphisms $a + b \cong b + a$.

Proof. This is [2, Cor. 3.10]. Note that we are abusing language slightly above. We must choose a specific coproduct for each pair of objects in \mathbf{X} and \mathbf{A} to give ${}_L\mathbf{Csp}(X)$ its tensor product. Given morphisms $i_1: La_1 \rightarrow x_1$ and $i_2: La_2 \rightarrow x_2$, their coproduct is really a morphism $i_1 + i_2: La_1 + La_2 \rightarrow x_1 + x_2$ between these chosen coproducts. But since L preserves coproducts, we can compose this morphism with the canonical isomorphism $L(a_1 + a_2) \cong La_1 + La_2$ to obtain the morphism that we call $i_1 + i_2: L(a_1 + a_2) \rightarrow x_1 + x_2$ above. \square

To apply this lemma to the situation at hand we need the following result.

Lemma 15. *Petri has small colimits.*

Proof. A small diagram $D: \mathbf{J} \rightarrow \mathbf{Petri}$ induces small diagrams $D_T: \mathbf{J} \rightarrow \mathbf{Set}$ on the transitions and $D_S: \mathbf{J} \rightarrow \mathbf{Set}$ on the places. We compute colimits in \mathbf{Petri} pointwise, so $\text{colim } D$ consists of the Petri net

$$\text{colim } D_T \xrightarrow[t]{s} \mathbb{N}[\text{colim } D_S]$$

made into a cocone using these morphisms of Petri nets for each $j \in \mathbf{J}$:

$$\begin{array}{ccc} D_T(j) & \xrightarrow[t_j]{s_j} & \mathbb{N}[D_S(j)] \\ f_j \downarrow & & \downarrow \mathbb{N}[g_j] \\ \text{colim } D_T & \xrightarrow[\tau]{\sigma} & \mathbb{N}[\text{colim } D_S]. \end{array}$$

All the above colimits are taken in \mathbf{Set} , and σ and τ are the unique maps induced by the universal property of $\text{colim } D_T$.

Given a Petri net

$$P = \left(T \xrightarrow[t]{s} \mathbb{N}[S] \right)$$

and a family of Petri net morphisms $\{(h_j, k_j): D(j) \rightarrow P\}$, we can prove there exists a unique morphism of Petri nets filling in the dashed arrows here:

$$\begin{array}{ccc} D_T(j) & \xrightarrow[t_j]{s_j} & \mathbb{N}[D_S(j)] \\ \downarrow h_j & & \downarrow \mathbb{N}[k_j] \\ T & \xrightarrow[t]{s} & \mathbb{N}[S] \\ \uparrow \text{dashed} & & \uparrow \text{dashed} \\ \text{colim } D_T & \xrightarrow[\tau]{\sigma} & \mathbb{N}[\text{colim } D_S] \end{array}$$

(Left dashed arrow is labeled f_j , right dashed arrow is labeled $\mathbb{N}[g_j]$)

The left dashed arrow arises from the universal property of $\text{colim } D_T$, while the right one is obtained by applying \mathbb{N} to the arrow arising from the universal property of $\text{colim } D_S$. \square

We now have all of the ingredients to apply Lemma 14 to the functor $L: \mathbf{Set} \rightarrow \mathbf{Petri}$. Thm. 13 follows from realizing that $\mathbf{Open}(\mathbf{Petri})$ as described in the theorem is the symmetric monoidal double category ${}_L\mathbf{Csp}(\mathbf{Petri})$. \square

4. OPEN PETRI CATEGORIES

In Section 2 we saw how a Petri net P gives a commutative monoidal category FP , and in Section 3 we constructed a double category $\mathbb{O}\mathbf{pen}(\mathbf{Petri})$ of open Petri nets. Now we construct a double category $\mathbb{O}\mathbf{pen}(\mathbf{PetriCat})$ of ‘open Petri categories’ and a map

$$\mathbf{Csp}(F): \mathbb{O}\mathbf{pen}(\mathbf{Petri}) \rightarrow \mathbb{O}\mathbf{pen}(\mathbf{PetriCat}).$$

This can be seen as providing a functorial semantics for open Petri nets in which any open Petri net is mapped to the commutative monoidal category it presents. The reachability semantics for open Petri nets is based on this more fundamental form of semantics.

The key is this commutative diagram of left adjoint functors:

$$\begin{array}{ccc} \mathbf{Set} & \xrightarrow{L} & \mathbf{Petri} \\ & \searrow^{L'} & \downarrow F \\ & & \mathbf{PetriCat} \end{array}$$

where $L' = FL$ sends any set to the free Petri category on this set: $L'X$ has $\mathbb{N}[X]$ as its set of objects, and only identity morphisms. Using Lemma 14, we can produce two symmetric monoidal double categories from this diagram. We have already seen one: $\mathbb{O}\mathbf{pen}(\mathbf{Petri}) = {}_L\mathbf{Csp}(\mathbf{Petri})$. We now introduce the other: $\mathbb{O}\mathbf{pen}(\mathbf{PetriCat}) = {}_{L'}\mathbf{Csp}(\mathbf{PetriCat})$.

Theorem 16. *There is a symmetric monoidal double category $\mathbb{O}\mathbf{pen}(\mathbf{PetriCat})$ for which:*

- *objects are sets*
- *vertical 1-morphisms are functions*
- *horizontal 1-cells from a set X to a set Y are open Petri categories $C: X \rightarrow Y$, that is, cospans in $\mathbf{PetriCat}$ of the form*

$$\begin{array}{ccc} & C & \\ i \nearrow & & \nwarrow o \\ L'X & & L'Y \end{array}$$

where C is a Petri category and i, o are strict monoidal functors,

- *2-morphisms $\alpha: C \Rightarrow C'$ are commutative diagrams in $\mathbf{PetriCat}$ of the form*

$$\begin{array}{ccccc} L'X & \xrightarrow{i} & C & \xleftarrow{o} & L'Y \\ L'f \downarrow & & \alpha \downarrow & & \downarrow L'g \\ L'X' & \xrightarrow{i} & C' & \xleftarrow{o'} & L'Y' \end{array}$$

and the rest of the structure is given as in Lemma 14.

Proof. To apply Lemma 14 to the functor $L': \mathbf{Set} \rightarrow \mathbf{PetriCat}$ we just need to check that $\mathbf{PetriCat}$ has finite colimits. In Lemma 10 we saw

$$F: \mathbf{Petri} \xrightarrow{\sim} \mathbf{PetriCat}$$

was an equivalence. Thus, colimits in $\mathbf{PetriCat}$ are given by applying F to the colimits in \mathbf{Petri} . \square

The functor $F: \mathbf{Petri} \rightarrow \mathbf{PetriCat}$ induces a map sending open Petri nets to open Petri categories. This map is actually part of a ‘symmetric monoidal double functor’, a concept recalled in Appendix A.

Lemma 17. *There is a symmetric monoidal double functor*

$$\mathbb{Csp}(F): \mathbb{Open}(\text{Petri}) \rightarrow \mathbb{Open}(\text{PetriCat})$$

that is the identity on objects and vertical 1-morphisms, and makes the following assignments on horizontal 1-cells and 2-morphisms:

$$\begin{array}{ccc} LX \xrightarrow{i} P \xleftarrow{o} LY & & L'X \xrightarrow{Fi} FP \xleftarrow{Fo} L'Y \\ Lf \downarrow & \alpha \downarrow & \downarrow Lg \\ LX' \xrightarrow{i'} P' \xleftarrow{o'} LY' & \mapsto & L'X' \xrightarrow{Fi'} FP' \xleftarrow{Fo'} L'Y' \end{array}$$

Proof. This follows from the theory of structured cospans. More generally, suppose \mathbf{A} is a category with finite coproducts and \mathbf{X}, \mathbf{X}' are categories with finite colimits. Suppose there is a commuting triangle of left adjoints

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{L} & \mathbf{X} \\ & \searrow L' & \downarrow F \\ & & \mathbf{X}' \end{array}$$

Then Lemma 14 gives us symmetric monoidal double categories ${}_L\mathbb{Csp}(\mathbf{X})$ and ${}_{L'}\mathbb{Csp}(\mathbf{X}')$, and Thm. 4.2 of [2] gives a symmetric monoidal double functor

$$\mathbb{Csp}(F): {}_L\mathbb{Csp}(\mathbf{X}) \rightarrow {}_{L'}\mathbb{Csp}(\mathbf{X}')$$

that is the identity on objects and vertical morphisms, and acts as follows on horizontal 1-cells and 2-morphisms:

$$\begin{array}{ccc} La \xrightarrow{i} x \xleftarrow{o} Lb & & L'a \xrightarrow{Fi} Fx \xleftarrow{Fo} L'b \\ Lf \downarrow & \alpha \downarrow & \downarrow Lg \\ La' \xrightarrow{i'} x' \xleftarrow{o'} Lb' & \mapsto & L'a' \xrightarrow{Fi'} Fx' \xleftarrow{Fo'} L'b' \end{array}$$

□

5. THE DOUBLE CATEGORY OF RELATIONS

Using the language of functorial semantics, $\mathbb{Open}(\text{Petri})$ can be thought of as a ‘syntax’ for describing open systems, and reachability as a choice of ‘semantics’. To implement this, we show that the reachability relation of a Petri net can be defined for open Petri nets in a way that gives a lax double functor from $\mathbb{Open}(\text{Petri})$ to the double category of relations constructed by Grandis and Paré [15, Sec. 3.4]. Here we recall this double category and give it a symmetric monoidal structure.

This double category, which we call \mathbb{Rel} , has:

- sets as objects,
- functions $f: X \rightarrow Y$ as vertical 1-morphisms from X to Y ,
- relations $R \subseteq X \times Y$ as horizontal 1-cells from X to Y ,

- squares

$$\begin{array}{ccc}
 X_1 & \xrightarrow{R \subseteq X_1 \times Y_1} & Y_1 \\
 f \downarrow & & \downarrow g \\
 X_2 & \xrightarrow{S \subseteq X_2 \times Y_2} & Y_2
 \end{array}$$

obeying $(f \times g)R \subseteq S$ as 2-morphisms.

The last item deserves some explanation. A preorder is a category such that for any pair of objects a, b there exists at most one morphism $\alpha: a \rightarrow b$. When such a morphism exists we usually write $a \leq b$. Similarly there is a kind of double category for which given any ‘frame’

$$\begin{array}{ccc}
 X_1 & \xrightarrow{M} & Y_1 \\
 f \downarrow & & \downarrow g \\
 X_2 & \xrightarrow{N} & Y_2
 \end{array}$$

there exists at most one 2-morphism

$$\begin{array}{ccc}
 X_1 & \xrightarrow{M} & Y_1 \\
 f \downarrow & \Downarrow \alpha & \downarrow g \\
 X_2 & \xrightarrow{N} & Y_2
 \end{array}$$

filling this frame. Following [1] we call this a **degenerate** double category. Our definition of the 2-morphism in $\mathbb{R}el$ will imply that this double category is degenerate.

Composition of vertical 1-morphisms in $\mathbb{R}el$ is the usual composition of functions, while composition of horizontal 1-cells is the usual composition of relations. Since composition of relations obeys the associative and unit laws strictly, $\mathbb{R}el$ will be a *strict* double category. Since $\mathbb{R}el$ is degenerate, there is at most one way to define the vertical composite of 2-morphisms

$$\begin{array}{ccc}
 X_1 & \xrightarrow{R \subseteq X_1 \times Y_1} & Y_1 \\
 f \downarrow & \Downarrow \alpha & \downarrow g \\
 X_2 & \xrightarrow{S \subseteq X_2 \times Y_2} & Y_2 \\
 f' \downarrow & \Downarrow \beta & \downarrow g' \\
 X_3 & \xrightarrow{T \subseteq X_3 \times Y_3} & Y_3
 \end{array}
 =
 \begin{array}{ccc}
 X_1 & \xrightarrow{R \subseteq X_1 \times Y_1} & Y_1 \\
 f'f \downarrow & \Downarrow \beta\alpha & \downarrow g'g \\
 X_3 & \xrightarrow{T \subseteq X_3 \times Y_3} & Y_3
 \end{array}$$

so we need merely check that a 2-morphism $\beta\alpha$ filling the frame at right exists. This amounts to noting that

$$(f \times g)R \subseteq S, (f' \times g')S \subseteq T \implies (f' \times g')(f \times g)R \subseteq T.$$

Similarly, there is at most one way to define the horizontal composite of 2-morphisms

$$\begin{array}{ccccc} X_1 & \xrightarrow{R \subseteq X_1 \times Y_1} & Y_1 & \xrightarrow{R' \subseteq Y_1 \times Z_1} & Z_1 \\ \downarrow f & & \downarrow g & & \downarrow h \\ X_2 & \xrightarrow{S \subseteq X_2 \times Y_2} & Y_2 & \xrightarrow{S' \subseteq Y_2 \times Z_2} & Z_2 \end{array} \quad = \quad \begin{array}{ccc} X_1 & \xrightarrow{R'R \subseteq X_1 \times Z_1} & Z_1 \\ \downarrow f & & \downarrow h \\ X_2 & \xrightarrow{S'S \subseteq X_2 \times Z_2} & Z_2 \end{array}$$

so we need merely check that a filler $\alpha' \circ \alpha$ exists, which amounts to noting that

$$(f \times g)R \subseteq S, (g \times h)R' \subseteq S' \implies (f \times h)(R'R) \subseteq S'S.$$

Theorem 18. *There exists a strict double category $\mathbb{R}el$ with the above properties.*

Proof. We use the definition of double category in Appendix A (Def. 25), which introduces two concepts not mentioned so far: the category of objects and the category of arrows. We define the category of objects $\mathbb{R}el_0$ to have sets as objects and functions as morphisms. We define the category of arrows $\mathbb{R}el_1$ to have relations as objects and squares

$$\begin{array}{ccc} X_1 & \xrightarrow{R \subseteq X_1 \times X_2} & X_2 \\ \downarrow f & & \downarrow g \\ Y_1 & \xrightarrow{S \subseteq Y_1 \times Y_2} & Y_2 \end{array}$$

with $(f \times g)R \subseteq S$ as morphisms. The source and target functors $S, T: \mathbb{R}el_1 \rightarrow \mathbb{R}el_0$ are clear. The identity-assigning functor $u: \mathbb{R}el_0 \rightarrow \mathbb{R}el_1$ sends a set X to the identity function 1_X and a function $f: X \rightarrow Y$ to the unique 2-morphism

$$\begin{array}{ccc} X & \xrightarrow{1_X} & X \\ \downarrow f & & \downarrow f \\ Y & \xrightarrow{1_Y} & Y \end{array}$$

The composition functor $\odot: \mathbb{R}el_1 \times_{\mathbb{R}el_0} \mathbb{R}el_1 \rightarrow \mathbb{R}el_1$ acts on objects by the usual composition of relations, and it acts on 2-morphisms by horizontal composition as described above. These functors can be shown to obey all the axioms of a double category. In particular, because $\mathbb{R}el$ is degenerate, all the required equations between 2-morphisms, such as the interchange law, hold automatically. \square

Next we make $\mathbb{R}\mathbf{el}$ into a symmetric monoidal double category. To do this, we first give $\mathbb{R}\mathbf{el}_0 = \mathbf{Set}$ the symmetric monoidal structure induced by the cartesian product. Then we give $\mathbb{R}\mathbf{el}_1$ a symmetric monoidal structure as follows. Given relations $R_1 \subseteq X_1 \times Y_1$ and $R_2 \subseteq X_2 \times Y_2$, we define

$$R_1 \times R_2 = \{(x_1, x_2, y_1, y_2) : (x_1, y_1) \in R_1, (x_2, y_2) \in R_2\} \subseteq X_1 \times X_2 \times Y_1 \times Y_2.$$

Given two 2-morphisms in $\mathbb{R}\mathbf{el}_1$:

$$\begin{array}{ccc} X_1 & \xrightarrow{R \subseteq X_1 \times Y_1} & Y_1 \\ \downarrow f & \Downarrow \alpha & \downarrow g \\ X_2 & \xrightarrow{S \subseteq X_2 \times Y_2} & Y_2 \end{array} \quad \begin{array}{ccc} X'_1 & \xrightarrow{R' \subseteq X'_1 \times Y'_1} & Y'_1 \\ \downarrow f' & \Downarrow \alpha' & \downarrow g' \\ X'_2 & \xrightarrow{S' \subseteq X'_2 \times Y'_2} & Y'_2 \end{array}$$

there is at most one way to define their product

$$\begin{array}{ccc} X_1 \times X'_1 & \xrightarrow{R \times R' \subseteq (X_1 \times X'_1) \times (Y_1 \times Y'_1)} & Y_1 \times Y'_1 \\ \downarrow f \times f' & \Downarrow \alpha \times \alpha' & \downarrow g \times g' \\ X_2 \times X'_2 & \xrightarrow{S \times S' \subseteq (X_2 \times X'_2) \times (Y_2 \times Y'_2)} & Y_2 \times Y'_2 \end{array}$$

because $\mathbb{R}\mathbf{el}$ is degenerate. To show that $\alpha \times \alpha'$ exists, we need merely note that

$$(f \times g)R \subseteq S, (f' \times g')R' \subseteq S' \implies (f \times f' \times g \times g')(R \times R') \subseteq S \times S'.$$

Theorem 19. *The double category $\mathbb{R}\mathbf{el}$ can be given the structure of a symmetric monoidal double category with the above properties.*

Proof. We have described $\mathbb{R}\mathbf{el}_0$ and $\mathbb{R}\mathbf{el}_1$ as symmetric monoidal categories. The source and target functors $S, T: \mathbb{R}\mathbf{el}_1 \rightarrow \mathbb{R}\mathbf{el}_0$ are strict symmetric monoidal functors. We must also equip $\mathbb{R}\mathbf{el}$ with two other pieces of structure. One, called χ , says how the composition of horizontal 1-cells interacts with the tensor product in the category of arrows. The other, called μ , says how the identity-assigning functor u relates the tensor product in the category of objects to the tensor product in the category of arrows. These are defined as follows. Given four horizontal 1-cells

$$\begin{aligned} R_1 &\subseteq X_1 \times Y_1, & R_2 &\subseteq Y_1 \times Z_1, \\ S_1 &\subseteq X_2 \times Y_2, & S_2 &\subseteq Y_2 \times Z_2, \end{aligned}$$

the globular 2-isomorphism $\chi: (R_2 \times S_2)(R_1 \times S_1) \Rightarrow (R_2 R_1) \times (S_2 S_1)$ is the identity 2-morphism

$$\begin{array}{ccc} X_1 \times X_2 & \xrightarrow{(R_2 \times S_2)(R_1 \times S_1)} & Z_1 \times Z_2 \\ \downarrow 1 & & \downarrow 1 \\ X_1 \times X_2 & \xrightarrow{(R_2 R_1) \times (S_2 S_1)} & Z_1 \times Z_2 \end{array}$$

The globular 2-isomorphism $\mu: u(X \times Y) \Rightarrow u(X) \times u(Y)$ is the identity 2-morphism

$$\begin{array}{ccc} X \times Y & \xrightarrow{1_{X \times Y}} & X \times Y \\ \downarrow 1 & & \downarrow 1 \\ X \times Y & \xrightarrow{1_X \times 1_Y} & X \times Y \end{array}$$

All the commutative diagrams in the definition of symmetric monoidal double category (Defs. 29 and 30) can be checked straightforwardly. In particular, all diagrams of 2-morphisms commute automatically because $\mathbb{R}el$ is degenerate. \square

6. THE REACHABILITY SEMANTICS

Now we explain how $\mathbb{O}pen(\text{Petri})$ provides a compositional approach to the reachability problem. In particular, we prove that the reachability semantics defines a lax double functor

$$\blacksquare: \mathbb{O}pen(\text{Petri}) \rightarrow \mathbb{R}el$$

which is symmetric monoidal.

Definition 20. Let P be a Petri net $(s, t: T \rightarrow \mathbb{N}[S])$. A **marking** of P is an element $m \in \mathbb{N}[S]$. Given a transition $\tau \in T$, a **firing** of τ is a tuple (τ, m, n) such that $n = m - s(\tau) + t(\tau)$. We say that a marking n is **reachable** from a marking m if for some $k \geq 1$ there is a sequence of markings $m = m_1, \dots, m_k = n$ and firings $\{(\tau_i, m_i, m_{i+1})\}_{i=1}^{k-1}$. In particular, taking $k = 1$, any marking is reachable from itself with no firings.

Given two markings of a Petri net, the problem of deciding whether one is reachable from the other is called the ‘reachability problem’. In 1984 Mayr showed that the reachability problem is decidable [21]. However, it is a very hard problem: in 1976 Lipton had showed that it requires at least exponential space, and in fact any EXPSPACE algorithm can be reduced in polynomial time to a Petri net reachability problem [19].

There is a close connection between reachability and the free commutative monoidal category on a Petri net constructed in Lemma 10.

Proposition 21. *If m and n are markings of a Petri net P , then n is reachable from m if and only if there is a morphism $f: m \rightarrow n \in FP$.*

Proof. If n is reachable from m , there is a sequence of markings $m = m_1, \dots, m_k = n$ and firings $\{(\tau_i, m_i, m_{i+1})\}_{i=1}^{k-1}$. For each firing (τ_i, m_i, m_{i+1}) there is a morphism in FP given by

$$\tau_i + 1_{m_i - s(\tau_i)}: m_i \rightarrow m_{i+1}.$$

Taking the composite of these morphisms gives a morphism $f: m \rightarrow n$ in FP .

Conversely, if $f: m \rightarrow n$ is a morphism in FP , it can be obtained by composition and addition (that is, the tensor product) from morphisms arising from the basic transitions and symmetry morphisms. Because $+$ is a functor, we have the interchange law

$$(f_1 \circ g_1) + (f_2 \circ g_2) = (f_1 + f_2) \circ (g_1 + g_2)$$

whenever f_1, g_1 and f_2, g_2 are pairs of composable morphisms in FP . We can use this inductively to simplify f into a composite of sums. If $f_1: a_1 \rightarrow b_1$ and $f_2: a_2 \rightarrow b_2$ are morphisms in FP , the interchange law also tells us that

$$f_1 + f_2 = (f_1 \circ 1_{a_1}) + (1_{b_2} \circ f_2) = (f_1 + 1_{b_2}) \circ (1_{a_1} + f_2).$$

This fact allows us to inductively simplify f to a composite of sums each containing one transition. The factors in this composite correspond to firings that make n reachable from m . (Here we allow the possibility of an empty composite, which corresponds to an identity morphism.) \square

Definition 22. We define the **reachability relation** of an open Petri net

$$LX \xrightarrow{i} P \xleftarrow{o} LY$$

to be the relation

$$\blacksquare P = \{(x, y) \in \mathbb{N}[X] \times \mathbb{N}[Y] \mid o(y) \text{ is reachable from } i(x)\} \subseteq \mathbb{N}[X] \times \mathbb{N}[Y].$$

Note that $\blacksquare P$ depends on the whole open Petri net $P: X \rightrightarrows Y$, not just its underlying Petri net P . By Prop. 21,

$$\blacksquare P = \{(x, y) \in \mathbb{N}[X] \times \mathbb{N}[Y] \mid \exists h: F(i)(x) \rightarrow F(o)(y)\}.$$

Here $F(i)(x)$ and $F(o)(y)$ are objects of the category FP , and the reachability relation holds iff there is a morphism in FP from the first of these to the second.

Theorem 23. *There is a lax double functor $\blacksquare: \mathbb{O}pen(\text{Petri}) \rightarrow \mathbb{R}el$, called the **reachability semantics**, that sends*

- any object X to the underlying set of the free commutative monoid $\mathbb{N}[X]$, which we denote simply as $\mathbb{N}[X]$,
- any vertical 1-morphism $f: X \rightarrow Y$ to the underlying function of $\mathbb{N}[f]$,
- any horizontal 1-cell, that is, any open Petri net

$$LX \xrightarrow{i} P \xleftarrow{o} LY,$$

to the reachability relation $\blacksquare P$.

- any 2-morphism $\alpha: P \Rightarrow P'$, that is any commuting diagram

$$\begin{array}{ccccc} LX & \xrightarrow{i} & P & \xleftarrow{o} & LY \\ Lf \downarrow & & \alpha \downarrow & & \downarrow Lg \\ LX' & \xrightarrow{i'} & P' & \xleftarrow{o'} & LY' \end{array},$$

to the square

$$\begin{array}{ccc} \mathbb{N}[X] & \xrightarrow{\blacksquare P \subseteq X \times Y} & \mathbb{N}[Y] \\ \mathbb{N}[f] \downarrow & & \downarrow \mathbb{N}[g] \\ \mathbb{N}[X'] & \xrightarrow{\blacksquare P' \subseteq X' \times Y'} & \mathbb{N}[Y']. \end{array}$$

Proof. We construct \blacksquare as the composite $G \circ \mathbb{C}sp(F)$ where

$$\mathbb{C}sp(F): \mathbb{O}pen(\text{Petri}) \rightarrow \mathbb{O}pen(\text{PetriCat})$$

is the double functor constructed in Lemma 17 and

$$G: \mathbb{O}pen(\text{PetriCat}) \rightarrow \mathbb{R}el$$

is defined as follows. Recall that we have categories of objects

$$\mathbb{O}pen(\text{PetriCat})_0 = \mathbb{R}el_0 = \text{Set}.$$

We define $G_0: \mathbb{O}pen(\text{PetriCat})_0 \rightarrow \mathbb{R}el_0$ to be the functor $\mathbb{N}: \text{Set} \rightarrow \text{Set}$. We define $G_1: \mathbb{O}pen(\text{PetriCat})_1 \rightarrow \mathbb{R}el_1$ as follows:

$$\begin{array}{ccc} L'X & \xrightarrow{i} & C \xleftarrow{o} L'Y \\ \downarrow L'f & & \downarrow \alpha \\ L'X' & \xrightarrow{i'} & C' \xleftarrow{o'} L'Y' \end{array} \quad \mapsto \quad \begin{array}{ccc} \mathbb{N}[X] & \xrightarrow{G_1 C \subseteq \mathbb{N}[X] \times \mathbb{N}[Y]} & \mathbb{N}[Y] \\ \downarrow \mathbb{N}[f] & & \downarrow \mathbb{N}[g] \\ \mathbb{N}[X'] & \xrightarrow{G_1 C' \subseteq \mathbb{N}[X'] \times \mathbb{N}[Y']} & \mathbb{N}[Y'] \end{array}$$

Recall that the set of objects of $L'X$ is $\mathbb{N}[X]$ and the set of objects of $L'Y$ is $\mathbb{N}[Y]$. We define $G_1 C$ to be the relation

$$\{(x, y) \in L'X \times L'Y \mid h: i(x) \rightarrow o(y) \text{ for some } h \text{ in } C\} \subseteq \mathbb{N}[X] \times \mathbb{N}[Y]$$

and $G_1 \alpha$ to be the inclusion

$$(\mathbb{N}[f] \times \mathbb{N}[g])G_1 C \subseteq G_1 C'.$$

To see that this inclusion is well-defined, suppose $(x, y) \in G_1 C$. Then there exists a morphism $h: i(x) \rightarrow o(y)$ in C . We thus have a morphism $\alpha(h): \alpha(i(x)) \rightarrow \alpha(o(y))$ in C' . However, on objects we have $\alpha \circ i = i' \circ L'f = i' \circ \mathbb{N}[f]$ and similarly $\alpha \circ o = o' \circ \mathbb{N}[g]$, so $\alpha(h): i'(\mathbb{N}[f](x)) \rightarrow o'(\mathbb{N}[g](y))$. It follows that $(\mathbb{N}[f] \times \mathbb{N}[g])(x, y) \in G_1 C'$.

Next we prove that G is a lax double functor. First note that by construction we have the following equalities:

$$S \circ G_1 = G_0 \circ S, \quad T \circ G_1 = G_0 \circ T.$$

Next we need the composition comparison required by Def. 27. Suppose we compose $C: X \rightarrow Y$ and $D: Y \rightarrow Z$ in $\mathbb{O}pen(\text{PetriCat})$:

$$\begin{array}{ccccc} & & C +_{L'Y} D & & \\ & \nearrow j_C & & \nwarrow j_D & \\ & C & & D & \\ \nearrow i_1 & & \nwarrow o_1 & & \nearrow i_2 \\ L'X & & L'Y & & L'Z. \\ & \nwarrow o_2 & & \swarrow i_1 & \end{array}$$

We need to prove that

$$G_1(D) \odot G_1(C) \subseteq G_1(D \odot C).$$

We have

$$G_1(D \odot C) = \{(x, z) \in L'X \times L'Z \mid \exists h: j_C i_1(x) \rightarrow j_D o_2(z)\}.$$

On the other hand,

$$G_1 C = \{(x, y) \in L'X \times L'Y \mid \exists m: i_1(x) \rightarrow o_1(y)\}$$

and

$$G_1 D = \{(y, z) \in L'Y \times L'Z \mid \exists n: i_2(y) \rightarrow o_2(z)\}$$

which compose to give the relation

$$G_1D \odot G_1C = \{(x, z) \in L'X \times L'Z \mid \exists y (x, y) \in G_1C \text{ and } (y, z) \in G_1D\}.$$

Suppose $(x, z) \in G_1D \odot G_1C$. Then there exist morphisms $m: i_1(x) \rightarrow o_1(y)$ in C and $n: i_2(y) \rightarrow o_2(z)$ in D . By commutativity of the pushout square, $j_C o_1 = j_D i_2$. Therefore, the codomain of $j_C(m)$ is $j_C o_1(y) = j_D i_2(y)$, which is also the domain of $j_D(n)$. This allows us to form the composite

$$j_D(n) \circ j_C(m): j_C i_1(x) \rightarrow j_D o_2(z).$$

Thus $(x, z) \in G_1(D \odot C)$ as desired.

We also need the identity comparison required by Def. 27. Thus, we need

$$U_{G_0(X)} \subseteq G_1(U_X)$$

for any set X . By definition, $U_X \in \mathbf{Open}(\mathbf{PetriCat})_1$ is the cospan


$$L'X \xrightarrow{1} L'X \xleftarrow{1} L'X.$$

Because $L'X$ has no non-identity morphisms, G_1 maps this to the identity relation on the set $\mathbb{N}[X]$. On the other hand, $G_0(X) = \mathbb{N}[X]$ and $U_{G_0(X)}$ is the identity relation on this set. So, the desired inclusion is actually an equality.


Finally, because \mathbf{Rel} is a degenerate double category, the composition and identity comparisons for G are trivially natural transformations. For the same reason, the diagrams in Def. 27 expressing compatibility with the associator, left unitor, and right unitor also commute trivially. It follows that G is a lax double functor.

To complete the proof, one simply computes the composite $\blacksquare = G \circ \mathbf{Csp}(F)$ and checks that it matches the description in the theorem statement. \square


The reachability semantics is only lax: given two open Petri nets $P: X \dashrightarrow Y$ and $Q: Y \dashrightarrow Z$, the composite of $\blacksquare Q$ and $\blacksquare P$ is in general a proper subset of $\blacksquare(QP)$. To see this, take P to be this open Petri net:



and take Q to be this:



Then their composite, $QP: X \rightarrow Z$, looks like this:



We have

$$\blacksquare P = \{(n, n, 0, 0) \mid n \in \mathbb{N}\} \subseteq \mathbb{N} \times \mathbb{N}^3$$

since tokens starting at A can only move to B , and similarly

$$\blacksquare Q = \{(0, 0, n, n) \mid n \in \mathbb{N}\} \subseteq \mathbb{N}^3 \times \mathbb{N}.$$

It follows that

$$\blacksquare Q \blacksquare P = \{(0, 0)\} \subseteq \mathbb{N} \times \mathbb{N}.$$

On the other hand

$$\blacksquare(QP) = \{(n, n) \mid n \in \mathbb{N}\} \subseteq \mathbb{N} \times \mathbb{N}$$

since in the composite open Petri net QP tokens can move from A to E . The point is that tokens can only accomplish this by leaving the open Petri net P , going to Q , then returning to P , then going to Q . The composite relation $\blacksquare Q \blacksquare P$ only keeps track of processes where tokens leave P , move to Q , and never reenter P .

On the other hand, the reachability semantics is maximally compatible with running Petri nets in parallel:

Theorem 24. *The reachability semantics $\blacksquare: \mathbb{O}pen(\text{Petri}) \rightarrow \mathbb{R}el$ is symmetric monoidal.*

Proof. Because $\mathbb{C}sp(F)$ is symmetric monoidal it suffices to show that

$$G: \mathbb{O}pen(\text{PetriCat}) \rightarrow \mathbb{R}el$$

is symmetric monoidal. This is simplified by that fact that $\mathbb{R}el$ is a degenerate double category. Following Def. 31, it suffices to show that

- $G_0: (\mathbb{S}et, +) \rightarrow (\mathbb{S}et, \times)$ is symmetric monoidal,
- $G_1: \mathbb{O}pen(\text{PetriCat})_1 \rightarrow \mathbb{R}el_1$ is symmetric monoidal,
- we have equations of monoidal functors

$$S \circ G_1 = G_0 \circ S, \quad T \circ G_1 = G_0 \circ T,$$

- the composition and unit comparisons are monoidal natural transformations.

To show these things, first recall that $G_0 = \mathbb{N} = K \circ J$ where $K: \mathbb{C}ommMon \rightarrow \mathbb{S}et$ is the forgetful functor and $J: \mathbb{S}et \rightarrow \mathbb{C}ommMon$ is its left adjoint. Since J is a left adjoint it preserves finite coproducts. Since $K: \mathbb{C}ommMon \rightarrow \mathbb{S}et$ is a right adjoint it preserves finite products. However, finite products in $\mathbb{C}ommMon$ are also finite coproducts. Thus, G_0 maps finite coproducts to finite products, and is thus a symmetric monoidal functor from $(\mathbb{S}et, +)$ to $(\mathbb{S}et, \times)$.

Next, suppose we are given two open Petri categories

$$L'X \xrightarrow{i} C \xleftarrow{o} L'Y, \quad L'X' \xrightarrow{i'} C' \xleftarrow{o'} L'Y'.$$

Their tensor product is

$$L'(X + X') \xrightarrow{i+i'} C + C' \xleftarrow{o+o'} L'(Y + Y') .$$

The set of objects of $L'(X+X')$ is naturally isomorphic to $\mathbb{N}[X] \times \mathbb{N}[X']$, and similarly for $L'(Y+Y')$, so we have natural isomorphisms

$$\begin{aligned} G_1(C + C') &\cong \\ \{((x, x', y, y') \in \mathbb{N}[X] \times \mathbb{N}[X'] \times \mathbb{N}[Y] \times \mathbb{N}[Y'] \mid \exists h: i(x) \rightarrow o(y) \text{ and } \exists h': i'(x') \rightarrow o'(y')\} & \\ \cong G_1(C) \times G_1(C'). & \end{aligned}$$

Using this fact one can check that G_1 is symmetric monoidal.

One can check that the equations $S \circ G_1 = G_0 \circ S$ and $T \circ G_1 = G_0 \circ T$ are equations of monoidal functors, and the composition and unit comparisons of G are trivially monoidal natural transformations because $\mathbb{R}el$ is degenerate. \square

7. CONCLUSIONS

The ideas presented here can be adapted to handle timed Petri nets, colored Petri nets with guards, and other kinds of Petri nets. One can also develop a reachability semantics for open Petri nets that are glued together along transitions as well as places. We hope to treat some of these generalizations in future work.

It would be valuable to have $\blacksquare(QP) = \blacksquare Q \blacksquare P$, since then the reachability relation for an open Petri net could be computed compositionally, not merely ‘approximated from below’ using $\blacksquare Q \blacksquare P \subseteq \blacksquare(QP)$. We conjecture that $\blacksquare(QP) = \blacksquare Q \blacksquare P$ if P and Q are ‘one-way’ open Petri nets. Here an open Petri net

$$LX \xrightarrow{i} P \xleftarrow{o} LY$$

is **one-way** if no place in the image of i appears in the target $t(\tau)$ of any transition τ of P , and no place in the image of o appears in the source $s(\tau)$ of any transition τ of P . One-way open Petri nets should be the horizontal 1-cells in a full sub-double category $\mathbb{OneWay}(\text{Petri})$ of $\mathbb{Open}(\text{Petri})$, and we conjecture that the reachability semantics restricts to an actual (not merely lax) double functor

$$\blacksquare: \mathbb{OneWay}(\text{Petri}) \rightarrow \mathbb{R}el.$$

Acknowledgements. We would like to thank Kenny Courser for help with double categories and for a careful reading of this paper. We thank Christina Vasilakopoulou for spending many hours helping us figure out how to turn Petri nets into commutative monoidal categories. We also thank Daniel Cicala, Joe Moeller, and Christian Williams for many insightful conversations.

APPENDIX A. DOUBLE CATEGORIES

What follows is a brief introduction to double categories. A more detailed exposition can be found in the work of Grandis and Paré [15, 16], and for monoidal double categories the work of Shulman [29]. We use ‘double category’ to mean what earlier authors called a ‘pseudo double category’.

Definition 25. A **double category** is a category weakly internal to Cat . More explicitly, a double category \mathbb{D} consists of:

- a **category of objects** \mathbb{D}_0 and a **category of arrows** \mathbb{D}_1 ,

- **source** and **target** functors

$$S, T: \mathbb{D}_1 \rightarrow \mathbb{D}_0,$$

an **identity-assigning** functor

$$U: \mathbb{D}_0 \rightarrow \mathbb{D}_1,$$

and a **composition** functor

$$\odot: \mathbb{D}_1 \times_{\mathbb{D}_0} \mathbb{D}_1 \rightarrow \mathbb{D}_1$$

where the pullback is taken over $\mathbb{D}_1 \xrightarrow{T} \mathbb{D}_0 \xleftarrow{S} \mathbb{D}_1$, such that

$$S(U_A) = A = T(U_A), \quad S(M \odot N) = SN, \quad T(M \odot N) = TM,$$

- natural isomorphisms called the **associator**

$$\alpha_{N, N', N''}: (N \odot N') \odot N'' \xrightarrow{\sim} N \odot (N' \odot N''),$$

the **left unitor**

$$\lambda_N: U_{T(N)} \odot N \xrightarrow{\sim} N,$$

and the **right unitor**

$$\rho_N: N \odot U_{S(N)} \xrightarrow{\sim} N$$

such that $S(\alpha), S(\lambda), S(\rho), T(\alpha), T(\lambda)$ and $T(\rho)$ are all identities and such that the standard coherence axioms hold: the pentagon identity for the associator and the triangle identity for the left and right unitor [20, Sec. VII.1].

If α, λ and ρ are identities, we call \mathbb{D} a **strict** double category.

Objects of \mathbb{D}_0 are called **objects** and morphisms in \mathbb{D}_0 are called **vertical 1-morphisms**. Objects of \mathbb{D}_1 are called **horizontal 1-cells** of \mathbb{D} and morphisms in \mathbb{D}_1 are called **2-morphisms**. A morphism $\alpha: M \rightarrow N$ in \mathbb{D}_1 can be drawn as a square:

$$\begin{array}{ccc} A & \xrightarrow{M} & B \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ C & \xrightarrow{N} & D \end{array}$$

where $f = S\alpha$ and $g = T\alpha$. If f and g are identities we call α a **globular 2-morphism**. These give rise to a bicategory:

Definition 26. Let \mathbb{D} be a double category. Then the **horizontal bicategory** of \mathbb{D} , denoted $H(\mathbb{D})$, is the bicategory consisting of objects, horizontal 1-cells and globular 2-morphisms of \mathbb{D} .

We have maps between double categories, and also transformations between maps:

Definition 27. Let \mathbb{A} and \mathbb{B} be double categories. A **double functor** $F: \mathbb{A} \rightarrow \mathbb{B}$ consists of:

- functors $F_0: \mathbb{A}_0 \rightarrow \mathbb{B}_0$ and $F_1: \mathbb{A}_1 \rightarrow \mathbb{B}_1$ obeying the following equations:

$$S \circ F_1 = F_0 \circ S, \quad T \circ F_1 = F_0 \circ T,$$

- natural isomorphisms called the **composition comparison**:

$$\phi(N, N'): F_1(N) \odot F_1(N') \xrightarrow{\sim} F_1(N \odot N')$$

and the **identity comparison**:

$$\phi_A: U_{F_0(A)} \xrightarrow{\sim} F_1(U_A)$$

whose components are globular 2-morphisms,

such that the following diagram commutes:

- a diagram expressing compatibility with the associator:

$$\begin{array}{ccc} (F_1(N) \odot F_1(N')) \odot F_1(N'') & \xrightarrow{\alpha} & F_1(N) \odot (F_1(N') \odot F_1(N'')) \\ \downarrow \phi(N, N') \odot 1 & & \downarrow 1 \odot \phi(N', N'') \\ F_1(N \odot N') \odot F_1(N'') & & F_1(N) \odot F_1(N' \odot N'') \\ \downarrow \phi(N \odot N', N'') & & \downarrow \phi(N, N' \odot N'') \\ F_1((N \odot N') \odot N'') & \xrightarrow{F_1(\alpha)} & F_1(N \odot (N' \odot N'')) \end{array}$$

- two diagrams expressing compatibility with the left and right unitors:

$$\begin{array}{ccc} F_1(N) \odot U_{F_0(A)} & \xrightarrow{\rho_{F_1(N)}} & F_1(N) \\ \downarrow 1 \odot \phi_A & & \uparrow F_1(\rho_N) \\ F_1(N) \odot F_1(U_A) & \xrightarrow{\phi(N, U_A)} & F_1(N \odot U_A) \end{array}$$

$$\begin{array}{ccc} U_{F_0(B)} \odot F_1(N) & \xrightarrow{\lambda_{F_1(N)}} & F_1(N) \\ \downarrow \phi_B \odot 1 & & \uparrow F_1(\lambda_N) \\ F_1(U_B) \odot F_1(N) & \xrightarrow{\phi(U_B, N)} & F_1(U_B \odot N). \end{array}$$

If the 2-morphisms $\phi(N, N')$ and ϕ_A are identities for all $N, N' \in \mathbb{A}_1$ and $A \in \mathbb{A}_0$, we say $F: \mathbb{A} \rightarrow \mathbb{B}$ is a **strict** double functor. If on the other hand we drop the requirement that these 2-morphisms be invertible, we call F a **lax** double functor.

Definition 28. Let $F: \mathbb{A} \rightarrow \mathbb{B}$ and $G: \mathbb{A} \rightarrow \mathbb{B}$ be lax double functors. A **transformation** $\beta: F \Rightarrow G$ consists of natural transformations $\beta_0: F_0 \Rightarrow G_0$ and $\beta_1: F_1 \Rightarrow G_1$ (both usually written as β) such that

- $S(\beta_M) = \beta_{SM}$ and $T(\beta_M) = \beta_{TM}$ for any object $M \in \mathbb{A}_1$,
- β commutes with the composition comparison, and
- β commutes with the identity comparison.

Shulman defines a 2-category **Dbf** of double categories, double functors, and transformations [29]. This has finite products. In any 2-category with finite products we can define a pseudomonoid [10], which is a categorification of the concept of monoid. For example, a pseudomonoid in **Cat** is a monoidal category.

Definition 29. A **monoidal double category** is a pseudomonoid in **Dbl**. Explicitly, a monoidal double category is a double category equipped with double functors $\otimes: \mathbb{D} \times \mathbb{D} \rightarrow \mathbb{D}$ and $I: * \rightarrow \mathbb{D}$ where $*$ is the terminal double category, along with invertible transformations called the **associator**:

$$A: \otimes \circ (1_{\mathbb{D}} \times \otimes) \Rightarrow \otimes \circ (\otimes \times 1_{\mathbb{D}}),$$

left unitor:

$$L: \otimes \circ (1_{\mathbb{D}} \times I) \Rightarrow 1_{\mathbb{D}},$$

and **right unitor**:

$$R: \otimes \circ (I \times 1_{\mathbb{D}}) \Rightarrow 1_{\mathbb{D}}$$

satisfying the pentagon axiom and triangle axioms.

This definition neatly packages a large quantity of information. Namely:

- \mathbb{D}_0 and \mathbb{D}_1 are both monoidal categories.
- If I is the monoidal unit of \mathbb{D}_0 , then U_I is the monoidal unit of \mathbb{D}_1 .
- The functors S and T are strict monoidal.
- \otimes is equipped with composition and identity comparisons

$$\chi: (M_1 \otimes N_1) \odot (M_2 \otimes N_2) \xrightarrow{\sim} (M_1 \odot M_2) \otimes (N_1 \odot N_2)$$

$$\mu: U_{A \otimes B} \xrightarrow{\sim} (U_A \otimes U_B)$$

making three diagrams commute as in Def. 27.

- The associativity isomorphism for \otimes is a transformation between double functors.
- The unit isomorphisms are transformations between double functors.

Definition 30. A **braided monoidal double category** is a monoidal double category equipped with an invertible transformation

$$\beta: \otimes \Rightarrow \otimes \circ \tau$$

called the **braiding**, where $\tau: \mathbb{D} \times \mathbb{D} \rightarrow \mathbb{D} \times \mathbb{D}$ is the twist double functor sending pairs in the object and arrow categories to the same pairs in the opposite order. The braiding is required to satisfy the usual two hexagon identities [20, Sec. XI.1]. If the braiding is self-inverse we say that \mathbb{D} is a **symmetric monoidal double category**.

In other words:

- \mathbb{D}_0 and \mathbb{D}_1 are braided (resp. symmetric) monoidal categories,
- the functors S and T are strict braided monoidal functors, and
- the braiding is a transformation between double functors.

Definition 31. A **monoidal lax double functor** $F: \mathbb{C} \rightarrow \mathbb{D}$ between monoidal double categories \mathbb{C} and \mathbb{D} is a lax double functor $F: \mathbb{C} \rightarrow \mathbb{D}$ such that

- F_0 and F_1 are monoidal functors,
- $SF_1 = F_0S$ and $TF_1 = F_0T$ are equations between monoidal functors, and
- the composition and unit comparisons $\phi(N_1, N_2): F_1(N_1) \odot F_1(N_2) \rightarrow F_1(N_1 \odot N_2)$ and $\phi_A: U_{F_0(A)} \rightarrow F_1(U_A)$ are monoidal natural transformations.

The monoidal lax double functor is **braided** if F_0 and F_1 are braided monoidal functors and **symmetric** if they are symmetric monoidal functors.

REFERENCES

- [1] J. Baez and K. Courser, Coarse-graining open Markov processes. Available as [arXiv:1710.11343](https://arxiv.org/abs/1710.11343). (Referred to on page 3, 15.)
- [2] J. Baez and K. Courser, Structured cospans, to appear. (Referred to on page 5, 9, 12, 14.)
- [3] J. Baez and B. Pollard, A compositional framework for reaction networks, *Rev. Math. Phys.* **29**, 1750028. Available as [arXiv:1704.02051](https://arxiv.org/abs/1704.02051). (Referred to on page 1.)
- [4] P. Baldan, A. Corradini, H. Ehrig, and R. Heckel, Compositional semantics for open Petri nets based on deterministic processes, *Math. Str. Comp. Sci.* **15** (2005), 1–35. (Referred to on page 1.)
- [5] P. Baldan, F. Bonchi, F. Gadducci and G. V. Monreale, Modular encoding of synchronous and asynchronous interactions using open Petri nets, *Sci. Comp. Prog.* **109** (2015), 96–124. (Referred to on page 1.)
- [6] R. Bruni, H. C. Melgratti and U. Montanari, A connector algebra for P/T nets interactions, in *Concurrency Theory (CONCUR '11)*, Lecture Notes in Computer Science 6901, Springer, Berlin, 2011, pp. 312–326. (Referred to on page 1.)
- [7] R. Bruni, J. Meseguer, U. Montanari and V. Sassone, Functorial models for Petri nets, *Information and Computation* **170** (2001), 207–236. (Referred to on page 1.)
- [8] F. Clerc, H. Humphrey and P. Panangaden, Bicategories of Markov processes, in *Models, Algorithms, Logics and Tools*, Lecture Notes in Computer Science **10460**, Springer, Berlin, 2017, pp. 112–124. (Referred to on page 3.)
- [9] K. Courser, A bicategory of decorated cospans, *Theory Appl. Categ.* **32** (2017), 995–1027. Available as [arXiv:1605.08100](https://arxiv.org/abs/1605.08100). (Referred to on page 3.)
- [10] B. Day and R. Street, Monoidal bicategories and Hopf algebroids, *Adv. Math.* **129** (1997), 99–157. (Referred to on page 25.)
- [11] P. Degano, J. Meseguer and U. Montanari, Axiomatizing net computations and processes, in *Logic in Computer Science, 1989*, IEEE, New Jersey, pp. 175–185. Available at <https://www.computer.org/csdl/proceedings/lics/1989/1954/00/00039172.pdf>. (Referred to on page 6.)
- [12] C. Ehresmann, Catégories structurées III: Quintettes et applications covariantes, *Cah. Top. Géom. Diff.* **5** (1963), 1–22. (Referred to on page 3.)
- [13] C. Ehresmann, *Catégories et Structures*, Dunod, Paris, 1965. (Referred to on page 3.)
- [14] C. Girault and R. Valk, *Petri Nets for Systems Engineering: a Guide to Modeling, Verification, and Applications*, Springer, Berlin, 2013. (Referred to on page 1.)
- [15] M. Grandis and R. Paré, Limits in double categories, *Cah. Top. Géom. Diff.* **40** (1999), 162–220. (Referred to on page 14, 23.)
- [16] M. Grandis and R. Paré, Adjoints for double categories, *Cah. Top. Géom. Diff.* **45** (2004), 193–240. (Referred to on page 23.)
- [17] K. Jensen and L. M. Kristensen, *Coloured Petri Nets: Modelling and Validation of Concurrent Systems*, Springer, Berlin 2009. (Referred to on page 1.)
- [18] E. Lerman and D. Spivak, An algebra of open continuous time dynamical systems and networks. Available as [arXiv:1602.01017](https://arxiv.org/abs/1602.01017). (Referred to on page 3.)
- [19] R. Lipton, The reachability problem is exponential-space-hard, Department of Computer Science, Yale University **62** (1976). (Referred to on page 18.)
- [20] S. Mac Lane, *Categories for the Working Mathematician*, Springer, Berlin, 1998. (Referred to on page 24, 26.)
- [21] E. Mayr, An algorithm for the general Petri net reachability problems, *SIAM J. Comput.* **13** (1984), 441–460. (Referred to on page 18.)
- [22] J. Meseguer and U. Montanari, Petri nets are monoids, *Information and Computation* **88** (1990), 105–155. (Referred to on page 2, 5.)
- [23] J. L. Peterson, *Petri Net Theory and the Modeling of Systems*, Prentice–Hall, New Jersey, 1981. (Referred to on page 1.)
- [24] J. Rathke, P. Sobociński and O. Stephens, Compositional reachability in Petri nets, in *International Workshop on Reachability Problems*, Lecture Notes in Computer Science **8762**, Springer, Berlin, 2014. Available at <http://users.ecs.soton.ac.uk/ps/papers/rp2014.pdf>. (Referred to on page 1.)
- [25] V. Sassone, Strong concatenable processes: an approach to the category of Petri net computations, *BRICS Report Series*, Dept. of Computer Science, U. Aarhus, 1994. Available at <https://tidsskrift.dk/brics/article/view/21610/19059>. (Referred to on page 6.)

- [26] V. Sassone, On the category of Petri net computations, in *Colloquium on Trees in Algebra and Programming*, Springer, Berlin, 1995. Available at <https://eprints.soton.ac.uk/261951/1/strong-conf.pdf>. (Referred to on page 6.)
- [27] V. Sassone, An axiomatization of the algebra of Petri net concatenable processes, in *Theoretical Computer Science* **170** (1996), 277–296. Available at <https://eprints.soton.ac.uk/261820/1/P-of-N-Off.pdf>. (Referred to on page 6.)
- [28] V. Sassone and P. Sobociński, A congruence for Petri nets, *Electronic Notes in Theoretical Computer Science* **127** (2005), 107–120. Available at <https://eprints.soton.ac.uk/262302/1/petriCongPNGToff.pdf>. (Referred to on page 1, 3, 6.)
- [29] M. Shulman, Constructing symmetric monoidal bicategories. Available as [arXiv:1004.0993](https://arxiv.org/abs/1004.0993). (Referred to on page 4, 23, 25.)
- [30] P. Sobociński and O. Stephens, Reachability via compositionality in Petri nets. Available as [arXiv:1303.1399](https://arxiv.org/abs/1303.1399). (Referred to on page 1.)