

Name: _____

Score: _____ / 100

Student ID: _____

DO NOT OPEN THE EXAM UNTIL YOU ARE TOLD TO DO SO

	1	2	3	4	5	6	7	8	9	10	11	Total
✓												100
Score												
Pts. Possible	10	10	10	10	10	10	10	10	10	10	10	110

INSTRUCTIONS FOR STUDENTS

- Questions are on both sides of the paper. This is an 11 question exam.
- Students have 2 hours and 30 minutes to complete the exam.
- The test will be out of **100** points. You may attempt as many problems or parts of problems as you would like. The highest possible score is therefore **110** points.
- In the above table, the row with the ✓, is for you to keep track of the problems you are attempting/completing.
- You may complete parts of problems, as partial credit will be given based on correctness, completeness, and ideas that are leading to the correct solutions.
- **PLEASE SHOW ALL WORK. Any unjustified claims will receive no credit. This means you need to state which test you are using for series questions!** Clearly box your final answer.
- No notes, textbooks, phones, calculators, etc. are allowed for the exam.
- The last page of the test can be used for scratch work.

GOOD LUCK!

FORMULAS:

$\frac{d^2y}{dx^2} = \frac{d}{dt} \left(\frac{dy}{dx} \right)$	$\frac{dy}{dx} = \frac{\frac{dr}{d\theta} \sin(\theta) + r \cos(\theta)}{\frac{dr}{d\theta} \cos(\theta) - r \sin(\theta)}$
$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n, \quad \text{for all } x < 1$	$\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}, \quad \text{for all } x \in \mathbb{R}$
$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}, \quad \text{for all } x \in \mathbb{R}$	$\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}, \quad \text{for all } x \in \mathbb{R}$
$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n}, \quad \text{for } x \in (-1, 1]$	$\arctan(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}, \quad \text{for } x \leq 1$
$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n, \quad \text{for } x-a < R$	$(1+x)^m = \sum_{n=0}^{\infty} \binom{m}{n} x^n, \quad \text{for } x < 1$

1) (10 pts.) Consider the parametric curve for $0 \leq t \leq \pi$, given by

$$x = \sin(t)$$

$$y = \cos(t)$$

(a) (3 pts.) Sketch the graph of the parametric curve and identify the direction for increasing values of t with an arrow.


(b) (4 pts.) Find the values of t where the the tangent line is vertical.

(c) (3 pts.) Determine for which values of t is the curve concave up.

2) (10 pts.) Consider the polar curve for $0 \leq t \leq \pi$, given by

$$r = 4 \sin(3\theta)$$



- (a) (4 pts.) Plot the curve on the given polar grid below.
(b) (6 pts.) Find the equation of the tangent line at $\theta = \frac{\pi}{6}$.



3) (10 pts.) Find the area of the region that lies inside one of the small petal and one large petal of

$$r = 1 + 2 \cos(2\theta)$$

Hint: Identities that may be helpful: 1) $\sin^2(2\theta) = \frac{1}{2} - \frac{1}{2} \cos(4\theta)$, 2) $\cos^2(2\theta) = \frac{1}{2} + \frac{1}{2} \cos(4\theta)$.



4) (5 pts.) (a) Determine whether the sequence converges or diverges:

$$a_n = \left(1 + \sin\left(\frac{1}{n}\right)\right)^{\cot\left(\frac{1}{n}\right)}.$$

(5 pts.) (b) Determine whether the sequence converges or diverges:

$$a_n = \arctan(2n).$$

5) (5 pts.) (a) Determine whether the series is convergent or divergent:

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt[n]{n}}.$$

(5 pts.) (b) Determine the rational number $\frac{p}{q}$, for $p, q \in \mathbb{Z}$ which represents the decimal expansion for $0.\overline{123} = 0.123123\dots$, **using infinite series**.

6) (5 pts.) Determine whether the series is convergent or divergent

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln(n))^3}.$$

(b) (5 pts.) Determine whether the series is convergent or divergent

$$\sum_{n=1}^{\infty} \sin\left(\frac{1}{n}\right).$$

7) (5 pts.) (a) Determine whether the series is absolutely convergent, conditionally convergent, or divergent:

$$\sum_{n=1}^{\infty} \frac{n!}{e^{n^2}}.$$

(5 pts.) (b) Determine whether the series is absolutely convergent, conditionally convergent, or divergent:

$$\sum_{n=0}^{\infty} \left(1 + \frac{1}{n}\right)^{n^2}.$$

8) (10 pts.) Determine whether the series is absolutely convergent, conditionally convergent, or divergent:

$$\sum_{n=2}^{\infty} (-1)^n \frac{n^3}{n^4 + 1}.$$

9) (10 pts.) Find the radius of convergence and interval of convergence for the following power series.

$$\sum_{n=0}^{\infty} (-1)^n \frac{(x-3)^n}{2n+1}$$

10) (10 pts.) Compute the first 4 non-zero terms of the Taylor series centered at $a = 3$ for the following function using the definition of a Taylor series.

$$f(x) = e^x$$

11) (5 pts.) (a) Compute the following integral using Taylor series, and leave the answer as a Taylor series.

$$\int \cos(x^2) dx$$

(5 pts.) (b) Find the Taylor series centered at $a = 0$ for the function and write as a single series.

$$f(x) = e^{-x^2} + \cos(x)$$

THIS PAGE IS LEFT BLANK FOR ANY SCRATCH WORK

END OF TEST