

INFINITE-ENERGY 2D STATISTICAL SOLUTIONS TO THE EQUATIONS OF INCOMPRESSIBLE FLUIDS

JAMES P. KELLIHER

ABSTRACT. We develop the concept of an infinite-energy statistical solution to the Navier-Stokes and Euler equations in the whole plane. We use a velocity formulation with enough generality to encompass initial velocities having bounded vorticity, which includes the important special case of vortex patch initial data. Our approach is to use well-studied properties of statistical solutions in a ball of radius R to construct, in the limit as R goes to infinity, an infinite-energy solution to the Navier-Stokes equations. We then construct an infinite-energy statistical solution to the Euler equations by making a vanishing viscosity argument.

Compiled on *September 15, 2009*

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| 2. Function spaces and the Biot-Savart Law | 3 |
| 3. Projection operators | 6 |
| 4. Weak deterministic solutions | 9 |
| 5. Deterministic expanding domain limit | 11 |
| 6. Definition of infinite-energy statistical solutions | 17 |
| 7. Construction of Navier-Stokes solutions | 19 |
| 8. Construction of Euler solutions | 27 |
| Acknowledgements | 32 |
| References | 32 |

1. INTRODUCTION

We develop the concept of a statistical solution to the Navier-Stokes (SSNS) or Euler equations (SSE) in the plane for an important class of velocity fields having sufficient decay of the vorticity at infinity to recover uniquely the velocity field from the vorticity. In particular, this class of velocity fields includes the important case of a vortex patch: a velocity field whose initial vorticity is the characteristic function of a bounded domain.

Our starting point is the velocity formulation of a SSNS on a bounded domain given by Foias in [5]. (A highly accessible account of the theory of SSNSs is given in [6], to which we refer often.) We adapt this formulation

2000 *Mathematics Subject Classification.* 76D06, 76D05.

slightly, of necessity changing the energy equality and using the same class of test functions as for homogeneous solutions in the plane (Section 6). Our definition of an infinite-energy SSE is the same with the viscosity set to zero. We construct our infinite-energy SSNS by showing that it is the limit, in a special sense, of a sequence of statistical solutions on balls of radius R as $R \rightarrow \infty$. At the core of our approach is the expanding domain limit for deterministic solutions to the Navier-Stokes equations established in [9], extended to handle infinite-energy solutions. We then construct our infinite-energy SSE by making a vanishing viscosity argument.

In the deterministic setting, the infinite-energy solutions that we consider correspond to an initial velocity lying in the space E_m of [2] and [3]. A vector v belongs to E_m if it is divergence-free and can be written in the form $v = \sigma + v'$, where v' is in $L^2(\mathbb{R}^2)$ and where σ is a smooth stationary solution to the Euler equations whose vorticity is radially symmetric and compactly supported. (See Section 2 for more details.) A unique global solution to the Navier-Stokes equations exists and remains in the space E_m for all time as long as the forcing has finite energy. The same can be said of solutions to the Euler equations if one imposes restrictions on the initial vorticity; for instance, that it lie in $L^1 \cap L^\infty$. This encompasses the case of a classical vortex patch—initial vorticity that equals the characteristic function of a bounded domain. (See also Corollary 2.3.)

A key parameter of any vortex patch is the total mass of its vorticity,

$$m = \int_{\mathbb{R}^2} \omega.$$

Only when $m = 0$ will the velocity field have finite energy (lie in L^2), which excludes the case of a classical vortex patch. In a sense, m measures how infinite the energy is.

When working with statistical solutions one would like to allow m to take on different values, because classical vortex patches that are nearly identical will typically have different values of m . Thus, we need to consider the spaces E_m for all values of m simultaneously.

We give a velocity rather than a vorticity formulation of our statistical solutions for several reasons. First, a vorticity formulation would require imposing higher regularity on the initial vorticity than required for solutions to the Navier-Stokes equations: it would be technically quite difficult to assume anything weaker than the initial vorticity lying in L^1 , as in [1]. Second, it would be hard to obtain convergence of the vorticity in the vanishing viscosity limit, even with higher regularity of the initial data, without knowing that the velocity decays at infinity, and this does not come from the Biot-Savart law. Third, a start in this direction has already been made in [4] for time-independent solutions to damped and driven Navier-Stokes and Euler equations in the vorticity formulation.

Constantin and Ramos do not specifically address infinite-energy solutions in [4]; however, their definition of such a solution requires no change at all to

encompass infinite-energy solutions and neither does their proof of the vanishing viscosity limit. Their construction of a stationary statistical solution to the Navier-Stokes equations as a long-time average of a deterministic solution to the damped and driven Navier-Stokes equations does assume finite energy. Nothing deep need be done, however, to extend their construction to allow infinite energy solutions (and so allow initial vortex patch data): one need only assume infinite-energy forcing.

This paper is organized as follows: In Section 2, we define the function spaces in which we will work. We characterize the projection operator we will use to construct the initial velocities in Ω_R in Section 3. In Section 4, we define weak deterministic solutions to the Navier-Stokes and Euler equations and give the basic well-posedness and regularity results for such solutions. The deterministic expanding domain limit of [9] is established for infinite-energy solutions in Section 5. We give the definition of a statistical solution to the Navier-Stokes equations in velocity form, for finite as well as infinite energy, in Section 6, and construct an infinite-energy statistical solution to the Navier-Stokes equations in Section 7, showing that it is unique. In Section 8 we construct an infinite-energy statistical solution to the Euler equations using a vanishing viscosity argument.

2. FUNCTION SPACES AND THE BIOT-SAVART LAW

Let

$\Omega_R =$ the disk of radius R centered at the origin,

with $\Omega_\infty = \mathbb{R}^2$, and define the classical function spaces of incompressible fluid mechanics,

$$\begin{aligned} H_R &= H(\Omega_R) = \{u \in (L^2(\Omega_R))^2 : \operatorname{div} u = 0, u \cdot \mathbf{n} = 0 \text{ on } \partial\Omega_R\}, \\ V_R &= V(\Omega_R) = \{u \in (H^1(\Omega_R))^2 : \operatorname{div} u = 0, u = 0 \text{ on } \partial\Omega_R\}, \\ H &= H_\infty = H(\mathbb{R}^2), \quad V = V_\infty = V(\mathbb{R}^2). \end{aligned}$$

We endow H_R with the L^2 -norm. For V_R , we use the H^1 -norm:

$$\|u\|_{V_R} = \|u\|_{L^2(\Omega_R)} + \|\nabla u\|_{L^2(\Omega_R)}. \quad (2.1)$$

Note, in particular, that

$$\|u\|_{L^2(\Omega_R)} \leq \|u\|_{V_R}, \quad \|\nabla u\|_{L^2(\Omega_R)} \leq \|u\|_{V_R}. \quad (2.2)$$

Had we used the Poincare inequality to replace Equation (2.1) with the equivalent norm that includes only the second term, as is normally done for a bounded domain, it would have introduced a factor of R in the right-hand side of the first inequality, preventing us from having a consistent norm with which to compare solutions on Ω_R for different values of R .

Our deterministic infinite energy solutions will lie in the space E_m of [3]. A vector v belongs to E_m if it is divergence-free and can be written in the

form $v = \sigma + v'$, where v' is in $L^2(\mathbb{R}^2)$ and where σ is a *stationary vector field*, meaning that σ is of the form,

$$\sigma = \left(-\frac{x^2}{r^2} \int_0^r \rho g(\rho) d\rho, \frac{x^1}{r^2} \int_0^r \rho g(\rho) d\rho \right) \quad (2.3)$$

with g in $C_0^\infty(\mathbb{R})$. The subscript $m \in \mathbb{R}$ is the integral over all space of the vorticity,

$$\omega(v) = \partial_1 v^2 - \partial_2 v^1.$$

E_m is an affine space; fixing an origin, σ , in E_m we can define a “norm” by $\|\sigma + v'\|_{E_m} = \|v'\|_{L^2(\Omega)}$. Convergence in E_m is equivalent to convergence in the L^2 -norm to a vector in E_m .

We will find it convenient to fix a choice of origin for E_m as follows. For E_1 we choose σ_1 of the form Equation (2.3) with $\omega(\sigma_1)$ supported in the unit disk and with

$$\int_{\Omega_1} \omega(\sigma_1) = \int_{\mathbb{R}^2} \omega(\sigma_1) = 1.$$

We can then use $\sigma_m = m\sigma_1$ as an origin for E_m .

Let ψ_{σ_1} be a given fixed stream function for σ_1 . As in [10], ψ_{σ_1} is radially symmetric with

$$\psi_{\sigma_1}(x) = C_2 + \frac{1}{2\pi} \log |x| \quad (2.4)$$

for all $|x| \geq 1$. For $|x| \geq 1$, $|\sigma_1(x)| = 1/|x|$ by Equation (2.3), and

$$\begin{aligned} \|\sigma_1\|_{H^1(\Omega_R \setminus \Omega_{R-1})}^2 &= 2\pi \int_{R-1}^R \frac{1}{r^2} r dr + 2\pi \int_{R-1}^R \frac{1}{r^4} r dr \\ &= 2\pi \log(R/(R-1)) + \pi [(R-1)^{-2} - R^{-2}] \rightarrow 0 \text{ as } R \rightarrow \infty. \end{aligned} \quad (2.5)$$

Equation (2.4) also gives $\Delta\sigma_1 = \Delta\nabla^\perp\psi_{\sigma_1} = \nabla^\perp\Delta\psi_{\sigma_1} = 0$ on Ω_1^C .

The spaces H_R and E_m , or V_R and $E_m \cap \dot{H}^1(\mathbb{R}^2)$, where $\dot{H}^1(\mathbb{R}^2)$ is the set of all functions whose gradient lies in $L^2(\mathbb{R}^2)$, are the appropriate ones for initial velocities for weak deterministic solutions to the Navier-Stokes equations, but for the Euler equations more regularity is required to obtain well-posedness. Rather than being as general as possible, we will assume that for deterministic solutions the initial vorticity lies in L^∞ for solutions on Ω_R and in $L^{p_0} \cap L^\infty$, for some $p_0 < 2$ for solutions on \mathbb{R}^2 . Slightly unbounded vorticities could be handled, as in [9], with little complication. This gives not only existence but uniqueness of the solutions. (The uniqueness of solutions for bounded initial vorticity is due to Yudovich [13], as is the uniqueness for unbounded vorticities [14].)

Thus, we fix $p_0 < 2$, and define the spaces

$$\mathbb{Y}_m = \{u \in E_m : \omega(u) \in L^{p_0} \cap L^\infty\},$$

with “norm”

$$\|u\|_{\mathbb{Y}_m} = \|u\|_{E_m} + \|\omega(u - \sigma_m)\|_{L^{p_0} \cap L^\infty}$$

and

$$\mathbb{Y}(\Omega_R) = \{u \in H(\Omega_R) \cap H^1(\Omega_R) : \omega(u) \in L^\infty\}$$

with norm

$$\|u\|_{\mathbb{Y}(\Omega_R)} = \|u\|_{H^1(\Omega_R)} + \|\omega(u)\|_{L^{p_0} \cap L^\infty(\Omega_R)}, \quad R < \infty.$$

Because $L^{p_0}(\Omega_R) \subseteq L^\infty(\Omega_R)$, using only the L^∞ -norm of $\omega(u)$ in the $\mathbb{Y}(\Omega_R)$ -norm would give a simpler, equivalent norm. We avoid doing this, however, for the same reason we avoided the use of Poincaré's inequality in defining the V_R -norm in Equation (2.1).

For statistical solutions in the whole plane, we do not want to assume that the value of m is fixed, so we must deal with larger spaces. For statistical solutions to the Navier-Stokes equations we will use

$$\mathbb{E} = \bigcup_{m \in \mathbb{R}} E_m \text{ and } \mathbb{E}^1 = \bigcup_{m \in \mathbb{R}} E_m \cap \dot{H}^1(\mathbb{R}^2)$$

and for statistical solutions to the Euler equations we will use

$$\mathbb{Y} = \bigcup_{m \in \mathbb{R}} \mathbb{Y}_m. \quad (2.6)$$

\mathbb{E} , \mathbb{Y} , and \mathbb{E}^1 are function spaces, being closed under addition, with the norms

$$\begin{aligned} \|\sigma_m + v\|_{\mathbb{E}} &= |m| + \|v\|_{L_2}, \quad \|u\|_{\mathbb{Y}} = \|u\|_{\mathbb{E}} + \|\omega(u)\|_{L^{p_0} \cap L^\infty}, \\ \|\sigma_m + v\|_{\mathbb{E}^1} &= \|\sigma_m + v\|_{\mathbb{E}} + \|\nabla v\|_{L^2(\mathbb{R}^2)}. \end{aligned}$$

These norms induce metrics on their respective spaces. Because H and V are separable, so too are \mathbb{E} and \mathbb{E}^1 . The space \mathbb{Y} , however, is not separable, because $L^\infty(\mathbb{R}^2)$ is not.

There exists a unique decomposition of any u in \mathbb{E} , \mathbb{Y} , or \mathbb{E}^1 of the form $u = \sigma_m + v$, $m \in \mathbb{R}$, $v \in H$. Given such a u , we define

$$m(u) = m, \quad \sigma(u) = \sigma_{m(u)}. \quad (2.7)$$

Definition 2.1. We say that the support of a measure μ on the function space X is *bounded in X* if

$$\text{supp } \mu \subseteq \{u \in X : \|u\|_X \leq M\} \text{ for some } M < \infty.$$

If Y is a subspace of X , we say that the support of a measure μ is (X, Y) -*bounded* if

$$\text{supp } \mu \subseteq \{u \in Y : \|u\|_X \leq M\} \text{ for some } M < \infty.$$

That is, the support of μ lies in the subspace, but only its norm in the full space is controlled.

Lemma 2.2. [*Biot-Savart law*] *Let p be in $[1, 2)$ and let $q > 2p/(2-p)$. For any vorticity ω in $L^p(\mathbb{R}^2)$ there exists a unique divergence-free vector field u*

in $L^p(\mathbb{R}^2) + L^q(\mathbb{R}^2)$ whose curl is ω , with u being given by the Biot-Savart law,

$$u = K * \omega. \quad (2.8)$$

Here, K is the Biot-Savart kernel, $K(x) = (1/2\pi)x^\perp/|x|^2$.

Proof. See the proof of Proposition 3.1.1 p. 44-45 of [3]. \square

Corollary 2.3. *For any vorticity ω in $L^1 \cap L^\infty(\mathbb{R}^2)$ there exists a unique divergence-free vector field u in $L^\infty(\mathbb{R}^2)$ whose curl is ω , with u given by Equation (2.8). If ω is also compactly supported then u lies in E_m , where $m = \int_{\mathbb{R}^2} \omega$.*

Proof. By Lemma 2.2 applied with $p = 1$, $K * \omega$ is the unique vector field in $L^1 \cap L^\infty$ whose vorticity is ω . But also,

$$\begin{aligned} \|K * \omega\|_{L^\infty} &\leq \|(\chi_{\Omega_1} K) * \omega\|_{L^\infty} + \|(1 - \chi_{\Omega_1})K * \omega\|_{L^\infty} \\ &\leq \|\chi_{\Omega_1} K\|_{L^1} \|\omega\|_{L^\infty} + \|(1 - \chi_{\Omega_1})K\|_{L^\infty} \|\omega\|_{L^1} \\ &\leq C \|\omega\|_{L^1 \cap L^\infty}. \end{aligned}$$

Here, χ_A is the characteristic function of A . This shows that, in fact, $K * \omega$ is in L^∞ and is the unique such vector field. The last statement in the corollary follows from Lemma 1.3.1 of [3]. \square

3. PROJECTION OPERATORS

Let $\mathbf{P}_{V_R} : \mathbb{E}^1 \rightarrow V_R$ be restriction to Ω_R followed by projection onto V_R . \mathbf{P}_{V_R} is well-defined because as Hilbert spaces V_R is a closed subspace of $H_{div}^1(\Omega_R)$, the space of all divergence-free vector fields in $(H^1(\Omega_R))^2$, endowed with the inner product,

$$\langle u, v \rangle_{H_{div}^1(\Omega_R)} = (u, v) + (\nabla u, \nabla v).$$

We can describe \mathbf{P}_{V_R} explicitly by characterizing V_R^\perp , the orthogonal complement of V_R in $H_{div}^1(\Omega_R)$. By definition, w is in V_R^\perp if and only if $\langle w, v \rangle_{H_{div}^1(\Omega_R)} = 0$ for all v in V_R . Treating Δw as a distribution, integrating by parts gives $(w - \Delta w, v) = 0$ for all v in \mathcal{V}_R , where $\mathcal{V}_R = V_R \cap \mathcal{D}(\Omega_R)$. It follows from this that w is in V_R^\perp if and only if

$$\Delta w - w = \nabla p \quad (3.1)$$

for some p in $L^2(\Omega)$ (see, for instance, Proposition I.1.1 of [12]) and, of course, $\operatorname{div} w = 0$.

Now let u lie in $H_{div}^1(\Omega_R)$ and let $\bar{u} = \mathbf{P}_{V_R} u$. Then $w = u - \bar{u}$ lies in V_R^\perp so from Equation (3.1),

$$\begin{cases} \Delta \bar{u} - \bar{u} = \Delta u - u + \nabla p & \text{in } \Omega_R, \\ \operatorname{div} \bar{u} = \Delta p = 0 & \text{in } \Omega_R, \\ \bar{u} = 0 & \text{on } \partial\Omega_R. \end{cases} \quad (3.2)$$

Equality is to hold in a weak sense in Equation (3.2). Since u is in H^1 , however, $f = \Delta u - u + \bar{u}$ is in H^{-1} , which is sufficient to conclude that

\bar{u} is in V and p is in L^2 . (See, for instance, Remark I.2.6 of [12].) Also, the solution to Equation (3.2) is unique because otherwise it would follow that -1 is an eigenvalue of the Stokes operator, $-\mathcal{P}\Delta$, where \mathcal{P} is the Leray projector. But the Stokes operator is positive-definite, so all its eigenvalues are positive.

The estimates involving the operator \mathbf{P}_{V_R} are hard to prove directly using this characterization. It is simpler to employ an approximate projection operator \mathbf{U}_R , and use the fact that projection into V_R gives the closest vector field in V_R to the vector field being projected. (The same idea is used for projection into H_R in [10, 7].)

To define \mathbf{U}_R we need two cutoff functions, φ_R and h_R .

Let φ_1 in $C^\infty(\Omega_1)$ take values in $[0, 1]$ and be defined so that $\varphi_1 = 1$ on $\Omega_{1/2}$ and so that φ_1 and $\nabla\varphi_1$ are both zero on $\partial\Omega_R$. Let $\varphi_R(\cdot) = \varphi_1(\cdot/R)$. Observe that φ_R and $\nabla\varphi_R$ both vanish on $\partial\Omega_R$.

Let g in $C^\infty([0, 3/4])$ taking values in $[0, 1]$ be defined so that $g(0) = g'(0) = 0$ and $g = 1$ on $[1/2, 1]$. Then define h_R in $C^\infty(\Omega_R)$ by $h_R(x) = g(R - |x|)$ for points x in $\Omega_R \setminus \Omega_{R-1}$ and $h_R = 1$ on Ω_{R-1} . Observe that

$$\|h_R\|_{C^k} \leq C_k, \quad (3.3)$$

$k = 0, 1, \dots$, for constants C_k independent of R in $[1, \infty)$. Also, $h_R = 0$ and $\nabla h_R = 0$ on $\partial\Omega_R$.

Definition 3.1. Define $\mathbf{U}_R: \mathbb{E}^1 \rightarrow V_R$ by

$$\mathbf{U}_R(u) = \nabla^\perp(h_R(\psi_{\sigma_m} - \psi_{\sigma_m}(R))) + \nabla^\perp(\varphi_R\psi_v)$$

for $u = \sigma_m + v$ in E_m . Here, ψ_v is the stream function for v chosen so that $\int_{\Omega_R} \psi_v = 0$ on $\partial\Omega_R$.

Lemma 3.2. \mathbf{P}_{V_R} maps \mathbb{E}^1 continuously onto V_R with

$$\|u - \mathbf{P}_{V_R}u\|_{H^1(\Omega_R)} \leq C \|u - \sigma(u)\|_{H^1(\Omega_R \setminus \Omega_{R/2})} + |m(u)|\beta(R), \quad (3.4)$$

where

$$\beta(R) = \|\sigma_1\|_{H^1(\Omega_R \setminus \Omega_{R-1})} \rightarrow 0 \text{ as } R \rightarrow \infty,$$

and

$$\|\mathbf{P}_{V_R}(u - \sigma_m)\|_{V_R} \leq \|u - \sigma_m\|_V, \quad (3.5)$$

$$\|\mathbf{P}_{V_R}u\|_{V_R} \leq \|u\|_{\mathbb{E}^1} + C|m(u)|. \quad (3.6)$$

Proof. That \mathbf{P}_{V_R} maps onto V_R is clear, and it is continuous because the restriction and the projection operators are continuous.

To prove Equation (3.4), let $u = \sigma_m + v$ in $E_m \cap \dot{H}^1(\mathbb{R}^2)$. Then

$$\|u - \mathbf{U}_R u\|_{H^1(\Omega_R)} \leq \|\sigma_m - \mathbf{U}_R \sigma_m\|_{H^1(\Omega_R)} + \|v - \mathbf{U}_R v\|_{H^1(\Omega_R)}. \quad (3.7)$$

It follows from the proof of Lemma 4.2 of [9] that

$$\|v - \mathbf{U}_R v\|_{H^1(\Omega_R)} \leq C \|u - \sigma(u)\|_{H^1(\Omega_R \setminus \Omega_{R/2})}.$$

Also, letting $\psi = \psi_{\sigma_m} - \psi_{\sigma_m}(R)$ and using Equation (3.3),

$$\begin{aligned} \|\sigma_m - \mathbf{U}_R \sigma_m\|_{H^1(\Omega_R)} &= \|\nabla^\perp \psi - \nabla^\perp(h_R \psi)\|_{H^1(\Omega_R)} \\ &\leq \|(1 - h_R)\sigma_m\|_{H^1(\Omega_R)} + \|\nabla^\perp h_R \psi\|_{H^1(\Omega_R)} \\ &\leq \|1 - h_R\|_{C^1} \|\sigma_m\|_{H^1(\Omega_R \setminus \Omega_{R-1})} + \|\nabla h_R\|_{C^1} \|\psi\|_{H^1(\Omega_R \setminus \Omega_{R-1})} \\ &\leq C \|\sigma_m\|_{H^1(\Omega_R \setminus \Omega_{R-1})} + C \|\psi\|_{H^1(\Omega_R \setminus \Omega_{R-1})} \\ &\leq C \|\sigma_m\|_{H^1(\Omega_R \setminus \Omega_{R-1})} + C \|\psi\|_{L^2(\Omega_R \setminus \Omega_{R-1})}. \end{aligned}$$

Because $\Omega_R \setminus \Omega_{R-1}$ has width 1 and ψ vanishes on its outer boundary, we can apply Poincaré's inequality with a constant that is independent of R to give

$$\|\psi\|_{L^2(\Omega_R \setminus \Omega_{R-1})} \leq C \|\nabla \psi\|_{L^2(\Omega_R \setminus \Omega_{R-1})} = C \|\sigma_m\|_{L^2(\Omega_R \setminus \Omega_{R-1})}.$$

Thus,

$$\|\sigma_m - \mathbf{U}_R \sigma_m\|_{H^1(\Omega_R)} \leq C \|\sigma_m\|_{H^1(\Omega_R \setminus \Omega_{R-1})} = C |m| \|\sigma_1\|_{H^1(\Omega_R \setminus \Omega_{R-1})},$$

which vanishes as $R \rightarrow \infty$ by Equation (2.5). This gives Equation (3.4) for \mathbf{U}_R . Projection into V_R gives the closest element in V_R , so Equation (3.4) holds for \mathbf{P}_{V_R} .

Equation (3.5) follows easily:

$$\|\mathbf{P}_{V_R}(u - \sigma_m)\|_{V_R} \leq \|u - \sigma_m\|_{V_R} \leq \|u - \sigma_m\|_V,$$

the first inequality holding simply because \mathbf{P}_{V_R} is an orthogonal projection operator.

To prove Equation (3.6), let $u = \sigma_m + v$ in $E_m \cap \dot{H}^1(\mathbb{R}^2)$. Then

$$\begin{aligned} \|\mathbf{P}_{V_R} u\|_{V_R} &\leq \|\mathbf{P}_{V_R} \sigma_m\|_{V_R} + \|\mathbf{P}_{V_R} v\|_{V_R} \leq \|\sigma_m\|_{V_R} + \|v\|_{V_R} \\ &\leq \|\sigma_m\|_V + \|v\|_V = |m| \|\sigma_1\|_V + \|v\|_V = C |m| + \|v\|_V \\ &\leq \|u\|_{\mathbb{E}^1} + C |m|. \end{aligned}$$

□

Lemma 3.3. $\mathbf{P}_{V_R} \sigma_m$ is a stationary solution to the Euler equations on Ω_R .

Proof. Let $\bar{\sigma}_m = \mathbf{P}_{V_R} \sigma_m$. Since ψ_{σ_m} and $\omega(\sigma_m)$ are radially symmetric, so too must $\psi_{\bar{\sigma}_m}$ and $\omega(\bar{\sigma}_m)$ be. But then

$$\omega(\bar{\sigma}_m \cdot \nabla \bar{\sigma}_m) = \bar{\sigma}_m \cdot \nabla \omega(\bar{\sigma}_m) = 0,$$

and thus $\bar{\sigma}_m \cdot \nabla \bar{\sigma}_m = \nabla p$ for some scalar field p . □

Remark 3.4. Equations (3.4) through (3.6) continue hold if the projection operator \mathbf{P}_{V_R} is replaced by the approximate projection operator \mathbf{U}_R , though a constant factor is introduced on the right-hand sides of Equation (3.5) and Equation (3.6). Equations (3.4) through (3.6) also hold with \mathbf{P}_{V_R} replaced by \mathbf{U}_R and H^1 replaced by L^2 . This gives control not only on the H^1 -norm but individual control on the L^2 -norm. See Remarks 5.3 and 5.6.

We will use the operator \mathbf{P}_{V_R} in establishing the deterministic expanding domain limit in Section 5 and in constructing statistical solutions to (NS) in Section 7. For solutions to the Euler equations, we will need the following approximate truncation operator of Definition 3.5 (or we could use projection into $Y(\Omega_R)$).

Definition 3.5. Define $\bar{\mathbf{U}}_R: \mathbb{Y} \rightarrow \mathbb{Y}(\Omega_R)$ by

$$\bar{\mathbf{U}}_R(\sigma_m + v) = \sigma_m|_{\Omega_R} + T_R v,$$

where $T_R: \mathbb{Y}_0 \rightarrow \mathbb{Y}(\Omega_R)$ is the operator in Lemma 4.2 of [9].

Because Ω_R is a disk, $\sigma_m|_{\Omega_R}$ is in H_R and so also in $\mathbb{Y}(\Omega_R)$. This is why $\bar{\mathbf{U}}_R: \mathbb{Y} \rightarrow \mathbb{Y}(\Omega_R)$. It is also why if we define \mathbf{P}_{H_R} to be the restriction to Ω_R followed by projection into H_R that for all u in \mathbb{E} ,

$$\mathbf{P}_{H_R} u = \sigma(u) + \mathbf{P}_{H_R}(u - \sigma(u)).$$

It follows from Lemma 4.2 of [9] that for all u in \mathbb{E} ,

$$\|\mathbf{P}_{H_R} u - u\|_{L^2(\Omega_R)} \rightarrow 0 \text{ as } R \rightarrow \infty. \quad (3.8)$$

Actually, Lemma 4.2 of [9] applies to an approximate projection operator into H_R , but we are using, as in in [10, 7], the fact that projection into H_R gives the closest vector field in H_R to the vector field being projected.

4. WEAK DETERMINISTIC SOLUTIONS

Definition 4.1 (Weak Navier-Stokes Solution). Given viscosity $\nu > 0$, initial velocity u_0 in H_R , and forcing f in $L^2_{loc}([0, \infty), H_R)$, u in $L^2([0, T]; V_R)$ with $\partial_t u$ in $L^2([0, T]; V'_R)$ is a weak solution to the Navier-Stokes equations on Ω_R if $u(0) = u_0$ and

$$(NS) \quad \int_{\Omega_R} \partial_t u \cdot v + \int_{\Omega_R} (u \cdot \nabla u) \cdot v + \nu \int_{\Omega_R} \nabla u \cdot \nabla v = (u, f)$$

for almost all t in $[0, T]$ and for all v in V_R . A weak solution on \mathbb{R}^2 is defined for u_0 in E_m with u lying in $L^2([0, T]; \dot{H}^1)$ and $\partial_t u$ in $L^2([0, T]; V')$, and with (NS) holding for all v in V .

Definition 4.2 (Weak Euler Solution). Given an initial velocity u_0 in $\mathbb{Y}(\Omega_R)$ and forcing f in $L^2_{loc}([0, \infty), H_R)$, u in $L^\infty([0, T]; H_R \cap H^1(\Omega_R))$ with $\partial_t u$ in $L^2([0, T]; V'_R)$ is a weak solution to the Euler equations if $u(0) = u_0$ and

$$(E) \quad \int_{\Omega_R} \partial_t u \cdot v + \int_{\Omega_R} (u \cdot \nabla u) \cdot v = (f, v)$$

for almost all t in $[0, T]$ and for all v in $H_R \cap H^1(\Omega_R)$. A weak solution on \mathbb{R}^2 is defined for u_0 in \mathbb{Y}_m with u lying in $L^\infty([0, T]; \mathbb{Y}_m)$ and $\partial_t u$ in $L^2([0, T]; V')$, and with (E) holding for all v in V .

Note that the test functions always have finite energy, even for solutions in the whole plane. (Test functions in E_m would be too large to define the integrals involving the nonlinear terms in (NS) and (E).)

Given a solution to (NS), there exists a distribution p (tempered, if the solution is in the whole plane) such that

$$\partial_t u + u \cdot \nabla u + \nabla p = \nu \Delta u + f, \quad (4.1)$$

equality holding in the sense of distributions. This follows from a result of Poincaré and de Rham that any distribution that is a curl-free vector is the gradient of some scalar distribution.

Given a solution to (E), there exists a pressure p such that

$$\partial_t u + u \cdot \nabla u + \nabla p = f, \quad (4.2)$$

but we can only interpret p as a distribution when working in the whole plane. Otherwise, we must view $\partial_t u + u \cdot \nabla u$ as lying in $H^{-1}(\Omega_R)$ and p as lying in $L^2(\Omega_R)$. (Equation (4.2) follows, for instance, from Remark I.1.9 p. 14 of [12].)

In both Equation (4.1) and Equation (4.2) the pressure is unique up to the addition of a function of time. We resolve this ambiguity on Ω_R by requiring that $\int_{\Omega_R} p(t) = 0$ and on \mathbb{R}^2 by requiring that $p(t)$ lie in $L^2(\Omega_R)$ for almost all t in $[0, T]$.

In referring to solutions on Ω_R we will say solutions for R in $[1, \infty)$ and in referring to solutions on \mathbb{R}^2 we will say solutions for $R = \infty$.

Theorem 4.3. (1) *Assume that u_0 is in $E_m \cap \dot{H}^1$. There exists a unique weak solution (u, p) to (NS) in the sense of Definition 4.1 with initial velocity u_0 for $R = \infty$ and initial velocity $\mathbf{P}_{V_R} u_0$ for R in $[1, \infty)$, with*

$$u - \sigma_m \in L^\infty([0, T]; H_R), \quad \nabla u \in L^\infty([0, T]; L^2(\Omega_R)). \quad (4.3)$$

Moreover, there is a bound on each of these norms that is independent of R in $[1, \infty)$ and that depends continuously on $\|u_0\|_{\mathbb{E}^1}$. For $R = \infty$, if u_0 is in $E_m \cap \dot{H}^1$ and $\omega(f)$ is in $L^1([0, T]; L^{p_0} \cap L^\infty)$ then $\omega(u)$ is in $L^\infty([0, T]; L^{p_0} \cap L^\infty)$.

(2) *Assume that u_0 is in \mathbb{Y}_m and that $\omega(f)$ is in $L^1([0, T]; L^{p_0} \cap L^\infty)$. There exists a unique weak solution (u, p) to (E) in the sense of Definition 4.2 with initial velocity u_0 for $R = \infty$ and initial velocity $\bar{\mathbf{U}}_{R} u_0$ for R in $[1, \infty)$. We have,*

$$\begin{aligned} u - \sigma_m &\in L^\infty([0, T]; H_R), & \nabla u &\in L^\infty([0, T]; L^2(\Omega_R)), \\ u &\in L^\infty([0, T] \times \Omega_R), & u &\in C([0, T] \times \bar{\Omega}_R), \\ \partial_t u &\in L^\infty([0, T]; H_R), & \nabla p &\in L^\infty([0, T]; L^2(\Omega_R)), \\ \omega(u) &\in L^\infty([0, T]; L^{p_0} \cap L^\infty(\Omega_R)), \end{aligned}$$

and there is a bound on each of these norms that is independent of R in $[1, \infty)$ and that depends continuously on $\|u_0\|_{\mathbb{Y}}$.

Proof. These results are standard for $R < \infty$, except for the independence of the norms on R . The independence of the first norm in Equation (4.3)

follows for solutions to (NS) from the energy inequality in Equation (5.4) along with Equation (3.4); for the second norm it follows from adapting slightly the proof of this same fact for finite energy in [9]. The stronger bounds for solutions to (E) follow from the vorticity equation for (E) . The results for $R = \infty$ are a minor modification of the same results for finite-energy: see, for instance, [2]. \square

Definition 4.4 (Solution operators). Fix f_R in $L^2([0, \infty); H_R)$ and write f for f_∞ . Let $S_R(t)$ be the solution operator for (NS) on Ω_R with $S = S_\infty$, and let $\bar{S}_R(t)$ be the solution operator for (E) on Ω_R with $\bar{S} = \bar{S}_\infty$.

In application, we will often start with f in $L^2([0, \infty); H)$ or even time-independent f in H and let $f_R = \mathbf{P}_{H_R} f$.

$S_R(0)$ is the identity operator, as is $\bar{S}_R(0)$. For all $t > 0$, S_R maps $H_R \rightarrow V_R$ and $V_R \rightarrow V_R$ and S maps $E_m \rightarrow E_m \cap \dot{H}^1$, $E_m \cap \dot{H}^1 \rightarrow E_m \cap \dot{H}^1$, $\mathbb{E} \rightarrow \mathbb{E}^1$, and $\mathbb{E}^1 \rightarrow \mathbb{E}^1$. For all $t \geq 0$, \bar{S}_R maps $Y(\Omega_R) \rightarrow Y(\Omega_R)$ and \bar{S} maps $\mathbb{Y}_m \rightarrow \mathbb{Y}_m$ and $\mathbb{Y} \rightarrow \mathbb{Y}$. Each of these maps is continuous.

Observe that $S_R(t)\mathbf{P}_{V_R}u_0 = u(t)$ for R in $[1, \infty)$ and $S(t)u_0 = u(t)$ for $R = \infty$ in part (1) of Theorem 4.3, while $\bar{S}_R(t)\bar{\mathbf{U}}_R u_0 = u(t)$ for R in $[1, \infty)$ and $\bar{S}(t)u_0 = u(t)$ for $R = \infty$ in part (2).

5. DETERMINISTIC EXPANDING DOMAIN LIMIT

First we establish the basic energy equality for deterministic solutions to (NS) in Ω_R and in all of \mathbb{R}^2 . In all of \mathbb{R}^2 , the energy is not finite, so we need to subtract σ_m from the velocity to produce an “energy” equality. To make these estimates uniform over R in $[1, \infty]$, we need, then, to subtract σ_m from the velocity for $R < \infty$ as well. Actually, it will be slightly more convenient to subtract

$$\bar{\sigma}_m = \mathbf{P}_{V_R}\sigma_m \text{ for } R \in [1, \infty), \quad \bar{\sigma}_m = \sigma_m \text{ for } R = \infty$$

instead, but because $\bar{\sigma}_m \rightarrow \sigma_m$ in the $H^1(\Omega_R)$ -norm as $R \rightarrow \infty$ by Equation (3.4), this amounts to the same thing.

Theorem 5.1. *Let u be a solution to (NS) as in Definition 4.1 with initial velocity u_0 in H_R for $R < \infty$ and u_0 in E_m for $R = \infty$. Then*

$$\begin{aligned} & \| (u - \bar{\sigma}_m)(t) \|_{L^2(\Omega_R)}^2 + 2\nu \int_0^t \| \nabla(u - \bar{\sigma}_m) \|_{L^2(\Omega_R)}^2 \\ &= \| u_0 - \bar{\sigma}_m \|_{L^2(\Omega_R)}^2 - 2 \int_0^t \int_{\Omega_R} ((u - \bar{\sigma}_m) \cdot \nabla \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m) \\ & \quad - 2\nu \int_0^t \int_{\Omega_R} \nabla \bar{\sigma}_m \cdot \nabla(u - \bar{\sigma}_m) + 2 \int_0^t \int_{\Omega} f_R \cdot (u - \bar{\sigma}_m) \end{aligned} \quad (5.1)$$

and

$$\begin{aligned} & \| (u - \bar{\sigma}_m)(t) \|_{L^2(\Omega_R)}^2 + \nu \int_0^t \| \nabla(u - \bar{\sigma}_m) \|_{L^2(\Omega_R)}^2 \\ & \leq \left(\| u_0 - \bar{\sigma}_m \|_{L^2(\Omega_R)}^2 + Cm^2 \nu t + \| f \|_{L^1([0,t]; L^2)}^2 \right)^{1/2} e^{(Cm+1)t}, \end{aligned} \quad (5.2)$$

where C depends only on $\| \nabla \sigma_1 \|_{L^2 \cap L^\infty}$.

Proof. Assume first that $R < \infty$. Using $u - \bar{\sigma}_m$, which is in V_R for all $t > 0$, as a test function in (NS) gives

$$\begin{aligned} & \int_{\Omega_R} \partial_t u \cdot (u - \bar{\sigma}_m) + \int_{\Omega_R} (u \cdot \nabla u) \cdot (u - \bar{\sigma}_m) + \nu \int_{\Omega_R} \nabla u \cdot \nabla(u - \bar{\sigma}_m) \\ & = \int_{\Omega_R} f \cdot (u - \bar{\sigma}_m). \end{aligned}$$

But,

$$\int_{\Omega_R} \partial_t u \cdot (u - \bar{\sigma}_m) = \int_{\Omega_R} \partial_t (u - \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m) = \frac{1}{2} \frac{d}{dt} \| u - \bar{\sigma}_m \|_{L^2(\Omega_R)}^2$$

and

$$\begin{aligned} & \int_{\Omega_R} (u \cdot \nabla u) \cdot (u - \bar{\sigma}_m) = \int_{\Omega_R} (u \cdot \nabla(u - \bar{\sigma}_m)) \cdot (u - \bar{\sigma}_m) \\ & \quad + \int_{\Omega_R} (u \cdot \nabla \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m) \\ & = \frac{1}{2} \int_{\Omega_R} u \cdot \nabla |u - \bar{\sigma}_m|^2 + \int_{\Omega_R} ((u - \bar{\sigma}_m) \cdot \nabla \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m) \\ & \quad + \int_{\Omega_R} (\bar{\sigma}_m \cdot \nabla \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m). \end{aligned}$$

The first integral in the right-hand side above is formally zero because $\operatorname{div} u = 0$ and $u \cdot \mathbf{n} = 0$ on $\partial\Omega_R$. More properly, we first observe that the integral is finite. This is because at time t in $[0, T]$ both u and $u - \bar{\sigma}_m$ are in $H^1(\Omega_R)$ and so in $L^4(\Omega_R)$ by Sobolev embedding. But $|\nabla |u - \bar{\sigma}_m|^2| \leq 2|u - \bar{\sigma}_m| |\nabla(u - \bar{\sigma}_m)|$, and applying Hölder's inequality gives the finiteness of the integral. Approximating by smooth functions and using the dominated convergence theorem shows that the integral is zero. (This is the approach of Lemmas II.1.1 and II.1.3 p. 108-109 of [12].) Since by Lemma 3.3, $\bar{\sigma}_m \cdot \nabla \bar{\sigma}_m$ is a gradient, the last integral above vanishes. We conclude that

$$\int_{\Omega_R} (u \cdot \nabla u) \cdot (u - \bar{\sigma}_m) = \int_{\Omega_R} ((u - \bar{\sigma}_m) \cdot \nabla \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m).$$

For the final term, we observe that

$$\begin{aligned} & \int_{\Omega_R} \nabla u \cdot \nabla(u - \bar{\sigma}_m) \\ & = \int_{\Omega_R} \nabla(u - \bar{\sigma}_m) \cdot \nabla(u - \bar{\sigma}_m) + \int_{\Omega_R} \nabla \bar{\sigma}_m \cdot \nabla(u - \bar{\sigma}_m). \end{aligned}$$

Combining all these equalities gives

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u - \bar{\sigma}_m\|_{L^2(\Omega_R)}^2 + \nu \|\nabla(u - \bar{\sigma}_m)\|_{L^2(\Omega_R)}^2 \\ &= - \int_{\Omega_R} ((u - \bar{\sigma}_m) \cdot \nabla \bar{\sigma}_m) \cdot (u - \bar{\sigma}_m) - \nu \int_{\Omega_R} \nabla \bar{\sigma}_m \cdot \nabla(u - \bar{\sigma}_m) \\ & \quad + \int_{\Omega_R} f_R \cdot (u - \bar{\sigma}_m). \end{aligned}$$

Integrating in time gives Equation (5.1). Also, we can bound the right-hand side by

$$\begin{aligned} & \|\nabla \bar{\sigma}_m\|_{L^\infty(\mathbb{R}^2)} \|u - \bar{\sigma}_m\|_{L^2(\Omega_R)}^2 + \nu \|\nabla \bar{\sigma}_m\|_{L^2(\mathbb{R}^2)} \|\nabla(u - \bar{\sigma}_m)\|_{L^2(\Omega_R)} \\ & \quad + \|f_R\|_{L^2(\Omega_R)} \|u - \bar{\sigma}_m\|_{L^2(\Omega_R)} \\ & \leq Cm \|u - \bar{\sigma}_m\|_{L^2(\Omega_R)}^2 + Cm^2 \nu + \frac{1}{2} \|f_R\|_{L^2(\Omega_R)}^2 + \frac{\nu}{2} \|\nabla(u - \bar{\sigma}_m)\|_{L^2(\Omega_R)}^2, \end{aligned}$$

where we used Young's inequality and $\bar{\sigma}_m = m\sigma_1$. Thus,

$$\begin{aligned} & \frac{d}{dt} \|u - \bar{\sigma}_m\|_{L^2(\Omega_R)}^2 + \nu \|\nabla(u - \bar{\sigma}_m)\|_{L^2(\Omega_R)}^2 \\ & \leq Cm^2 \nu + \|f_R\|_{L^2(\Omega_R)}^2 + C \|u - \bar{\sigma}_m\|_{L^2(\Omega_R)}^2. \end{aligned} \tag{5.3}$$

Integrating in time and applying Gronwall's inequality gives Equation (5.2).

The energy argument above works equally as well when $R = \infty$ with the exception of the term $(1/2) \int_{\mathbb{R}^2} u \cdot \nabla |u - \bar{\sigma}_m|^2$, which must be handled slightly differently, because at time t in $[0, T]$ we no longer have u in $H^1(\Omega_R)$, only in $\dot{H}^1(\mathbb{R}^2)$. So we divide the integral in two, writing

$$\begin{aligned} \frac{1}{2} \int_{\mathbb{R}^2} u \cdot \nabla |u - \bar{\sigma}_m|^2 &= \frac{1}{2} \int_{\mathbb{R}^2} (u - \bar{\sigma}_m) \cdot \nabla |u - \bar{\sigma}_m|^2 \\ & \quad + \frac{1}{2} \int_{\mathbb{R}^2} \bar{\sigma}_m \cdot \nabla |u - \bar{\sigma}_m|^2. \end{aligned}$$

The first integral is finite and, in fact, zero, using the same reasoning as with the similar term for $R < \infty$. The vector field $\bar{\sigma}_m$ is in $L^\infty(\mathbb{R}^2)$ and $\nabla |u - \bar{\sigma}_m|^2$ is in $L^1(\mathbb{R}^2)$ since $|\nabla |u - \bar{\sigma}_m|^2| \leq 2|u - \bar{\sigma}_m| |\nabla(u - \bar{\sigma}_m)|$; hence, the second integral is finite. We then have

$$\begin{aligned} & \left| \frac{1}{2} \int_{\mathbb{R}^2} \bar{\sigma}_m \cdot \nabla |u - \bar{\sigma}_m|^2 \right| = \lim_{R \rightarrow \infty} \left| \frac{1}{2} \int_{\Omega_R} \bar{\sigma}_m \cdot \nabla |u - \bar{\sigma}_m|^2 \right| \\ & = \lim_{R \rightarrow \infty} \left| \frac{1}{2} \int_{\partial\Omega_R} (\bar{\sigma}_m \cdot \mathbf{n}) |u - \bar{\sigma}_m|^2 \right|. \end{aligned}$$

This integral vanishes, since $\bar{\sigma}_m \cdot \mathbf{n} = 0$ on $\partial\Omega_R$. \square

Corollary 5.2. *Let f lie in $L^2([0, \infty); H)$ and let $f_R = \mathbf{P}_{H_R} f$. Let u_0 be in $E_m \cap \mathbb{E}^1$ and let u be a solution to (NS) as in Definition 4.1 with initial*

velocity $\mathbf{P}_{V_R}u_0$ when R is in $[1, \infty)$ and initial velocity u_0 when $R = \infty$. Then for a constant C independent of R and u_0 ,

$$\begin{aligned} & \|(u - \bar{\sigma}_m)(t)\|_{L^2(\Omega_R)}^2 + \nu \int_0^t \|\nabla(u - \bar{\sigma}_m)\|_{L^2(\Omega_R)}^2 \\ & \leq \left(\|u_0 - \sigma_m\|_{V_R}^2 + Cm^2\nu t + \|f\|_{L^1([0,t];L^2)}^2 \right)^{1/2} e^{(Cm+1)t}. \end{aligned} \quad (5.4)$$

Proof. Apply Theorem 5.1 with initial velocity $\mathbf{P}_{V_R}u_0 = \mathbf{P}_{V_R}(u_0 - \sigma_m) + \mathbf{P}_{V_R}\sigma_m$ and use Equation (3.5). \square

Remark 5.3. Were we to use an initial velocity of $\mathbf{U}_R u_0$ instead of $\mathbf{P}_{V_R}u_0$ in Corollary 5.2 we could replace $\|u_0 - \sigma_m\|_{V_R}$ on the right-hand side of Equation (5.4) with $C\|u_0 - \sigma_m\|_{H_R}$. See Remark 3.4.

We can control the decay of the tail of solutions to (NS) at time t based, ultimately, on their decay at time zero:

Lemma 5.4. For all u_0 in \mathbb{E}^1 ,

$$\|S(t)u_0 - \sigma(S(t)u_0)\|_{L^\infty([0,T];L^2(\Omega_R^c))} \rightarrow 0 \text{ as } R \rightarrow \infty$$

and

$$\|S(t)u_0 - \sigma(S(t)u_0)\|_{L^2([0,T];H^1(\Omega_R^c))} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

Proof. This is a minor adaptation of Lemma 7.1 of [9] to account for infinite-energy, and follows by a standard argument. \square

Theorem 5.5. Assume that u_0 lies in \mathbb{E}^1 and let $u_R(t) = S_R(t)\mathbf{P}_{V_R}u_0$ and $u(t) = S(t)u_0$. Then

$$\|u_R - u\|_{L^\infty([0,T];H^1(\Omega_R))} \rightarrow 0 \text{ as } R \rightarrow \infty, \quad (5.5)$$

$$\|\nabla(u_R - u)\|_{L^2([0,T];L^2(\Omega_R))} \rightarrow 0 \text{ as } R \rightarrow \infty, \quad (5.6)$$

and

$$\|F(t, u) - F_R(t, u_R)\|_{L^2([0,T];V_R'(\Omega_R))} \rightarrow 0 \text{ as } R \rightarrow \infty. \quad (5.7)$$

In addition, the supremum over all u_0 in any bounded subset of \mathbb{E}^1 and over all R in $[1, \infty]$ of each of the quantities,

$$\begin{aligned} & \|u_R - \sigma(u)\|_{L^\infty([0,T];L^2(\Omega_R))}, \quad \|\nabla u_R\|_{L^2([0,T];L^2(\Omega_R))}, \\ & \|F_R(t, u_R)\|_{L^2([0,T];V_R'(\Omega_R))} \end{aligned} \quad (5.8)$$

is finite.

Proof. The first two bounds in Equation (5.8) follow from Equation (5.4). Equations (5.5) and (5.6) follow from Theorem 8.1 of [9] extended to infinite energy solutions using Equation (5.4) in place of the standard finite-energy energy bounds.

We now prove Equation (5.7). We have,

$$\begin{aligned} \|F(t, u) - F_R(t, u_R)\|_{V'_R(\Omega_R)} &\leq \|Au(t) - A_R u_R(t)\|_{V'_R(\Omega_R)} \\ &\quad + \|Bu(t) - B_R u_R(t)\|_{V'_R(\Omega_R)} + \|f - f_R\|_{V'_R(\Omega_R)}. \end{aligned}$$

Let v be in V_R with $\|v\|_{V_R} = 1$. Then

$$\begin{aligned} |(Au(t) - A_R u_R(t), v)| &= \nu |(\Delta u(t) - \Delta u_R(t), v)| \\ &= \nu |(\nabla u(t) - \nabla u_R(t), \nabla v)| \leq \nu \|\nabla u(t) - \nabla u_R(t)\|_{L^2(\Omega_R)} \|v\|_{V_R}. \end{aligned}$$

Thus,

$$\|Au(t) - A_R(u_R)(t)\|_{L^2([0,T];V'_R(\Omega_R))} \leq \nu \|\nabla u - \nabla u_R\|_{L^2([0,T];L^2(\Omega_R))},$$

which vanishes as $R \rightarrow \infty$ by Equation (5.6).

For the nonlinear term,

$$\begin{aligned} |(Bu(t) - B_R u_R(t), v)| &= |(u(t) \cdot \nabla u(t) - u_R(t) \cdot \nabla u_R(t), v)| \\ &= |(\operatorname{div}(u(t) \otimes u(t) - u_R(t) \otimes u_R(t)), v)| \\ &= |(u(t) \otimes u(t) - u_R(t) \otimes u_R(t), \nabla v)| \\ &\leq \|u(t) \otimes u(t) - u_R(t) \otimes u_R(t)\|_{L^2(\Omega_R)} \|v\|_{V_R}. \end{aligned}$$

But,

$$\begin{aligned} &\|u(t)^i u(t)^j - u_R(t)^i \otimes u_R(t)^j\|_{L^2(\Omega_R)} \\ &\leq \|u(t)^i (u(t)^j - u_R(t)^j)\|_{L^2(\Omega_R)} + \|(u(t)^i - u_R(t)^i) u_R(t)^j\|_{L^2(\Omega_R)} \\ &\leq C \|u(t)\|_{L^4(\Omega_R)} \|u(t) - u_R(t)\|_{L^2(\Omega_R)} \\ &\leq C \|u(t)\|_{L^2(\Omega_R)}^{1/2} \|\nabla u(t)\|_{L^2(\Omega_R)}^{1/2} \|u(t) - u_R(t)\|_{L^2(\Omega_R)}^{1/2} \\ &\quad \times \|\nabla(u(t) - u_R(t))\|_{L^2(\Omega_R)}^{1/2} \\ &\leq C \|\nabla u(t)\|_{L^2(\Omega_R)}^{1/2} \|\nabla(u(t) - u_R(t))\|_{L^2(\Omega_R)}^{1/2}, \end{aligned}$$

where we used Ladyzhenskaya's inequality (which gives no dependence on R for the constant C). Thus,

$$\begin{aligned} &\|Bu(t) - B_R(u_R)(t)\|_{L^2([0,T];V'_R(\Omega_R))} \\ &\leq C \|\nabla u\|_{L^1([0,T];L^2(\Omega_R))} \|\nabla(u(t) - u_R(t))\|_{L^1([0,T];L^2(\Omega_R))} \\ &\quad Ct \|\nabla u\|_{L^2([0,T];L^2(\Omega_R))} \|\nabla(u(t) - u_R(t))\|_{L^2([0,T];L^2(\Omega_R))}, \end{aligned}$$

which vanishes as $R \rightarrow \infty$ by Equation (5.6).

For the forcing term,

$$\|f - f_R\|_{L^2([0,T];V'_R(\Omega_R))} \leq \|f - f_R\|_{L^2([0,T];H_R(\Omega_R))},$$

which vanishes as $R \rightarrow \infty$ by Equation (3.8).

From these bounds, Equation (5.7) follows.

It remains to establish the last bound in Equation (5.8). But this follows from an argument similar to that we just made to prove Equation (5.7), using the first two bounds in Equation (5.8). \square

Remark 5.6. Together, the limits in Equations (5.5) and (5.6) are called the *expanding domain limit* in [9]. Most of the estimates involved in establishing these limits require only that the initial velocity lie in H (or \mathbb{E} for the infinite-energy extension). The key exception is that the regularity of the pressure is insufficient to complete the argument unless the initial velocity is in V (or \mathbb{E}^1 for the infinite-energy extension).

Had we used \mathbf{U}_R in place of \mathbf{P}_{V_R} in defining the initial velocity, the expanding domain limit would still hold (indeed, this is how the limit was established in [9]). An advantage of using \mathbf{U}_R is that the resulting bound on the rate of convergence is slightly improved, since H^1 -norms are replaced by L^2 -norms in certain constants that appear in the bound. But this is unimportant in our use of the limit, so we preferred to use \mathbf{P}_{V_R} , since it has a more natural definition.

Corollary 5.7. *For all u_0 in \mathbb{E}^1 ,*

$$\|S_R(t)\mathbf{P}_{V_R}u_0 - \mathbf{P}_{V_R}S(t)u_0\|_{L^\infty([0,T];H_R)} \rightarrow 0 \text{ as } R \rightarrow \infty \quad (5.9)$$

and

$$\|S_R(t)\mathbf{P}_{V_R}u_0 - \mathbf{P}_{V_R}S(t)u_0\|_{L^2([0,T];V_R)} \rightarrow 0 \text{ as } R \rightarrow \infty. \quad (5.10)$$

Proof. By Equations (5.5) and (5.6)

$$\lim_{R \rightarrow \infty} \|S_R(t)\mathbf{P}_{V_R}u_0 - S(t)u_0\|_{L^2([0,T];V_R)} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

But by Equation (3.4),

$$\begin{aligned} & \|\mathbf{P}_{V_R}S(t)u_0 - S(t)u_0\|_{L^2([0,T];V_R)} \\ & \leq C \|S(t)u_0 - \sigma(S(t)u_0)\|_{L^2([0,T];H^1(\Omega_R \setminus \Omega_{R/2}))} + T^{1/2} |m(u)| \beta(R) \\ & \leq C \|S(t)u_0 - \sigma(S(t)u_0)\|_{L^2([0,T];H^1(\Omega_{R/2}^C))} + T^{1/2} |m(u)| \beta(R), \end{aligned}$$

which also vanishes as $R \rightarrow \infty$ by Lemma 5.4. Equation (5.10) then follows from the triangle inequality. The proof of Equation (5.9) is similar. \square

Remark 5.8. It is only in the proof of Corollary 5.7 where we directly use the uniform decay over time of the tail of the velocity for solutions to (NS) . It was, however, already used in the extension of the expanding domain limit from finite to infinite energy alluded to in the proof of Theorem 5.5.

Lemma 5.9. *For all u in \mathbb{E}^1 .*

$$\|F(t, u) - F_R(t, \mathbf{P}_{V_R}u)\|_{V'_R(\Omega_R)} \rightarrow 0 \text{ as } R \rightarrow \infty,$$

and the supremum over any bounded subset of \mathbb{E}^1 and over all R in $[1, \infty]$ of

$$\|F_R(t, \mathbf{P}_{V_R}u)\|_{V'_R(\Omega_R)} \quad (5.11)$$

is finite.

Proof. The proof is the same as that of Equation (5.7), with no need to introduce the L^2 -norm over $[0, T]$, and using Lemma 3.2 in place of the bounds in Equations (5.5) and (5.6). \square

6. DEFINITION OF INFINITE-ENERGY STATISTICAL SOLUTIONS

Following [6] p. 264-265, we define a statistical solution to (NS) on Ω_R , first defining the space of test functions.

Definition 6.1. The space \mathcal{T}_R of test functions, $R < \infty$, is the set of all functions $\Phi: H_R \rightarrow \mathbb{R}$ such that $\Phi(u) = \phi((u, g_1), \dots, (u, g_k))$ for some ϕ in $C^1(\mathbb{R}^k)$ and some g_1, \dots, g_k in V_R . The Fréchet derivative of such a Φ is given by

$$\Phi'(u) = \sum_{j=1}^k \partial_j \phi((u, g_1), \dots, (u, g_k)) g_j,$$

which lies in V_R since each g_j is in V_R . When $R = \infty$, we require that each of g_1, \dots, g_k be compactly supported in \mathbb{R}^2 , so that $\Phi: \mathbb{E} \rightarrow \mathbb{R}$, and we also write \mathcal{T} for \mathcal{T}_∞ . This is the same class of test functions as for homogeneous solutions in the whole space (Definition 2.3 p. 278 of [6]).

Observe that because each $\partial_j \phi$ is bounded,

$$\|\Phi'(u)\|_{V_R} \leq C(\Phi) \tag{6.1}$$

for all u in V_R .

For $t \geq 0$ and u in H_R let

$$F_R(t, u) = f_R(t) - \nu A_R u - B_R(u),$$

where A_R is the Stokes operator and B_R is the classical linear operator associated with the nonlinear term in (NS) on Ω_R . (See, for instance, p. 38 of [6].) We also write F for F_∞ , A for A_∞ , and B for B_∞ .

For $R = \infty$, we will assume for simplicity that f is in $L^2_{loc}([0, \infty); H)$; that is, we do not allow infinite forcing.

Definition 6.2 (Statistical solution to (NS)). Assume that μ_0 is a Borel probability measure on H_R . Then a family,

$$\mu = \{\mu_t\}_{t \geq 0},$$

of Borel probability measures on H_R , $R < \infty$, is a statistical solution to (NS) (SSNS) on Ω_R if each of the following is satisfied:

- (1) For all Φ in \mathcal{T}_R and all $t \geq 0$,

$$\begin{aligned} \int_{H_R} \Phi(u) d\mu_t(u) &= \int_{H_R} \Phi(u) d\mu_0(u) \\ &\quad + \int_0^t \int_{H_R} (F_R(s, u), \Phi'(u)) d\mu_s(u) ds. \end{aligned}$$

- (2) For all $t \geq 0$,

$$\begin{aligned} \int_{H_R} \|u\|_{L^2}^2 d\mu_t(u) &+ 2\nu \int_0^t \int_{H_R} \|\nabla u\|_{L^2}^2 d\mu_s(u) ds \\ &= \int_0^t \int_{H_R} (f(s), u) d\mu_s(u) ds + \int_{H_R} \|u\|_{L^2}^2 d\mu_0(u). \end{aligned}$$

(3) The map

$$t \mapsto \int_{H_R} \phi(u) d\mu_t(u)$$

is measurable for all $t \geq 0$ and all ϕ in $C^0(H_R)$.

(4) The map

$$t \mapsto \int_{H_R} \|u\|_{H_R}^2 d\mu_t(u)$$

lies in $L_{loc}^\infty([0, \infty))$.

(5) The map

$$t \mapsto \int_{H_R} \|\nabla u\|_{L^2}^2 d\mu_t(u)$$

lies in $L_{loc}^1([0, \infty))$.

When $R = \infty$, we make two changes in the definition. First, we replace H_R by \mathbb{E} throughout. Second, the energy equality in (2) is replaced by

(2') For all $t \geq 0$,

$$\begin{aligned} & \int_{\mathbb{E}} \|u - \sigma(u)\|_{L^2}^2 d\mu_t(u) + 2\nu \int_0^t \int_{\mathbb{E}} \|\nabla(u - \sigma(u))\|_{L^2}^2 d\mu_s(u) ds \\ &= \int_0^t \int_{\mathbb{E}} (f(s), u - \sigma(u)) d\mu_s(u) ds + \int_{\mathbb{E}} \|u - \sigma(u)\|_{L^2}^2 d\mu_0(u) \\ & \quad - 2 \int_0^t \int_{\mathbb{E}} ((u - \sigma(u)) \cdot \nabla \sigma(u), u - \sigma(u)) d\mu_s(u) ds \\ & \quad - 2\nu \int_0^t \int_{\mathbb{E}} \nabla \sigma(u) \cdot \nabla(u - \sigma(u)) d\mu_s(u) ds, \end{aligned}$$

where $\sigma(u)$ is defined in Equation (2.7).

The following is from Theorems 1.1 and 1.2 Chapter V of [6]:

Theorem 6.3. *Let μ_0 be as in Definition 6.2, $R < \infty$, with kinetic energy*

$$\int_{H_R} \|u\|_{H_R}^2 d\mu_0(u) < \infty$$

and assume that f lies in $L_{loc}^2([0, \infty); H_R)$. There exists a SSNS, μ , as in Definition 6.2. If the support of μ_0 is (H_R, V_R) -bounded as in Definition 2.1 (meaning that the containment in Equation (6.3) holds for $t = 0$) and f in H_R is time-independent then $\mu_t = S_R(t)\mu_0$ for all $t \geq 0$ is a SSNS. Furthermore, this solution is the unique SSNS satisfying Equations (6.2)

through (6.4):

$$t \mapsto \int_{H_R} \varphi(u) d\mu_t(u) \text{ is continuous on } [0, \infty) \text{ for all } \varphi \text{ in } C(H_R^w), \quad (6.2)$$

$$\text{supp } \mu_t \subseteq \left\{ u \in V_R : \|u\|_{H_R} \leq M \right\} \text{ for all } t \geq 0 \text{ for some } M, \quad (6.3)$$

$$\begin{aligned} \int_{H_R} \Psi(t, u) d\mu_t(u) &= \int_{H_R} \Psi(0, u) d\mu_0(u) \\ &+ \int_0^t \int_{H_R} [\Psi'_s(s, u) + (F(u), \Psi'_u(s, u))] d\mu_s(u) ds. \end{aligned} \quad (6.4)$$

H_R^w is the space H_R in the weak topology. In Equation (6.4), equality holds for all Fréchet-differentiable continuous real-valued functions on $[0, \infty) \times V_R$ (see the discussion following Equation V.1.16 in [6] for more details).

For statistical solutions to (E), we consider only solutions in the whole plane. For solutions to (E) there is no term involving the Stokes operator, so we define

$$F(t, u) = f(t) - B(u).$$

Definition 6.4 (Statistical solution to (E) in the plane). Assume that μ_0 is a Borel probability measure on \mathbb{E} . A statistical solution to the Euler equations (SSE) on \mathbb{E} satisfies all the properties of a SSNS in Definition 6.2 for $R = \infty$ except that the terms involving ν in property (2') are eliminated.

7. CONSTRUCTION OF NAVIER-STOKES SOLUTIONS

Let $S(t)$ be the solution operator on \mathbb{E} as in Definition 4.4. Given that we expect the analog of Theorem 6.3 to hold for infinite-energy solutions in \mathbb{R}^2 , we would expect that

$$\mu_t = S(t)\mu_0 \quad (7.1)$$

is the unique SSNS associated to the initial measure μ_0 if we assume that the support of the initial Borel probability measure μ_0 is $(\mathbb{E}, \mathbb{E}^1)$ -bounded as in Definition 2.1. We show that this is, in fact, the case. Our approach will be to use the SSNS on Ω_R and take a limit as $R \rightarrow \infty$ in a careful way to demonstrate that μ_t is a SSNS on all of \mathbb{R}^2 .

We start by defining the initial probability measure μ_0^R on H_R by

$$\mu_0^R(E) = \mu_0(\mathbf{P}_{V_R}^{-1}E)$$

for all Borel measurable subsets E of H_R . Then μ_0^R is a probability measure, for $\mu_0^R(H_R) = \mu_0(\mathbf{P}_{V_R}^{-1}H_R) = \mu_0(\mathbb{E}) = 1$. Since we are treating initial probability distributions supported on \mathbb{E}^1 , we use projection into V_R . When working with SSNSs as weak as those of Definition 6.2, projection into H_R would be used instead (though the limiting argument in that case is considerably more involved).

Similarly, we define the forcing term f_R in F_R by letting

$$f_R = \mathbf{P}_{H_R} f, \quad (7.2)$$

where \mathbf{P}_{H_R} is projection into H_R . For simplicity, we assume that f is time-independent. Then

$$\|f - f_R\|_{H_R} \rightarrow 0 \text{ as } R \rightarrow \infty \quad (7.3)$$

from Lemma 4.2 of [9] and the observation that projection into H_R gives the closest element in H_R .

We let μ^R be the associated SSNS on Ω_R , so that, by Theorem 6.3,

$$\mu_t^R = S_R(t)\mu_0^R,$$

meaning that $S_R(t)\mu_0^R(E) = \mu_0^R(S_R^{-1}(t)E)$ for any Borel measurable subset E of H_R .

Let Φ be in \mathcal{T} as in Definition 6.1. Because each g_j is compactly supported, for all sufficiently large R , we can define a test function Φ_R in \mathcal{T}_R by

$$\Phi_R(v) \stackrel{\text{def}}{=} \phi((v, g_1|_{\Omega_R}), \dots, (v, g_k|_{\Omega_R})) = \Phi(\mathcal{E}_R v) \quad (7.4)$$

for all v in H_R , where $\mathcal{E}_R v$ is extension by zero of v in H_R to all of \mathbb{R}^2 . It follows that for all v in H_R ,

$$\mathcal{E}_R \Phi'_R(v) = \Phi'(\mathcal{E}_R v). \quad (7.5)$$

From now on, we always assume that R is sufficiently large that Equation (7.4) holds.

For all u in \mathbb{E}^1 ,

$$\begin{aligned} & |\Phi_R(\mathbf{P}_{V_R} u) - \Phi(u)| \\ &= |\phi((\mathbf{P}_{V_R} u, g_1|_{\Omega_R}), \dots, (\mathbf{P}_{V_R} u, g_k|_{\Omega_R})) - \phi((u, g_1), \dots, (u, g_k))| \\ &\leq \|\nabla \phi\|_{L^\infty} |(\mathbf{P}_{V_R} u - u, g_1), \dots, (\mathbf{P}_{V_R} u - u, g_k)| \\ &\leq \|\phi\|_{C^1} \|\mathbf{P}_{V_R} u - u\|_{L^2(\Omega_R)} \left(\|g_1\|_H^2 + \dots + \|g_k\|_H^2 \right)^{1/2} \\ &\leq C \|\mathbf{P}_{V_R} u - u\|_{L^2(\Omega_R)}. \end{aligned}$$

Thus from Lemma 3.2,

$$\Phi_R(\mathbf{P}_{V_R} u) \rightarrow \Phi(u) \text{ as } R \rightarrow \infty. \quad (7.6)$$

Similarly, for all u in \mathbb{E}^1 ,

$$\begin{aligned}
 & \|\Phi'_R(\mathbf{P}_{V_R}u) - \Phi'(u)\|_{V_R} \\
 & \leq \sum_{j=1}^k |\partial_j \phi((\mathbf{P}_{V_R}u, g_1), \dots, (\mathbf{P}_{V_R}u, g_k)) - \partial_j \phi((u, g_1), \dots, (u, g_k))| \|g_j\|_{V_R} \\
 & \leq C \sum_{j=1}^k \mu_j (|(\mathbf{P}_{V_R}u, g_1), \dots, (\mathbf{P}_{V_R}u, g_k) - (u, g_1), \dots, (u, g_k)|) \\
 & = C \sum_{j=1}^k \mu_j (|(\mathbf{P}_{V_R}u - u, g_1), \dots, (\mathbf{P}_{V_R}u - u, g_k)|) \\
 & \leq C \sum_{j=1}^k \mu_j \left(\|\mathbf{P}_{V_R}u - u\|_{H_R} \left(\|g_1\|_H^2 + \dots + \|g_k\|_H^2 \right)^{1/2} \right),
 \end{aligned}$$

where μ_j is the modulus of continuity of $\partial_j \phi$. Thus by Lemma 3.2,

$$\|\Phi'_R(\mathbf{P}_{V_R}u) - \Phi'(u)\|_{V_R} \rightarrow 0 \text{ as } R \rightarrow \infty. \quad (7.7)$$

Before proceeding, we mention one logical simplification that we cannot make. It might seem reasonable to try to show that

$$\begin{aligned}
 S(t)\mu_0 &= \lim_{R \rightarrow \infty} \mathbf{P}_{V_R} \circ S_R(t) \circ \mathbf{P}_{V_R}^{-1} \mu_0 \\
 &= \lim_{R \rightarrow \infty} \mu_0 \circ \mathbf{P}_{V_R}^{-1} \circ S_R(t)^{-1} \circ \mathbf{P}_{V_R}
 \end{aligned} \quad (7.8)$$

by showing that equality holds on any Borel measurable set E . We note, however, that if μ_0 is supported on a singleton set $E = \{u_0\}$ in \mathbb{E} , u_0 nonzero, then $\mathbf{P}_{V_R}^{-1} \circ S_R(t)^{-1} \circ \mathbf{P}_{V_R}(S_R(t)u_0)$ will in general never equal u_0 . Thus, the right-hand side above applied to E will evaluate to 0 for all R , while the left-hand side will evaluate to 1.

Observe that Equation (7.8) is equivalent to saying that

$$\int_{\mathbb{E}} \Phi(u) d\mu_t(u) = \lim_{R \rightarrow \infty} \int_{H_R} \Phi(u) d\nu_t^R(u), \quad (7.9)$$

where $\nu_t = \mathbf{P}_{V_R} \mu_t$ for all test functions Φ (which are dense in the set of bounded continuous functions). We will prove instead that

$$\int_{\mathbb{E}} \Phi(u) d\mu_t(u) = \lim_{R \rightarrow \infty} \int_{H_R} \Phi_R(u) d\mu_t^R(u), \quad (7.10)$$

and that similar limits hold for the other integral in Property (1) of Definition 6.2, thus circumventing our difficulty. In this weak sense, the expanding domain limit could be said to hold for statistical solutions.

Theorem 6.3 continues to hold for $R = \infty$ if we impose at the outset the condition that the initial velocity μ_0 is supported in \mathbb{E}^1 . This condition is required to allow us to take advantage of the expanding domain limit and related bounds from Section 5. (See, however, Remark 7.2.) The idea for proving existence is to first assume that the measure has bounded support

in \mathbb{E}^1 , apply the results of Section 5, which require such support, then use the linearity of properties (1)-(5) of Definition 6.2 to drop the boundedness assumption. This leads to Theorem 7.1.

Theorem 7.1. *Let μ_0 be as in Definition 6.2 for $R = \infty$, but supported in \mathbb{E}^1 , and having “energy”*

$$\int_{\mathbb{E}} \|u\|_{\mathbb{E}}^2 d\mu_0(u) < \infty.$$

Assume that f lies in $L^2_{loc}([0, \infty); V)$. There exists a SSNS, μ , as in Definition 6.2. If the support of μ_0 is $(\mathbb{E}, \mathbb{E}^1)$ -bounded as in Definition 2.1 (meaning that Equation (7.12) holds for $t = 0$) and f in V is time-independent then $\mu_t = S(t)\mu_0$ for all $t \geq 0$ is a SSNS. Furthermore, this solution is the unique SSNS satisfying Equations (7.11) through (7.13):

$$t \mapsto \int_{\mathbb{E}} \varphi(u) d\mu_t(u) \text{ is continuous on } [0, \infty) \text{ for all } \varphi \text{ in } C(H^w), \quad (7.11)$$

$$\text{supp } \mu_t \subseteq \{u \in \mathbb{E}^1 : \|u\|_{\mathbb{E}} \leq M(t)\} \text{ for all } t \geq 0, \quad (7.12)$$

$$\begin{aligned} \int_{\mathbb{E}} \Psi(t, u) d\mu_t(u) &= \int_{\mathbb{E}} \Psi(0, u) d\mu_0(u) \\ &+ \int_0^t \int_X [\Psi'_s(s, u) + (F(u), \Psi'_u(s, u))] d\mu_s(u) ds. \end{aligned} \quad (7.13)$$

In Equation (7.12), M is continuous on $[0, \infty)$. In Equation (7.12), equality holds for all Fréchet-differentiable continuous real-valued functions on $[0, \infty) \times V$.

Proof. Existence: Assume first that the initial Borel probability measure μ_0 has bounded support in \mathbb{E}^1 , meaning that

$$\text{supp } \mu_0 \subseteq \{u \in \mathbb{E}^1 : \|u\|_{\mathbb{E}^1} \leq M\}, \text{ for some } M,$$

and define μ_t by Equation (7.1) for $t \geq 0$. By Equation (4.3) it follows that

$$\text{supp } \mu_t \subseteq \{u \in \mathbb{E}^1 : \|u\|_{\mathbb{E}^1} \leq M(t)\} \quad (7.14)$$

for some continuous function M .

In Theorem 5.5, for initial velocity u_0 in \mathbb{E}^1 we defined $u_R(t) = S_R(t)\mathbf{P}_{V_R}u_0$ and $u(t) = S(t)u_0$. In this proof we will be integrating over all initial velocities in \mathbb{E} and calling the initial velocity u , to agree with the notation of [6]. In this notation, Equation (5.8) and Equation (5.11) become

$$\begin{aligned} \|S_R(t)\mathbf{P}_{V_R}u - \sigma(u)\|_{L^\infty([0, T]; L^2(\Omega_R))}, & \quad \|\nabla S_R(t)\mathbf{P}_{V_R}u\|_{L^2([0, T]; L^2(\Omega_R))}, \\ \|F_R(t, S_R(t)\mathbf{P}_{V_R}u)\|_{L^2([0, T]; V'_R(\Omega_R))}, & \quad \|F_R(t, \mathbf{P}_{V_R}S(t)u)\|_{V'_R(\Omega_R)} \end{aligned} \quad (7.15)$$

are bounded on $\text{supp } \mu_0$ uniformly over all R in $[1, \infty]$. This will allow us to apply the dominated convergence theorem in several steps in our proof.

Using Equation (7.4), for all u in \mathbb{E}^1 ,

$$\begin{aligned} \Phi \left(\lim_{R \rightarrow \infty} S_R(t) \mathbf{P}_{V_R} u \right) &= \Phi \left(\lim_{R \rightarrow \infty} \mathcal{E}_R S_R(t) \mathbf{P}_{V_R} u \right) \\ &= \lim_{R \rightarrow \infty} \Phi (\mathcal{E}_R S_R(t) \mathbf{P}_{V_R} u) = \lim_{R \rightarrow \infty} \Phi_R (S_R(t) \mathbf{P}_{V_R} u), \end{aligned} \quad (7.16)$$

where we used the continuity of Φ . Thus we have,

$$\begin{aligned} \int_{\mathbb{E}} \Phi(u) d\mu_t(u) &= \int_{\mathbb{E}} \Phi(S(t)u) d\mu_0(u) \\ &= \int_{\mathbb{E}} \Phi \left(\lim_{R \rightarrow \infty} S_R(t) \mathbf{P}_{V_R} u \right) d\mu_0(u) \\ &= \int_{\mathbb{E}} \lim_{R \rightarrow \infty} \Phi_R (S_R(t) \mathbf{P}_{V_R} u) d\mu_0(u) \\ &= \lim_{R \rightarrow \infty} \int_{\mathbb{E}} \Phi_R (S_R(t) \mathbf{P}_{V_R} u) d\mu_0(u) \\ &= \lim_{R \rightarrow \infty} \int_{H_R} \Phi_R (S_R(t)v) d\mu_0^R(v) \\ &= \lim_{R \rightarrow \infty} \int_{H_R} \Phi_R (u) d\mu_t^R(u) \end{aligned}$$

giving Equation (7.10). The first equality follows from Equation (7.1), since the space of bounded continuous functions is dual to the space of Borel probability measures. The limit in the second equality follows from Theorem 5.5. The third equality follows from Equation (7.16). The fourth equality follows by the dominated convergence theorem, since Φ_R is uniformly bounded over R in $[1, \infty]$ and μ_0 is a finite measure. The fifth equality follows from Lemma 7.3. The sixth and final equality follows in the same way as does the first.

This shows that Equation (7.10) holds for all $t \geq 0$, so if we can show that

$$\begin{aligned} \int_0^t \int_{\mathbb{E}} (F(s, u), \Phi'(u)) d\mu_s(u) ds \\ = \lim_{R \rightarrow \infty} \int_0^t \int_{H_R} (F_R(s, u), \Phi'_R(u)) d\mu_s^R(u) ds \end{aligned} \quad (7.17)$$

then we will have established the first property of Definition 6.2 for μ .

Toward this end,

$$\begin{aligned}
& \int_0^t \int_{\mathbb{E}} (F(s, u), \Phi'(u)) d\mu_s(u) ds \\
&= \int_0^t \int_{\mathbb{E}} (F(s, S(s)u), \Phi'(S(s)u)) d\mu_0(u) ds \\
&= \int_0^t \int_{\mathbb{E}} \lim_{R \rightarrow \infty} (F_R(s, \mathbf{P}_{V_R} S(s)u), \Phi'(S(s)u)) d\mu_0(u) ds \quad (7.18) \\
&= \int_0^t \int_{\mathbb{E}} \lim_{R \rightarrow \infty} (F_R(s, \mathbf{P}_{V_R} S(s)u), \Phi'_R(\mathbf{P}_{V_R} S(s)u)) d\mu_0(u) ds \\
&= \lim_{R \rightarrow \infty} \int_0^t \int_{\mathbb{E}} (F_R(s, \mathbf{P}_{V_R} S(s)u), \Phi'_R(\mathbf{P}_{V_R} S(s)u)) d\mu_0(u) ds.
\end{aligned}$$

The first equality follows from Lemma 7.4. The second equality follows from Lemma 5.9 and Equation (7.15), since $\Phi'(S(s)u)$ is bounded and compactly supported—and so also we can view the pairings as being in either the duality between V and V' or between V_R and V'_R . The third equality follows from Equation (7.7). The fourth equality follows from the dominated convergence theorem using Equation (7.15).

We would like to commute the roles of the projection operator and the solution operator in the right-hand side of Equation (7.18) to allow us to apply Lemma 7.3. To do this, we estimate,

$$\begin{aligned}
D(s, u) &= |(F_R(s, \mathbf{P}_{V_R} S(s)u), \Phi'_R(\mathbf{P}_{V_R} S(s)u)) \\
&\quad - (F_R(s, S_R(s)\mathbf{P}_{V_R}u), \Phi'_R(S_R(s)\mathbf{P}_{V_R}u))| \\
&\leq |(F_R(s, \mathbf{P}_{V_R} S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u), \Phi'_R(\mathbf{P}_{V_R} S(s)u))| \\
&\quad + |(F_R(s, S_R(s)\mathbf{P}_{V_R}u), \Phi'_R(\mathbf{P}_{V_R} S(s)u) - \Phi'_R(S_R(s)\mathbf{P}_{V_R}u))| \\
&\leq \|F_R(s, \mathbf{P}_{V_R} S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R} \|\Phi'_R(\mathbf{P}_{V_R} S(s)u)\|_{V_R} \\
&\quad + \|F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R} \|\Phi'_R(\mathbf{P}_{V_R} S(s)u) - \Phi'_R(S_R(s)\mathbf{P}_{V_R}u)\|_{V_R}.
\end{aligned}$$

Letting

$$h(R, u, s) := \|\Phi'_R(\mathbf{P}_{V_R} S(s)u) - \Phi'_R(S_R(s)\mathbf{P}_{V_R}u)\|_{V_R}, \quad (7.19)$$

we have

$$\begin{aligned}
& \int_0^t \int_{\mathbb{E}} D(s, u) d\mu_0(u) ds \\
&\leq C \int_0^t \int_{\mathbb{E}} \|F_R(s, \mathbf{P}_{V_R} S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R} d\mu_0(u) ds \quad (7.20) \\
&\quad + \int_0^t \int_{\mathbb{E}} \|F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R} h(R, u, s) d\mu_0(u) ds.
\end{aligned}$$

Applying the Cauchy-Schwarz inequality, we can bound the last term by

$$\begin{aligned} & \int_{\mathbb{E}} \|F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{L^2([0,t];V'_R)} \|h(R, u, \cdot)\|_{L^2([0,t])} d\mu_0(u) \\ & \leq C \int_{\mathbb{E}} \|h(R, u, \cdot)\|_{L^2([0,t])} d\mu_0(u), \end{aligned}$$

using Equation (7.15).

By Equation (7.5),

$$\begin{aligned} h(R, u, s) &= \|\Phi'_R(\mathbf{P}_{V_R}S(s)u) - \Phi'_R(S_R(s)\mathbf{P}_{V_R}u)\|_{V_R} \\ &= \|\Phi'(\mathcal{E}_R\mathbf{P}_{V_R}S(s)u) - \Phi'(\mathcal{E}_RS_R(s)\mathbf{P}_{V_R}u)\|_V. \end{aligned}$$

By Equation (6.1), $\|\Phi'(v)\|_V \leq C_0$ for all v in H_R , for some C_0 independent of R . Thus, $h(R, u, \cdot) \leq 2C_0$. Also, because $\Phi': H \rightarrow V$ is continuous it follows from Equation (5.9) that $h(R, u, \cdot) \rightarrow 0$ as $R \rightarrow \infty$ for all u in \mathbb{E}^1 . Hence, for all u in \mathbb{E}^1 ,

$$\|h(R, u, \cdot)\|_{L^2([0,t])}^2 = \int_0^t h(R, u, s)^2 ds \rightarrow 0$$

by the dominated convergence theorem. But then also $\|h(R, u, \cdot)\|_{L^2([0,t])} \leq 2C_0t^{1/2}$ and applying the dominated convergence theorem again gives

$$\int_{\mathbb{E}} \|h(R, u, \cdot)\|_{L^2([0,t])} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

We conclude that the second term on the right-hand side of Equation (7.20) vanishes as $R \rightarrow \infty$.

For the first term in the right-hand side of Equation (7.20),

$$\begin{aligned} & \|F_R(s, \mathbf{P}_{V_R}S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R} \\ & \leq \|F_R(s, \mathbf{P}_{V_R}S(s)u) - F(s, S(s)u)\|_{V'_R} \\ & \quad + \|F(s, S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R}. \end{aligned}$$

Since $d\mu_0(u) ds$ is a finite measure on $[0, t] \times \mathbb{E}$ and the first term on the right-hand side is both bounded and vanishes as $R \rightarrow \infty$ by Lemma 5.9 and Equation (7.15), after being integrated over $[0, t] \times \mathbb{E}$ the first term vanishes as $R \rightarrow \infty$. The $L^2([0, t])$ -norm of the second term on the right-hand side is bounded on the support of μ_0 by Equation (7.15) and vanishes as $R \rightarrow \infty$ by Equation (5.7); applying the Cauchy-Schwarz inequality followed by the dominated convergence theorem shows that

$$\begin{aligned} & \int_0^t \int_{\mathbb{E}} \|F(s, S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{V'_R} d\mu_0(u) ds \\ & \leq t^{1/2} \int_{\mathbb{E}} \|F(s, S(s)u) - F_R(s, S_R(s)\mathbf{P}_{V_R}u)\|_{L^2([0,t];V'_R)} d\mu_0(u) \end{aligned}$$

vanishes as $R \rightarrow \infty$.

We conclude that $D(s, u)$ integrates to zero in the limit as $R \rightarrow \infty$, meaning that

$$\begin{aligned} & \int_0^t \int_{\mathbb{E}} (F(s, u), \Phi'(u)) d\mu_s(u) ds \\ &= \lim_{R \rightarrow \infty} \int_0^t \int_{\mathbb{E}} (F_R(s, S_R(s)\mathbf{P}_{V_R}u), \Phi'_R(S_R(s)\mathbf{P}_{V_R}u)) d\mu_0(u) ds. \end{aligned}$$

Finally, using Lemma 7.3 and Lemma 7.4,

$$\begin{aligned} & \int_0^t \int_{\mathbb{E}} (F_R(s, S_R(s)\mathbf{P}_{V_R}u), \Phi'_R(S_R(s)\mathbf{P}_{V_R}u)) d\mu_0(u) ds \\ &= \int_0^t \int_{H_R} (F_R(s, S_R(s)v), \Phi'_R(S_R(s)v)) d\mu_0^R(v) ds \\ &= \int_0^t \int_{H_R} (F_R(s, v), \Phi'_R(v)) d\mu_s^R(v) ds, \end{aligned}$$

giving Equation (7.17), completing the demonstration that property (1) of Definition 6.2 is satisfied for μ .

The other properties in Definition 6.2 follow more easily, using the dominated convergence theorem and the first two bounds in Equation (5.8). Thus, we have established the existence of a SSNS for $R = \infty$ when the initial probability measure has bounded support in \mathbb{E}^1 . But we can drop this restriction by exploiting the inherent linearity in the definition of a SSNS, as done on p. 318 of [6]. This establishes the existence part of the theorem.

Higher regularity: We now add the assumption that the support of μ_0 is $(\mathbb{E}, \mathbb{E}^1)$ -bounded. Equation (7.11) and Equation (7.12) follow much as did properties (2) through (5) of Definition 6.2. Adding the assumption that f is time-independent, Equation (7.13) follows for $R = \infty$ in the same way it does for $R < \infty$.

Uniqueness: The proof of uniqueness for $R < \infty$ on p. 319-321 of [6] applies with the following two changes: First, in the Galerkin approximation we use a basis for \mathbb{E}^1 in place of the eigenfunctions of the Stokes operator (the spectrum no longer being discrete). Second, we use the energy bound in Equation (5.2) for $R = \infty$ in place of the bound involving the eigenvalue, λ_m , of the Stokes operator. \square

Remark 7.2. It is possible to drop the assumption in Theorem 7.1 that μ_0 is supported in \mathbb{E}^1 and still obtain existence and to weaken the assumption that the support of μ_0 is $(\mathbb{E}, \mathbb{E}^1)$ -bounded to the support of μ_0 being bounded in \mathbb{E} and still obtain uniqueness. The argument relies on using the boundary condition, $u \cdot \mathbf{n} = \omega(u) = 0$ on $\partial\Omega_R$, in place of no-slip boundary conditions in solutions to the Navier-Stokes equations. Such boundary conditions allow one to bound the gradient of the pressure in $L^1([0, T]; L^2(\Omega_R))$ uniformly in R and so obtain the expanding domain limit of Theorem 5.5 assuming only that u_0 lies in \mathbb{E} . Key to bounding the pressure in this way is Lemma 1 of [11].

We used the following two elementary lemmas in the proof of Theorem 7.1. Note that when we say that equality holds between two integrals when the integrands are only Borel measurable, we mean that either both integrals are defined and equal or that both integrals are undefined. We state the lemmas this way because in their application we do not always know a priori that the integrands are integrable.

Lemma 7.3. *For any Borel measurable function f on H_R ,*

$$\int_{\mathbb{E}} f(\mathbf{P}_{V_R} u) d\mu_0(u) = \int_{H_R} f(v) d\mu_0^R(v). \quad (7.21)$$

Proof. First observe that $f \circ \mathbf{P}_{V_R}$ is Borel measurable on \mathbb{E} because \mathbf{P}_{V_R} is Borel measurable (in fact, continuous) and f is Borel measurable, so the left-hand side of Equation (7.21) is well-defined. When $f = \chi_E$, the characteristic function of a Borel measurable subset E of H_R ,

$$\int_{\mathbb{E}} f(\mathbf{P}_{V_R} u) d\mu_0(u) = \mu_0(\mathbf{P}_{V_R}^{-1} E) = \mu_0^R(E) = \int_{H_R} f(v) d\mu_0^R(v).$$

Equation (7.21) then holds for simple functions by linearity, for nonnegative functions by the monotone convergence theorem, and hence for all Borel measurable functions. \square

Lemma 7.4. *For any function f that is Borel measurable on H_R ,*

$$\int_{H_R} f(u) d\mu_t^R(u) = \int_{H_R} f(S_R(t)u) d\mu_0^R(u).$$

When f is Borel measurable on X ,

$$\int_X f(u) d\mu_t(u) = \int_X f(S(t)u) d\mu_0(u).$$

Proof. As in the proof of Lemma 7.3, equality holds for simple functions, then nonnegative functions, then all Borel measurable functions. \square

8. CONSTRUCTION OF EULER SOLUTIONS

We construct infinite-energy statistical solutions to the Euler equations by making a vanishing viscosity argument, using the infinite-energy statistical solutions to the Navier-Stokes equations that we constructed in Section 7.

For initial velocities as in Theorem 7.1, we have the following for SSNSs:

Theorem 8.1. *Assume that the support of the initial velocity μ_0 for a SSNS with $R = \infty$ is bounded in \mathbb{Y} as in Definition 2.1 and that f is time-independent and lies in \mathbb{Y}_0 . Then the SSNS also satisfies*

$$\text{supp } \mu_t \subseteq \{u \in \mathbb{Y} : \|u\|_{\mathbb{Y}} \leq M(t)\}, \quad (8.1)$$

for a continuous function M independent of ν , and for all p in $[p_0, \infty]$,

$$\int_{\mathbb{E}} \|\omega(u)\|_{L^p} d\mu_t(u) \leq \int_{\mathbb{E}} \|\omega(u)\|_{L^p} d\mu_0(u) + \int_0^t \|\omega(f(s))\|_{L^p} ds. \quad (8.2)$$

Proof. It is a standard result that

$$\|\omega(S(t)u)\|_{L^p} \leq \|\omega(u)\|_{L^p} + \int_0^t \|\omega(f(t))\|_{L^p} \quad (8.3)$$

for all u in \mathbb{Y}_0 . To prove it for $p = r/q$ in lowest terms, with r even, one takes the vorticity of Equation (4.1), $\partial_t \omega + u \cdot \nabla \omega = \nu \Delta \omega + \omega(f)$, multiplies both sides by ω^{p-1} , and integrates over space and time formally to give

$$\|\omega(t)\|_{L^p}^p + p(p-1) \int_0^t \|\omega^{p/2-1} \nabla \omega\|_{L^2(\mathbb{R}^2)}^2 = \|\omega_0\|_{L^p}^p + p \int_0^t (\omega(f), \omega^{p-1}).$$

An approximation and smoothing argument is required to establish the equality rigorously, and it then follows for all p in $[p_0, \infty]$ by the continuity of the L^p norm as a function of p . Applying Hölder's inequality gives Equation (8.3).

Now assume that $u = \sigma_m + v$ is in \mathbb{Y}_m . Then $\partial_t u = \partial_t v$ and $\Delta u = \Delta v$ on Ω_1^C , where $\Delta \sigma_m$ vanishes. Thus, the only additional complication in the argument above is the presence of the additional term $(\sigma_m \cdot \nabla \omega, \omega^{p-1}) = (1/p)(\sigma_m, \nabla \omega^p)$. But this vanishes formally by the divergence theorem, since $\sigma_m \cdot \mathbf{n} = 0$ on $\partial \Omega_R$, hence this term need not be accounted for in the approximation and smoothness argument.

Integrating Equation (8.3) over \mathbb{E} gives

$$\int_{\mathbb{E}} \|\omega(S(t)u)\|_{L^p} d\mu_0(u) \leq \int_{\mathbb{E}} \|\omega(u)\|_{L^p} d\mu_0(u) + \int_0^t \|\omega(f(t))\|_{L^p}.$$

(The last term has no dependence on u so the integral over \mathbb{E} disappears, μ_0 being a probability measure.) But $\|\omega(\cdot)\|_{L^p} : \mathbb{Y} \rightarrow [0, \infty)$ is a bounded continuous function on $\text{supp } \mu_0$ so Equation (8.2) follows from $\mu_t = S(t)\mu_0$, and Equation (8.1) follows from Equation (8.2). \square

Theorem 8.2. *Assume that μ_0 is supported in \mathbb{Y} with*

$$\int_{\mathbb{E}} \|u\|_{\mathbb{E}}^2 d\mu_0(u) < \infty,$$

and assume that f is time-independent and lies in \mathbb{Y}_0 . There exists a SSE, μ , as in Definition 6.4. One such solution is $\mu_t = \bar{S}(t)\mu_0$ for all $t \geq 0$, where $\bar{S}(t)$ is the solution operator for the two-dimensional Euler equations in \mathbb{R}^2 as in Definition 4.4. Furthermore, if the support of μ_0 is bounded in \mathbb{Y} as in Definition 2.1 and f is time-independent then this solution satisfies Equation (8.1) for some function M continuous on $[0, \infty)$ and Equation (8.2).

Proof. Assume first that the support of μ_0 is bounded in \mathbb{Y} . Define $\bar{\mu}_t = \bar{S}(t)\mu_0$, and let μ be the unique SSNS for $R = \infty$ given by Theorem 7.1 with the same forcing and initial data as for the Euler equations. Let $\Phi =$

$\phi((u, g_1), \dots, (u, g_k))$ lie in \mathcal{T} . Then g_1, \dots, g_k are in V and

$$\begin{aligned}\Phi'(u) &= \sum_{j=1}^k \partial_j \phi((u, g_1), \dots, (u, g_k)) g_j \in V, \\ \nabla \Phi'(u) &= \sum_{j=1}^k \partial_j \phi((u, g_1), \dots, (u, g_k)) \nabla g_j \in L^2,\end{aligned}$$

with

$$\|\Phi'(u)\|_V \leq C, \quad \|\nabla \Phi'(u)\|_{L^2} \leq C \quad (8.4)$$

for some constant C independent of u in \mathbb{E} .

Now,

$$\int_{\mathbb{Y}} \Phi(u) d\mu_t(u) = \int_{\mathbb{Y}} \Phi(u) d\mu_0(u) + \int_0^t \int_{\mathbb{Y}} (F(s, u), \Phi'(u)) d\mu_s(u) ds$$

so, using $F = f - \nu Au - Bu$ and $\bar{F} = f - Bu$,

$$\begin{aligned}\int_{\mathbb{Y}} \Phi(u) d\mu_t(u) - \int_{\mathbb{Y}} \Phi(u) d\bar{\mu}_0(u) &- \int_0^t \int_{\mathbb{Y}} (\bar{F}(s, u), \Phi'(u)) d\bar{\mu}_s(u) ds \\ &= \int_{\mathbb{Y}} \Phi(u) d(\mu_0 - \bar{\mu}_0)(u) + \int_0^t \int_{\mathbb{Y}} (F(s, u) - \bar{F}(s, u), \Phi'(u)) d\mu_s(u) ds \\ &\quad - \int_0^t \int_{\mathbb{Y}} (Bu, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) ds \\ &= \int_{\mathbb{Y}} \Phi(u) d(\mu_0 - \bar{\mu}_0)(u) - \nu \int_0^t \int_{\mathbb{Y}} (Au, \Phi'(u)) d\mu_s(u) ds \\ &\quad - \int_0^t \int_{\mathbb{Y}} (Bu, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) ds.\end{aligned}$$

But $\mu_0 = \bar{\mu}_0$, so

$$\begin{aligned}\int_{\mathbb{Y}} \Phi(u) d(\mu_t - \bar{\mu}_t)(u) &= -\nu \int_0^t \int_{\mathbb{Y}} (Au, \Phi'(u)) d\mu_s(u) ds \\ &\quad - \int_0^t \int_{\mathbb{Y}} (Bu, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) ds.\end{aligned}$$

We have,

$$(Bu, \Phi'(u)) = (u \cdot \nabla u, \Phi'(u))$$

and

$$(Au, \Phi'(u)) = -(\Delta u, \Phi'(u)) = (\nabla u, \nabla \Phi'(u)),$$

since $\Phi'(u)$ is in V . Thus,

$$\begin{aligned}\int_{\mathbb{Y}} \Phi(u) d(\mu_t - \bar{\mu}_t)(u) &= -\nu \int_0^t \int_{\mathbb{Y}} (\nabla u, \nabla \Phi'(u)) d\mu_s(u) ds \\ &\quad - \int_0^t \int_{\mathbb{Y}} (u \cdot \nabla u, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) ds.\end{aligned}$$

We have,

$$\int_{\mathbb{Y}} (\nabla u, \nabla \Phi'(u)) d\mu_s(u) \leq C \int_{\mathbb{Y}} \|\nabla u\|_{L^2} d\mu_s(u) \leq C,$$

where we used Equation (8.4) followed by Equation (8.2) and the boundedness of the support of μ_0 in \mathbb{Y} . The same bound holds when integrating against $\bar{\mu}_s$. Thus,

$$\begin{aligned} & \left| \int_{\mathbb{Y}} \Phi(u) d(\mu_t - \bar{\mu}_t)(u) \right| \\ & \leq Rvt + \left| \int_0^t \int_{\mathbb{Y}} (u \cdot \nabla u, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) ds \right|, \end{aligned} \quad (8.5)$$

where R is proportional to the right-hand side of Equation (8.2), which we note increases with time unless there is zero forcing.

For any Borel measurable function G on H ,

$$\begin{aligned} \int_{\mathbb{Y}} G(u) d(\mu_s - \bar{\mu}_s)(u) &= \int_{\mathbb{Y}} G(u) d\mu_s(u) - \int_{\mathbb{Y}} G(u) d\bar{\mu}_s(u) \\ &= \int_{\mathbb{Y}} G(S(s)u) d\mu_0(u) - \int_{\mathbb{Y}} G(\bar{S}(s)u) d\bar{\mu}_0(u) \\ &= \int_{\mathbb{Y}} (G(S(s)u_0) - G(\bar{S}(s)u_0)) d\mu_0(u_0) \\ &= \int_{\mathbb{Y}} (G(u(s)) - G(\bar{u}(s))) d\mu_0(u_0). \end{aligned}$$

In the last integral, we are defining $u(t)$ and $\bar{u}(t)$ to be $S(t)u_0$ and $\bar{S}(t)u_0$, respectively. These are the solutions to (NS) and (E) given the initial velocity u_0 . (The support of μ_0 lying in \mathbb{Y} insures that $\bar{S}(t)u_0$ is well-defined and continuous for μ_0 -almost all u_0 .)

Thus,

$$\begin{aligned} & \int_{\mathbb{Y}} (u \cdot \nabla u, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) \\ &= \int_{\mathbb{Y}} [(u(s) \cdot \nabla u(s), \Phi'(u(s))) - (\bar{u}(s) \cdot \nabla \bar{u}(s), \Phi'(\bar{u}(s)))] d\mu_0. \end{aligned}$$

Letting $w = u - \bar{u}$, we have

$$\begin{aligned} & (u \cdot \nabla u, \Phi'(u)) - (\bar{u} \cdot \nabla \bar{u}, \Phi'(\bar{u})) = (u \cdot \nabla w, \Phi'(u)) \\ & + (u \cdot \nabla \bar{u}, \Phi'(u) - \Phi'(\bar{u})) + (w \cdot \nabla \bar{u}, \Phi'(\bar{u})) \\ & = -(u \cdot \nabla \Phi'(u), w) + (u \cdot \nabla \bar{u}, \Phi'(u) - \Phi'(\bar{u})) + (w \cdot \nabla \bar{u}, \Phi'(\bar{u})), \end{aligned}$$

so

$$\begin{aligned}
 & |(u(s) \cdot \nabla u(s), \Phi'(u(s))) - (\bar{u}(s) \cdot \nabla \bar{u}(s), \Phi'(\bar{u}(s)))| \\
 & \leq \|u(s)\|_{L^\infty} \|\nabla \Phi'(u(s))\|_{L^2} \|w(s)\|_H \\
 & \quad + \|u(s)\|_{L^\infty} \|\nabla \bar{u}(s)\|_{L^2} \|\Phi'(u(s)) - \Phi'(\bar{u}(s))\|_H \\
 & \quad + \|w(s)\|_H \|\nabla \bar{u}(s)\|_{L^2} \|\Phi'(\bar{u}(s))\|_{L^\infty}.
 \end{aligned}$$

Now, Equation (8.3) holds for solutions to (E) : it can be derived as for (NS) or by viewing (E) as a non-homogeneous transport equation for the vorticity. Since $\text{supp } \mu_0$ is bounded in \mathbb{Y} , it follows from Equation (8.3) that u and \bar{u} are bounded in the $L^\infty([0, T] \times \mathbb{R}^2)$ -norm uniformly over $\text{supp } \mu_0$, as is $\nabla \bar{u}$ in the $L^\infty([0, T]; L^2)$ -norm. This is discussed more fully in [2] or [8], where it is shown, moreover, that there exists a continuous function $\rho : [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$, nondecreasing in t , with $\rho(0, t) = 0$ for all $t \geq 0$, such that for all $t > 0$,

$$\|w(t)\|_H \leq \rho(\nu, t). \quad (8.6)$$

(For sufficiently small νt , $\rho(\nu, t) = (C\nu t)^{(1/2)e^{-Ct}}$.)

Also,

$$\begin{aligned}
 & \|\Phi'(u(s)) - \Phi'(\bar{u}(s))\|_H \\
 & \leq \sum_{j=1}^k |\partial_j \phi((u(s), g_1), \dots, (u(s), g_k)) - \partial_j \phi((\bar{u}(s), g_1), \dots, (\bar{u}(s), g_k)))| \|g_j\|_H.
 \end{aligned}$$

Now,

$$|(u(s), g_j) - (\bar{u}(s), g_j)| \leq \|w(s)\|_H \|g_j\|_H \leq \rho(\nu, s) \|g_j\|_H,$$

so since each $\partial_j \phi$ is continuous, it follows that

$$\|\Phi'(u(s)) - \Phi'(\bar{u}(s))\|_H \rightarrow 0 \text{ as } \nu \rightarrow 0 \text{ uniformly over } [0, T].$$

Combining all these facts shows that

$$\int_0^t \int_{\mathbb{Y}} (u \cdot \nabla u, \Phi'(u)) d(\mu_s - \bar{\mu}_s)(u) \rightarrow 0 \text{ as } \nu \rightarrow 0$$

and hence that

$$\begin{aligned}
 & \lim_{\nu \rightarrow 0} \int_{\mathbb{Y}} \Phi(u) d\mu_t(u) \\
 & = \int_{\mathbb{Y}} \Phi(u) d\bar{\mu}_0(u) + \int_0^t \int_{\mathbb{Y}} (\bar{F}(s, u), \Phi'(u)) d\bar{\mu}_s(u) ds.
 \end{aligned} \quad (8.7)$$

On the other hand,

$$\begin{aligned}
\lim_{\nu \rightarrow 0} \int_{\mathbb{Y}} \Phi(u) d\mu_t(u) &= \lim_{\nu \rightarrow 0} \int_{\mathbb{Y}} \Phi(S(t)u) d\mu_0(u) \\
&= \int_{\mathbb{Y}} \lim_{\nu \rightarrow 0} \Phi(S(t)u) d\mu_0(u) = \int_{\mathbb{Y}} \Phi(\bar{S}(t)u) d\mu_0(u) \quad (8.8) \\
&= \int_{\mathbb{Y}} \Phi(u) d\bar{\mu}_t(u).
\end{aligned}$$

In the second equality we used the dominated convergence theorem. For the third equality, we used

$$\begin{aligned}
&|\Phi(S(t)u) - \Phi(\bar{S}(t)u)| \\
&= |\phi((S(t)u, g_1), \dots, (S(t)u, g_k)) - \phi((\bar{S}(t)u, g_1), \dots, (\bar{S}(t)u, g_k))| \\
&\leq \|\nabla\phi\|_{L^\infty} |((S(t)u, g_1), \dots, (S(t)u, g_k)) - ((\bar{S}(t)u, g_1), \dots, (\bar{S}(t)u, g_k))| \\
&\leq C |((S(t)u - \bar{S}(t)u, g_1), \dots, (S(t)u - \bar{S}(t)u, g_k))| \\
&\leq C \|S(t)u - \bar{S}(t)u\|_H \leq C\rho(\nu, t) \rightarrow 0 \text{ as } \nu \rightarrow 0,
\end{aligned}$$

the last inequality just being another way of writing Equation (8.6). Hence, the right-hand sides of Equations (8.7) and (8.8) are equal, establishing the first property in Definition 6.4.

Equations (8.1) and (8.2) follow as in the proof of Theorem 8.1.

As in the proof of Theorem 7.1, we can drop the restriction that the support of μ_0 is bounded in \mathbb{Y} by exploiting the inherent linearity in the definition of a SSE, as done on p. 318 of [6]. The remaining properties in Definition 6.4 follow using the dominated convergence theorem in a manner similar to what we did above. \square

The proof of Theorem 8.2 shows that

$$\int_{\mathbb{Y}} \Phi(u) d\mu_t(u) \rightarrow \int_{\mathbb{Y}} \Phi(u) d\bar{\mu}_t(u) \text{ as } \nu \rightarrow 0.$$

Since the space \mathcal{T} of test functions is dense in the space of all bounded continuous functions on \mathbb{E} , it follows that $\mu \rightarrow \bar{\mu}$ as measures as $\nu \rightarrow 0$; that is, the vanishing viscosity limit holds for statistical solutions to the Navier-Stokes and Euler equations.

ACKNOWLEDGEMENTS

The author was supported in part by NSF grant DMS-0705586 during the period of this work.

REFERENCES

- [1] Matania Ben-Artzi. Global solutions of two-dimensional Navier-Stokes and Euler equations. *Arch. Rational Mech. Anal.*, 128(4):329–358, 1994.
- [2] Jean-Yves Chemin. A remark on the inviscid limit for two-dimensional incompressible fluids. *Comm. Partial Differential Equations*, 21(11-12):1771–1779, 1996.

- [3] Jean-Yves Chemin. *Perfect incompressible fluids*, volume 14 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press Oxford University Press, New York, 1998. Translated from the 1995 French original by Isabelle Gallagher and Dragos Iftimie.
- [4] P. Constantin and F. Ramos. Inviscid limit for damped and driven incompressible Navier-Stokes equations in \mathbb{R}^2 . *Comm. Math. Phys.*, 275(2):529–551, 2007.
- [5] C. Foias. Statistical study of Navier-Stokes equations. I, II. *Rend. Sem. Mat. Univ. Padova*, 48:219–348 (1973); *ibid.* 49 (1973), 9–123, 1972.
- [6] C. Foias, O. Manley, R. Rosa, and R. Temam. *Navier-Stokes equations and turbulence*, volume 83 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2001.
- [7] Dragos Iftimie and James P. Kelliher. Remarks on the vanishing obstacle limit for a 3d viscous incompressible fluid. *Proceedings of the American Mathematical Society*, 137(2):685–694, 2009.
- [8] James P. Kelliher. The inviscid limit for two-dimensional incompressible fluids with unbounded vorticity. *Math. Res. Lett.*, 11(4):519–528, 2004.
- [9] James P. Kelliher. Expanding domain limit for incompressible fluids in the plane. *Comm. Math. Phys.*, 278(3):753–773, 2008.
- [10] James P. Kelliher, M. C. Lopes Filho, and H. J. Nussenzweig Lopes. Vanishing viscosity limit for an expanding domain in space. *To appear, Annales de l'Institut Henri Poincaré (C) Analyse Non Linéaire*.
- [11] Namkwon Kim. Large friction limit and the inviscid limit of 2-D Navier-Stokes equations under Navier friction condition. *To appear in SIAM Math Analysis*.
- [12] Roger Temam. *Navier-Stokes equations, theory and numerical analysis*. AMS Chelsea Publishing, Providence, RI, 2001. Reprint of the 1984 edition.
- [13] V. I. Yudovich. Non-stationary flows of an ideal incompressible fluid. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 3:1032–1066 (Russian), 1963.
- [14] V. I. Yudovich. Uniqueness theorem for the basic nonstationary problem in the dynamics of an ideal incompressible fluid. *Math. Res. Lett.*, 2(1):27–38, 1995.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, 900 UNIVERSITY AVE., RIVERSIDE, CA 92521

Current address: Department of Mathematics, University of California, Riverside, 900 University Ave., Riverside, CA 92521

E-mail address: kelliher@math.ucr.edu