TANGENTIAL THICKNESS OF HOMOTOPY LENS SPACES

Sławomir Kwasik Department of Mathematics, Tulane University New Orleans, Louisiana 70118, U. S. A. kwasik@math.tulane.edu

> Reinhard Schultz Department of Mathematics University of California, Riverside Riverside, California 92521, U. S. A. schultz@math.ucr.edu

Given two nonhomeomorphic topological spaces X and Y, it is often interesting or important to specify necessary or sufficient conditions for $X \times \mathbb{R}$ and $Y \times \mathbb{R}$ to be homeomorphic, where as usual \mathbb{R} denotes the real line. More generally, it is also useful to have criteria for determining whether $X \times \mathbb{R}^l$ and $Y \times \mathbb{R}^k$ are homeomorphic for some k > 1. If X and Y are closed manifolds, the following result, which is due to B. Mazur in the smooth and piecewise linear (PL) categories [91, 93], provides an abstract answer. and to R. Kirby and L. Siebenmann in the topological category [51], provides an abstract answer. In this result CAT refer to the category of smooth, piecewise linear or topological manifolds and a CAT-isomorphism is a diffeomorphism, piecewise linear homeomorphism or homeomorphism respectively:

Theorem 1. STABLE EQUIVALENCE THEOREM. Let M and N be closed CAT-manifolds. Then $M \times \mathbb{R}^k$ and $N \times \mathbb{R}^k$ are CAT isomorphic for some $k \ge 1$ if and only if M and N are stably tangentially homotopy equivalent.

For the sake of completeness, we note that two manifolds are stably tangentially homotopy equivalent if and only if they are homotopy equivalent such that the stable tangent bundle of one pulls back to the stable tangent bundle of the other; other words, the direct sum of the tangent bundle with a trivial line bundle on the codomain pulls back to the corresponding bundle on the domain under the homotopy equivalence. As shown in [12], there are pairs of homotopy equivalent manifolds such that the homotopy equivalence is stably tangential but the unstable tangent bundle of the codomain does not pull back to the unstable tangent bundle of the domain.

In fact, if f exists then for some k the map $f \times id(\mathbb{R}^k)$ is properly homotopic to a CATisomorphism; the topological version of this result follows from [?, ?] if [?]

Given two manifolds M and N satisfying the conditions of the stable Equivalence Theorem, it is natural to ask the following:

OPTIMAL VALUE QUESTION. For a given tangential homotopy equivalence $f : M \to N$, what is the least value of k such that $f \times id(\mathbb{R}^k)$ is properly homotopic to a CAT isomorphism?

If n is the common dimension of M and N, the standard embedding and stable tubular neighborhood theorems for CAT-manifolds imply that n + 1 is a universal upper bound for k in each category.

In [56] special cases of the Optional Value Question were considered for linear spherical space forms in the topological category, and in particular it was shown that if M and N are linear

space forms such that $M \times \mathbb{R}^2$ is homeomorphic to $N \times \mathbb{R}^2$, then M and N are homeomorphic but $M \times \mathbb{R}^3$ and $N \times \mathbb{R}^3$ are not diffeomorphic. These results already reflect the relative complexity of this problem.

Our first result concerns a linear lens spaces.

Theorem 2. Let $f : M \to N$ be tangential homotopy equivalence of linear lens spaces with prime order fundamental groups. Then $f \times id_{R^3}$ is properly homotopic to a homeomorphism.

The restriction to linear lens spaces as opposed to homotopy lens spaces is crucial. It is well know that there are fake lens spaces L_1 and L_2 are not homeomorphic but $L_1 \times \mathbb{R}$ and $L_2 \times \mathbb{R}$ are diffeomorphic [?]

In this paper we shall study the Optimal Value Question for homotopy lens spaces. We will concentrate on the case of (odd) prime order fundamental groups, although many of our results hold without this restriction. Qualitatively, one can describe the results in terms of the concept we call **tangential thickness**. Specifically, two CAT manifolds M and N are said to have tangential thickness $\leq k$ if the following holds: Given a manifold M, let $\mathbb{TT}_k(M)$ denote the isomorphism classes of manifolds N such that $\{M, N\}$ has tangential thickness $\leq k$, and let $\mathbf{TT}(M)$ denote the isomorphism classes of manifolds that are stably tangentially homotopy equivalent to M. One then has an increasing sequence of sets $TT_k(M)$:

$$\{\operatorname{class}(M)\} = \mathbf{TT}_0(M) \subset \mathbf{TT}_1(M) \subset \cdots \subset \mathbf{TT}_k(M) \subset \mathbf{TT}(M)$$

The sequence stabilizes for $k \ge \dim M + 1$ by Mazur's result, so that

$$\mathbf{TT}_k(M) = \mathbf{TT}_{k+i}(M) = \mathbf{TT}(M)$$
, $i = 1, 2, 3, \cdots$

In particular, given a manifold M then the classification of all manifolds having the tangential thickness tt(M) = k is equivalent to the computation of the set $\mathbf{TT}_k(M) - \mathbf{TT}_{k-1}(M)$. We are ready now to state results of this paper.

Theorem 3. Let M^{2N-1} , $N \ge 3$ be a homotopy lens space (arbitrary fake spherical space form of dimension ≥ 3). Then $\mathbf{TT}^{TOP}(M^{2n-1})$ consists of manifolds h-cobordant to M^{2n-1} . These manifolds are classified by $Wh(\pi_1(M2n-1))$ via realization of Whitehead torsion by h-cobordant i.e. free torsion of $Wh(\pi_1(M^{2n-1}))$ on M^{2n-1} .

Theorem 4. Let M^{2n-1} , $n \ge 3$ be a fake lens space. Then a manifold N^{2n-1} is in $\mathbf{TT}_2^{TOP}(M^{2n-1})$ if and only if $N^{2n-1} \times R$ is properly h-cobordant to $M^{2n-1} \times R$. The set $\mathbf{TT}_2^{TOP}(M^{2n-1})$ - $\mathbf{TT}_1^{TOP}(M^2n - 1)$ is in one-to-one correspondence with $H^0(K_0Z[\prod_1 M^{2n-1})]$. Moreover all possible manifolds in $\mathbf{TT}_2^{TOP}(M^{2n-1})$ - $\mathbf{TT}_1^{TOP}(M^{2n-1})$ are obtained by a free action of $H_0(K_02[_1)M^{2n-1}]$ on $M^{2n-1} \times R$ as the realization of Whitehead torsion by proper h-cobordisms.

<u>Theorem 3</u>. Let $M^{2n-1}, n \ge 3$ be a fake lens space with $\prod_1 (M^{2n-1}) \cong 2p, p$ -odd prime. The the set

TO BE COMPLETED

References

- D. Acosta and T. Lawson, Even non-spin manifolds, spin^c structures, and duality. Enseign. Math. (2) 43 (1997), 27–32.
- [2] J. F. Adams, On the groups J(X) IV. Topology 4 (1966), 21–71.
- [3] J. F. Adams, Lectures on generalised cohomology. "Category Theory, Homology Theory and Their Applications III (Battelle Institute Conference, Seattle, WA, 1968, Vol. 3)," pp. 1–138. Lecture Notes in Math., 99. Springer-Verlag, Berlin-etc., 1969.
- [4] J. P. Alexander, *The bisection problem*. Ph.D. Dissertation, University of California, Berkeley, 1971.
- [5] M. F. Atiyah, Geometrical aspects of gauge theories. "Proceedings of the International Congress of Mathematicians (Helsinki, 1978)," pp. 881–885, Acad. Sci. Fennica, Helsinki, 1980.
- [6] A. Bak, The computation of surgery groups of finite groups with abelian 2hyperelementary subgroups. "Algebraic K-theory (Proc. Conf., Northwestern Univ., Evanston, Ill., 1976)," pp. 384–409. Lecture Notes in Math., 551. Springer-Verlag, Berlinetc., 1976.
- [7] A. Baker, "Matrix Groups: An Introduction to Lie Group Theory." Springer Undergraduate Mathematics Series. Springer-Verlag, Berlin-etc., 2002.
- [8] D. Barden, On the structure and classification of differential manifolds. Ph.D. Thesis, University of Cambridge, 1964.
- [9] M. Berger, "Geometry I (Transl. from the 1977 French original by M. Cole and S. Levy, corrected reprint of the 1987 transl.)," Universitext. Springer-Verlag, Berlin-etc., 1994.
- [10] R. H. Bing, "The geometric topology of 3-manifolds." Amer. Math. Soc. Colloq. Publ. No. 40. Amer. Math. Soc., Providence, RI, 1983.
- [11] M. Brown, Locally flat imbeddings of topological manifolds. Ann. of Math. 75 (1962), 331–341.
- [12] Y. Byun. Tangent bundle of a manifold and its homotopy type. J. London Math. Soc.
 (2) 60 (1969), 303–307.
- [13] S. Cappell, R. Lashof, and J. Shaneson, A splitting theorem and the structure of 5manifolds. "Symposia Mathematica, Vol. X (Convegno di Geometria Differenziale, IN-DAM, Rome, 1971)," pp. 47–58. Academic Press, London, 1972.
- [14] S. E. Cappell, and J. L. Shaneson, On four dimensional surgery and applications. Comment. Math. Helv. 46 (1971), 500–528.
- [15] S. E. Cappell, and J. L. Shaneson, On 4-dimensional s-cobordisms. J. Diff. Geom. 22 (1985), 97–115.
- [16] A. J. Casson and C. McA. Gordon, On slice knots in dimension three. "Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, CA, 1976), Part 2," pp. 39–53, Proc. Sympos. Pure Math. Vol. XXXII, Part 2, Amer. Math. Soc., Providence, RI, 1978.
- [17] F. R. Cohen, J. C. Moore, and J. A. Neisendorfer, The double suspension and exponents of the homotopy groups of spheres. Ann. of Math. (2) 110 (1979), 549–565.
- [18] F. R. Cohen, J. C. Moore, and J. A. Neisendorfer, Note on higher torsion in the homotopy groups of single suspensions. Illinois J. Math. 24 (1980), 619–623.
- [19] M. M. Cohen, "A course in simple-homotopy theory." Graduate Texts in Mathematics, Vol. 10. Springer-Verlag, Berlin-etc., 1973.
- [20] E. H. Connell, A topological h-cobordism theorem for $n \ge 5$. Illinois J. Math. 11 (1967), 300–309.

- [21] H. S. M. Coxeter, "Regular polytopes (Third Ed.)." Dover Publications, New York, NY, 1973.
- [22] S. K. Donaldson, An application of gauge theory to four-dimensional topology. J. Diff. Geom. 18 (1983), 279–315.
- [23] S. K. Donaldson, Connections, cohomology and the intersection forms of 4-manifolds. J. Diff. Geom. 24 (1986), 275–341.
- [24] S. K. Donaldson, Irrationality and the h-cobordism conjecture. J. Differential Geom. 26 (1987), 141–168.
- [25] S. K. Donaldson, The Seiberg-Witten equations and 4-manifold topology. Bull. Amer. Math. Soc. (2) 33 (1996), 45–70.
- [26] J. Ewing, S. Moolgavkar, L. Smith, and R. E. Stong, Stable parallelizability of lens spaces. J. Pure Appl. Algebra 10 (1977/78), 177–191.
- [27] F. T. Farrell and J. B. Wagoner, Algebraic torsion for infinite simple homotopy types. Comment. Math. Helv. 47 (1972), 502–513.
- [28] R. Fintushel, and T. Lawson, Compactness of moduli spaces for orbifold instantons. Topology Appl. 23 (1986), 305–312.
- [29] R. Fintushel and R. Stern, *Pseudofree orbifolds*. Ann. of Math. 122 (1985), 335–364.
- [30] R. Fintushel and R. Stern, *Definite 4-manifolds*. J. Diff. Geom. 28 (1988), 133–141.
- [31] J. Folkman, "Equivariant maps of spheres into the classical groups" (with a preface by J. Minor). Memoirs of the Amer. Math. Soc., No. 95. American Mathematical Society, Providence, R.I., 1971.
- [32] M. H. Freedman and F. S. Quinn, A quick proof of the 4-dimensional stable surgery theorem. Comment. Math. Helv. 55 (1980), 668–671.
- [33] M. H. Freedman, The topology of four-dimensional manifolds, J. Differential Geom. 17 (1982), 357–453.
- [34] M. H. Freedman and F. S. Quinn, "Topology of 4-manifolds," Princeton Mathematical Series, 39. Princeton University Press, Princeton, NJ, 1990.
- [35] D. Gay, 4-manifolds which are homeomorphic but not diffeomorphic. Expository paper, Riemannian Geometry Seminar, University of California at Berkeley, 1995. Online, available at http://www.mths.uct.ac.za/~dgay/homnotdiff.pdf.
- [36] E. Giroux, What is ... an Open Book? Notices Amer. Math. Soc. 52 (2005), 42–47.
- [37] W. Haken, Some results on surfaces in 3-manifolds. "Studies in Modern Topology," pp. 39 – 98. Mathematical Association of American, distr. by Prentice-Hall, Englewood Cliffs, NJ, 1968.
- [38] A. Hatcher, "Algebraic Topology (Third Paperback Printing)," Cambridge University Press, New York, NY, 2002. Updated version with corrections available online at http:// www.math.cornell.edu/~hatcher/AT/ATpage.html.
- [39] A. Hatcher and T. Lawson, Stability theorems for "concordance implies isotopy" and "h-cobordism implies diffeomorphism". Duke Math. J. 43 (1976), 555–560.
- [40] J. Hempel, "3-manifolds (New Edition)." AMS Chelsea Publications, American mathematical Society, Providence, RI, 2004.
- [41] D. Hilbert and S. Cohn-Vossen, "Geometry and the imagination (Transl. by P. Neményi)." Chelsea Publishing, New York, NY, 1952.
- [42] M. W. Hirsch, "Differential topology. Corrected reprint of the 1976 original." Graduate Texts in Mathematics No. 33. Springer-Verlag, Berlin-etc., 1994.
- [43] N. Hitchin, PROJECTIVE GEOMETRY, b3 course 2003. Maths. Inst., Oxford University, 2003. Online, available at http://www.maths.ox.ac.uk/~hitchin/hitchinnotes/ Projective_geometry/ Chapter_1_Projective_geometry.pdf.

- [44] J. G. Hocking and G. S. Young, "Topology (2nd Ed.)." Dover Publications, New York, 1988.
- [45] R. Holzsager, Stable splitting of K(G, 1). Proc. Amer. Math. Soc. **31** (1972), 305–306.
- [46] W.-C. Hsiang, and R. H. Szczarba, On embedding surfaces in four-manifolds. "Algebraic topology (Proc. Sympos. Pure Math., Vol. XXII, Univ. Wisconsin, Madison, WI, 1970)," pp. 97–103. Amer. Math. Soc., Providence, RI, 1971.
- [47] W. H. Jaco, and P. B. Shalen, Seifert fibered spaces in 3-manifolds. Mem. Amer. Math. Soc. 21 (1979), No. 220.
- [48] K. Johannson, "Homotopy equivalences of 3-manifolds with boundaries." Lecture Notes in Mathematics, 761. Springer-Verlag, Berlin-etc., 1979.
- [49] T. Kambe, The structure of K_{Λ} -rings of the lens spaces and their applications, J. Math. Soc. Japan **18** (1966), 135–146.
- [50] M. A. Kervaire, and J. W. Milnor, On 2-spheres in 4-manifolds. Proc. Nat. Acad. Sci. U. S. A. 47 (1961), 1651–1657.
- [51] R. C. Kirby and L. C. Siebenmann, "Foundational Essays on Topological Manifolds, Smoothings, and Triangulations," Ann. of Math. Studies No. 88. Princeton University Press, Princeton, NJ, 1977.
- [52] R. C. Kirby, (ed.), Problems in low-dimensional topology, "AMS/IP Stud. Adv. Math. 2.2, Geometric topology (Athens, GA, 1993)," Amer. Math. Soc., Providence, RI, 1997, pp. 35–473. (An updated version of this is available online at http://math.berkeley.edu/~kirby/problems.ps.gz.
- [53] F. Krauß and T. Lawson, Examples of homogeneous C^{*}-algebras. "Recent advances in the representation theory of rings and C^{*}-algebras by continuous sections (Sem., Tulane Univ., New Orleans, La., 1973)," pp. 153–164. Mem. Amer. Math. Soc., No. 148, Amer. Math. Soc., Providence, RI, 1974.
- [54] M. Kreck, and W. Lück, "The Novikov conjecture. Geometry and algebra." Oberwolfach Seminars, 33. Birkhäuser Verlag, Basel, 2005.
- [55] S. Kwasik and T. Lawson, Nonsmoothable Z_p actions on contractible 4-manifolds. J. reine angew. Math. 437 (1993), 29–54.
- [56] S. Kwasik and R. Schultz, Toral and exponential stabilization for homotopy spherical spaceforms. Math. Proc. Cambridge Philos. Soc. 137 (2004), 571–593.
- [57] K. Kuga, Representing homology classes of $S^2 \times S^2$. Topology 23 (1984), 133–137.
- [58] S. Lang, "Differential manifolds." Addison-Wesley, Reading, Mass. and London-Don Mills, Ont., 1972.
- [59] H. B. Lawson, *Foliations*. Bull. Amer. Math. Soc. **80** (1974), 369–418.
- [60] T. Lawson, Some examples of nonfinite diffeomorphism groups. Proc. Amer. Math. Soc. 34 (1972), 570–572.
- [61] T. Lawson, Remarks on the pairings of Bredon, Milnor, and Milnor-Munkres-Novikov. Indiana Univ. Math. J. 22 (1972/73), 833–843.
- [62] T. Lawson, Inertial h-cobordisms with finite cyclic fundamental group. Proc. Amer. Math. Soc. 44 (1974), 492–496.
- [63] T. Lawson, Remarks on the four and five dimensional s-cobordism conjectures. Duke Math. J. 41 (1974), 639–644.
- [64] T. Lawson, Splitting isomorphisms of mapping tori. Trans. Amer. Math. Soc. 205 (1975), 285–294.
- [65] T. Lawson, Homeomorphisms of $B^k \times T^n$. Proc. Amer. Math. Soc. 56 (1976), 349–350.
- [66] T. Lawson, Applications of decomposition theorems to trivializing h-cobordisms. Canad. Math. Bull. 20 (1977), 389–391.
- [67] T. Lawson, Trivializing h-cobordisms by stabilization. Math. Z. 156 (1977), 211–215.

- [68] T. Lawson, Open book decompositions for odd dimensional manifolds. Topology 17 (1978), 189–192.
- [69] T. Lawson, Decomposing 5-manifolds as doubles. Houston J. Math. 4 (1978), 81–84.
- [70] T. Lawson, Trivializing 5-dimensional h-cobordisms by stabilization. Manuscr. Math. 29 (1979), 305–321.
- [71] T. Lawson, Splitting spheres as codimension r doubles. Houston J. Math. 8 (1982), 205– 220.
- [72] T. Lawson, Splitting S⁴ on ℝP² via the branched cover of ℂP² over S⁴. Proc. Amer. Math. Soc. 86 (1982), 328–330.
- [73] T. Lawson, Detecting the standard embedding of \mathbb{RP}^2 in S^4 . Math. Ann. **267** (1984), 439–448.
- [74] T. Lawson, Normal bundles for an embedded ℝP² in a positive definite 4-manifold. J. Diff. Geom. 22 (1985), 215–231.
- [75] T. Lawson, Invariants for families of Brieskorn varieties. Proc. Amer. Math. Soc. 99 (1987), 187–192.
- [76] T. Lawson, Representing homology classes of almost definite 4-manifolds. Michigan Math. J. 34 (1987), 85–91.
- [77] T. Lawson, h-cobordisms between simply connected 4-manifolds. Topology Appl. 28 (1988), 75–82.
- [78] T. Lawson, Compactness results for orbifold instantons. Math. Z. 200 (1988), 123–140.
- [79] T. Lawson, Smooth embeddings of 2-spheres in 4-manifolds. Exposition. Math. 10 (1992), 289–309.
- [80] T. Lawson, A note on trigonometric sums arising in gauge theory. Manuscr. Math. 80 (1993), 265–272.
- [81] T. Lawson, The minimal genus problem. Exposition. Math. 15 (1997), 385–431.
- [82] T. Lawson, "Topology: a geometric approach." Oxford Graduate Texts in Mathematics No. 9. Oxford University Press, Oxford, 2003.
- [83] R. Lee, and D. Wilczyński, Locally flat 2-spheres in simply connected 4-manifolds. Comment. Math. Helv. 65 (1990), 388–412.
- [84] M. Mahowald, "The metastable homotopy of S^n ." Memoirs of the Amer. Math. Soc., No. 72. American Mathematical Society, Providence, R.I., 1967.
- [85] M. Mahowald, A new infinite family in $_{2}\pi_{*}^{s}$. Topology 16 (1977), 249–256.
- [86] R. Mandelbaum, Four-dimensional topology: An introduction. Bull. Amer. Math. Soc. (2) 2 (1980), 1–159.
- [87] R. Mandelbaum and B. Moishezon, On the topology of simply-connected algebraic surfaces. Trans. Amer. Math. Soc. 260 (1980), 195–222.
- [88] A. A. Markov, Insolubility of the problem of homeomorphy. (Russian) Proc. Internat. Congress Math. (Edinburgh, 1958) pp. 300–306. Cambridge Univ. Press, New York, 1960.
- [89] A. A. Markov, The insolubility of the problem of homeomorphy (Russian). Dokl. Akad. Nauk SSSR 121 (1958) 218–220.
- [90] W. S. Massey, Proof of a conjecture of Whitney. Pacific J. Math. 31 (1969), 143–156.
- [91] B. Mazur, Stable equivalence of differentiable manifolds. Bull. Amer. Math. Soc. 67 (1961), 377–384.
- [92] B. Mazur, A note on some contractible 4-manifolds. Ann. of Math. (2) 73 (1961), 221– 228.
- [93] B. Mazur, The method of infinite repetition in pure topology. II. Stable applications. Ann. of Math. (2) 83 (1966), 387–401.

- [94] J. Milnor, "Lectures on the *h*-cobordism theorem (Notes by L. Siebenmann and J. Sondow)," Princeton Mathematical Series No. 1. Princeton University Press, Princeton, NJ, 1965.
- [95] J. Milnor and J. Stasheff, "Characteristic Classes," Annals of Mathematics Studies Vol. 76. Princeton Univ. Press, Princeton, NJ, 1974.
- [96] E.E. Moise, "Geometric topology in dimensions 2 and 3," Graduate Texts in Mathematics, Vol. 147. Springer-Verlag, Berlin-etc., 1977.
- [97] J. W. Morgan, The work of Grigory Perelman. Notices Amer. Math. Soc, 54 (2007), 393–399.
- [98] J. R. Munkres, "Topology (Second Ed.)." Prentice-Hall, Upper Saddle River, NJ, 2000.
- [99] J. A. Neisendorfer, Primary homotopy theory. Mem. Amer. Math. Soc. 25 (1980), No. 232.
- [100] J. A. Neisendorfer, 3-primary exponents. Math. Proc. Cambridge Philos. Soc. 90 (1981), 63–83.
- [101] W. D. Neumann, and D. B. Zagier, A note on an invariant of Fintushel and Stern. "Geometry and topology (College Park, Md., 1983/84)," pp. 241–244, Lecture Notes in Math., 1167. Springer-Verlag, Berlin-etc., 1985.
- [102] S. P. Novikov, Classical and modern topology. Topological phenomena in real world physics. "GAFA 2000 (Tel Aviv, 1999)," Geom. Funct. Anal. 2000, Special Volume, Part I, pp. 406-424. Birkhäuser Verlag, Basel, 2000.
- [103] P. Orlik, "Seifert manifolds." Lecture Notes in Mathematics, 291. Springer-Verlag, Berlin-etc., 1972.
- [104] C. D. Papakyriakopoulos, On Dehn's lemma and the asphericity of knots. Ann. of Math. 66 (1957), 1–26.
- [105] F. Quinn, Open book decompositions, and the bordism of automorphisms. Topology 18 (1979), 55–73.
- [106] D. Repovš, The recognition problem for topological manifolds: a survey. Kodai Math. J. 17 (1994), 538–548.
- [107] J. J. Rotman, "Introduction to the Theory of Groups (Fourth Ed.)." Graduate Texts in Mathematics No. 148. Springer-Verlag, Berlin-etc., 1995.
- [108] D. C. Ravenel, "Complex Cobordism and Stable Homotopy Groups of Spheres (Second Edition)." AMS Chelsea Publishing, Providence, RI, 2004.
- [109] C. P. Rourke, The Hauptvermutung according to Casson and Sullivan. "The Hauptvermutung book," K-Monogr. Math., 1, Kluwer Acad. Publ., Dordrecht, NL, 1996, pp. 129–164.
- [110] C. P. Rourke and B. J. Sanderson, "Introduction to piecewise-linear topology," Springer Study Edition, Ergebnisse Math. (2), Band 69, Springer-Verlag, Berlin-etc., 1982.
- [111] T. Rowland, "Free Action." MathWorld–A Wolfram Web Resource. Online, available at http://mathworld.wolfram.com/FreeAction.html .
- [112] T. Rowland, "Group Action." MathWorld-A Wolfram Web Resource. Online, available at http://mathworld.wolfram.com/GroupAction.html.
- [113] T. Rowland and E. W. Weisstein, "Connected Sum." MathWorld–A Wolfram Web Resource. Online, available at http://mathworld.wolfram.com/ConnectedSum.html .
- [114] T. B. Rushing, "Topological embeddings." Pure and Applied Mathematics, Vol. 52. Academic Press, New York and London, 1973.
- [115] L. Schläfli, Theorie der vielfachen Kontinuität. "Denkschriften der Schweizerischen naturforschenden Gesellschaft," Vol. 38, pp. 1-237, Zürcher und Furrer, Zürich, 1901.

[Originally Written in 1850-1852.] — Reprinted in: Ludwig Schläfli, 1814-1895, Gesammelte Mathematische Abhandlungen, Bd. I, Birkhäuser, Basel, 1950, pp. 167-387. Online: http://historical.library.cornell.edu/cgi-bin/cul.math/docviewer?did=Schl022&seq=7.

- [116] R. Schultz, Some recent results on topological manifolds. Amer. Math. Monthly 78 (1971), 941–952.
- [117] R. Schultz, Smoothable submanifolds of smooth manifolds. Department of Mathematics, University of California, Riverside, 2007. Online, available at http://math.ucr.edu/~res/miscpapers/smoothablesubmflds.pdf.
- [118] A. Scorpan, "The Wild World of 4-Manifolds." American Mathematical Society, Providence, RI, 2005.
- [119] Selick, Paul, Odd primary torsion in $\pi_k(S^3)$. Topology 17 (1978), 407–412.
- [120] L. C. Siebenmann, On detecting open collars. Trans. Amer. Math. Soc. 142 (1969), 201–227.
- [121] L. C. Siebenmann, Disruption of low-dimensional handlebody theory by Rochlin's theorem. Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969), pp. 57–76. Markham, Chicago, 1970.
- [122] L. C. Siebenmann, Infinite simple homotopy types. Nederl. Akad. Wetensch. Proc. (A)
 73 = Indag. Math. 32 (1970), 479–495.
- [123] L. C. Siebenmann, Topological manifolds. "Actes du Congrès International des Mathématiciens (Nice, 1970)," Tome 2, pp. 133–163. Gauthier-Villars, Paris, 1971.
- [124] D. S. Silver, Finding stable boundaries for open five-dimensional manifolds. Amer. J. Math. 105 (1983), 1309–1324.
- [125] S. Smale, On the structure of manifolds. Amer. J. Math. 84 (1962), 387–399.
- [126] R. Solomon, On the finite simple groups and their classification. Notices Amer. Math. Soc. 42 (1995), 231–239.
- [127] A. I. Suciu, Immersed spheres in \mathbb{CP}^2 and $S^2 \times S^2$. Math. Zeitschrift **196** (1987), 51–57.
- [128] I. Tamura, Spinnable structures on differentiable manifolds. Proc. Japan Acad. 48 (1972), 293–296.
- [129] I. Tamura, Foliations and spinnable structures on manifolds. "Colloque International sur l'Analyse et la Topologie Différentielle (Colloques Internationaux du Centre National de la Recherche Scientifique, Strasbourg, 1972)." Ann. Inst. Fourier (Grenoble) 23 (1973), 197–214.
- Ph.D. [130] L. R. Taylor, "Surgery on Paracompact Manifolds." The-University California, Berkeley, 1972. Available online sis, of at http://www.maths.ed.ac.uk/~aar/surgery/taylor.pdf.
- [131] H. Toda, "Composition Methods in Homotopy Groups of Spheres." Ann. of Math. Studies, No. 49. Princeton University Press, Princeton, NJ, 1962.
- [132] F. Waldhausen, On irreducible 3-manifolds which are sufficiently large. Ann. of Math. 87 (1968), 56–88.
- [133] C. T. C. Wall, On simply-connected 4-manifolds. J. London Math. Soc. 39 (1964), 141– 149.
- [134] C. T. C. Wall, "Surgery on compact manifolds (Second edition; edited and with a foreword by A. A. Ranicki)." Mathematical Surveys and Monographs, 69. American Mathematical Society, Providence, RI, 1999.
- [135] E. W. Weisstein, "Classification Theorem of Finite Groups." MathWorld-A Wolfram Web Resource. Online, available at http://mathworld.wolfram.com/ ClassificationTheoremofFiniteGroups.html.
- [136] E. W. Weisstein, "Klein Bottle." MathWorld–A Wolfram Web Resource. Online, available at http://mathworld.wolfram.com/KleinBottle.html .

- [137] H. Whitney, The self-intersections of a smooth n-manifold in 2n-space. Ann. of Math. 45 (1944), 220–246.
- [138] R. L. Wilder, "Topology of Manifolds," Amer. Math. Soc. Colloquium Publications Vol. 32. American Mathematical Society, New York, NY, 1949.
- [139] H. E. Winkelnkemper, Manifolds as open books. Bull. Amer. Math. Soc. 79 (1973), 45–51.
- [140] H. E. Winkelnkemper. The History and Applications of Open Books. Appendix to: A. A. Ranicki, "High-dimensional Knot Theory. Algebraic surgery in codimension 2." Springer Monographs in Mathematics. Springer-Verlag, Berlin-etc., 1998.
- [141] H. E. Winkelnkemper. Artin Presentations, I: Gauge Theory, 3 + 1 TQFT's and the Braid Groups. J. Knot Theory and Ramifications 11 (2002), 223–275.