

## §B. Rational decomposition of nilpotent transformations

### Review of §A.

Let  $T: V \rightarrow V$  be a linear transformation, where  $V$  is a finite dimensional complex vector space. Then  $V$  has a direct sum decomposition  $\bigoplus_{\lambda} V_{\lambda}$  (where  $\lambda$  runs through the eigenvalues)

such that  $T$  maps each  $V_{\lambda}$  into itself and on  $V_{\lambda}$  we have  $(T - \lambda I)^{n(\lambda)} = 0$  for some positive integer  $n(\lambda)$ .

The preceding allows us to find a "block diagonal" matrix representing  $T$ .

Let  $B_{\lambda}$  be an ordered basis for  $V_{\lambda}$  and order the eigenvalues as  $\lambda_1, \dots, \lambda_k$ .

Then on each  $V_{\lambda_j}$  we have  $T = \lambda_j I + N_j$  where  $N_j^{m(\lambda_j)} = 0$ . We can then assemble these into the following matrix for  $T$ :

$$\begin{array}{c}
 B_{\lambda_1} \\
 B_{\lambda_2} \\
 \vdots \\
 B_{\lambda_k}
 \end{array}
 \begin{pmatrix}
 \overset{B_{\lambda_1}}{\lambda_1 I + B_1} & \overset{B_{\lambda_2}}{0} & \dots & \overset{B_{\lambda_k}}{0} \\
 0 & \lambda_2 I + B_2 & & 0 \\
 \vdots & & \ddots & \vdots \\
 0 & \dots & & \lambda_k I + B_k
 \end{pmatrix}$$

Each of the matrices  $B_j$  is nilpotent.\*

To complete the picture, we want matrices of the  $B_j$  for the subspaces  $\mathbb{B}V_{\lambda_j}$  ~~which~~ <sup>which</sup> have as many zero entries as possible. We can do this individually for each  $j$ .

Historical note The term "nilpotent" (accent on the second syllable), like much of linear algebra, is due to J. J. Sylvester (1814-1897). Biographical information about Sylvester is available online, either through Wikipedia or the University of St. Andrews history of mathematics site:

\*  $B$  is nilpotent if  $B^q = 0$  for some  $q$ . The least  $q$  is called the index of nilpotency.

[www-groups.dcs.st-and.ac.uk/history/biography/](http://www-groups.dcs.st-and.ac.uk/history/biography/)

~~James Joseph Sylvester~~  
Sylvester.html

### Iterations of linear transformations

Since  $T^m v = 0 \Rightarrow T^{m+1} v = 0$

( $T: V \rightarrow V$  a linear transformation)

we have

$$\text{Kernel } T \subseteq \text{Kernel } T^2 \subseteq \dots \subseteq V.$$

Suppose that  $\dim V = n$ , and let  $d_j = \dim$

( $\text{Ker } T^j$ ). Then we have

$$d_1 \leq d_2 \leq \dots \leq n = \dim V.$$
 Since the  $d_j$ 's

are integers, this sequence is eventually constant.

In fact, if two consecutive terms are equal, then all ~~remain~~ further terms take the same value.

Lemma 1 If  $\text{Ker } T^m = \text{Ker } T^{m+1}$ ,  
then  $\text{Ker } T^{m+j} = \text{Ker } T^m$  for all  $j \geq 0$ .

PROOF: Will show the conclusion by

induction on  $j$ . The result is known if  
 $j = 0$  or  $1$ . If it is true for  $j = p$ , then

we need to show  $\text{Ker } T^{p+m} = \text{Ker } T^{p+m+1}$ .

We know  $(\subseteq)$  is true. To check  $(\supseteq)$ ,

Suppose  $T^{p+m+1}x = 0$ . Then

$T^p x \in \text{Ker } T^{m+1}$ . But the latter is

$\text{Ker } T^m$ , and hence  $T^m T^p x = 0$ ,

so that  $\text{Ker } T^{p+m+1} = \text{Ker } T^{p+m}$ , and

we know the latter is just  $\text{Ker } T^m$ .  $\square$

Specialize to the nilpotent case.

Lemma 2 (i) If  $T$  is nilpotent, then

$$T^n = 0.$$

(ii) If  $T$  is nilpotent and  $k > 0$  is minimal so that  
 $T^k = 0$ , then  $0 < d_1 < \dots < d_k = n$ .

PROOF. The second conclusion follows because of Lemma 1. To see the second note that if  $0 \leq d_1 < \dots < d_h = n$ , then

$h \leq n$ , so it suffices to prove that

$d_1 > 0$ ; i.e.,  $\text{Ker } T \neq \{0\}$ . However,

if  $\text{Ker } T \neq \{0\}$  then  $T$  is invertible and

each power of  $T$  has  $\text{Ker } T^k = \{0\}$ .

This is impossible if  $T$  is nilpotent, for then  $\text{Ker } T^s = V$  for some large  $s$ .  $\blacksquare$

Corollary 3  $T$  nilpotent  $\Rightarrow T$  is not invertible.  $\blacksquare$

Let's analyze the special case of a nilpotent  $N: V \rightarrow V$  such that  $N^2 = 0$ .

Note first that  $\text{Image } N \subseteq \text{Kernel } N$

in such cases. Let  $x_1, \dots, x_p$  be a basis for  $\text{Im}(N)$  and expand it to a basis for  $\text{Ker}(N)$  by

adjoining  $y_1, \dots, y_q$ . Finally let  $z_1, \dots, z_p$  be such that  $Tz_i = x_i$ . Then we can represent  $N$  by the following matrix:

	<del><math>x_1</math></del>	<del><math>z_1</math></del>	<del><math>x_2</math></del>	<del><math>z_2</math></del>	<del><math>\dots</math></del>	<del><math>x_p</math></del>	<del><math>z_p</math></del>	$y_1$	$\dots$	$y_q$
<del><math>x_1</math></del>	0	1								
<del><math>z_1</math></del>	0	0								
$x_2$			0	1						
$z_2$			0	0						
$\dots$					$\dots$					
$x_p$						0	1			
$z_p$						0	0			
$y_1$										
$\dots$										
$y_q$										

This is a minimally sparse matrix for a linear transformation with rank  $p$ . (At least  $p$  columns must have ~~non~~ zero entries!)

Strategy for the case where  $N^3 = 0$ . ( $N^2 \neq 0$ ).

Let  $V_0 = \text{Image } N$ . Then  $N$  sends  $V_0$  to itself. If  $N_0: V_0 \rightarrow V_0$  is given by the same formula as  $N$ , then  $N_0^2 = 0$

$$(N^2(Ny) = N^3y = 0y = 0).$$

Construct a basis for  $V_0$  with respect to  $N_0$ :

$$\begin{aligned} z_1, \dots, z_p, y_1, \dots, y_q \\ Nz_1, \dots, Nz_p \end{aligned}$$

Now choose  $z'_i, y'_j \in V$  such that

$$Nz'_i = z_i, \quad Ny'_j = y_j$$

obtaining a new (linearly independent) set in  $V$ :

$$X = \left\{ \begin{array}{ll} z'_1, \dots, z'_p & y'_1, \dots, y'_q \\ Nz'_1, \dots, Nz'_p & Ny'_1, \dots, Ny'_q \\ N^2z'_1, \dots, N^2z'_p & \end{array} \right\}$$

We need to verify this set is linearly independent.

Suppose that

$$\begin{aligned} \sum a_j z_j' + \sum b_j N z_j' + \sum c_j N^2 z_j' \\ + \sum d_j y_j' + \sum e_j N y_j' = 0 \end{aligned}$$

If we apply  $N$ , we get

$$\sum a_j z_j + \sum b_j N z_j + \sum d_j y_j = 0$$

and by construction the  $a_j$ 's,  $b_j$ 's and  $d_j$ 's are all 0. Substituting this into the top equation, we get

$$\sum c_j N z_j + \sum e_j y_j = 0$$

and again by construction this yields  $c_j = 0, e_j = 0$  for all  $j$ .

NOW expand  $X$  to a basis for  $V$  by adjoining  $\{w_1, \dots, w_r\}$ . The final step is to modify these slightly, obtaining  $\{w_1, \dots, w_r\}$  with  $N w_j = 0$  for all  $j$ .



Specifically, we know

$$Nw_j = Nv_j \text{ for some } v_j \in \text{Span}(X)$$

If  $w_j = w_j - v_j$  then  $X \cup \{w_1, \dots, w_n\}$  is also a basis for  $V$ , but now  $Nw_j = 0$  all  $j$ .

Thus there is a basis for  $V$  of the form.

$$\left\{ \begin{array}{ccc} N^i z_j' & N^i y_j' & w_j \\ i=0, 1, 2 & i=0, 1 & \dots \end{array} \right\} \cdot \blacksquare$$

$$\left\{ \begin{array}{ccc} N^3 z_j' = 0 & N^2 y_j' = 0 & Nw_j = 0 \end{array} \right\}$$

NOTE. Either the second or third column (or both) might be empty!

### RECURSIVE REFORMULATION

Induction on the index of nilpotency (= least  $p > 0$  such that  $N^p = 0$ ).

Examine the proof when  $p = 3$ , then generalize the conclusion.

We have a basis  $N^i y_{k,j}$  for  $V$

each piece of which has the form

$$y_{k,j} = N^0 y_{k,j}, N^1 y_{k,j}, \dots, N^{m(k)-1} y_{k,j}$$

such that  $N^{m(k)} y_{k,j} = 0$ , where  $1 \leq j \leq q(k)$ ,

and  $p = m(1) \geq m(2) \geq \dots (\geq 1)$ .

Assume this is known if  $N^p = 0$ , and [ $p = \text{index of nil } p$ ]  
 now suppose we have  $N$  with  $N^{p+1} = 0$  but  $N^p \neq 0$ .

Let  $V_0 = \text{Image } N$ , so that  $N[V_0] \subseteq V_0$ ,  
 and let  $N_0: V_0 \rightarrow V_0$  be defined by  $N_0(x) = Nx$ .  
 Then  $y \in V_0 \Rightarrow y = Nx \Rightarrow N^p y = N^{p+1} x = 0$ ,  
 so that  $N_0^p = 0$ . Also, if  $z \in V$  is such  
 that  $N^p z \neq 0$ , then  $N_0^{p-1}(Nz) \neq 0$ , so the  
 index of nilpotency for  $N_0$  is equal to  $p$ .

Now apply the induction hypothesis

Let  $\{N^i y_{k,j} \mid i, j, k\}$  be the basis for  $V_0$  as described on the preceding page, and write  $y_{k,j} = N x_{k,j}$  (can do since  $V_0 = N[V]$ ). Consider the set with

$$x_{k,j} = N^0 x_{k,j}, \quad N^1 x_{k,j}, \quad \dots, \quad N^{m(k)} x_{k,j}$$

$\parallel$   
 $y_{k,j}$ 
 $\parallel$   
 $N^{m(k)-1} y_{k,j}$

As in the proof when  $p+1=3$ , this set is linearly independent. Now expand it to a basis for  $V$  by adjoining  $z'_1, \dots, z'_r$ . Then each  $z'_\ell$  has the property  $Nz'_\ell = w'_\ell$  where the latter is a linear combination of the given basis for  $V_0$ , and  $w'_\ell = Nw_\ell$  where  $w_\ell$  is a linear comb. of the  $N^i x_{k,j}$ 's. The new set, still spans  $V$  (hence is also a basis),

$z_\ell = z'_\ell - w_\ell$  but now  $N(z_\ell) = 0$ , all  $\ell$ .

sub-script  
"ell"

This completes the proof that there is a basis for  $V$  of the form appearing on p. 10, and hence completes the proof of the inductive step. ■

### Conclusion, stated in terms of matrices

If  $N$  is nilpotent, then  $N$  can be represented by a block sum of matrices which are zero off the diagonal and have diagonal pieces like the following:

$$\begin{pmatrix} 0 & 1 & 0 & & \\ & 0 & 1 & 0 & \\ & & 0 & 0 & 1 \\ & & & \ddots & \\ & & & & 0 & \ddots \\ & & & & & & 0 \end{pmatrix}$$

Formally,  $a_{ij} = 1$  if  $j = i+1$  and 0 otherwise. ■