

**SOLUTIONS TO SELECTED  
ADDITIONAL EXERCISES FOR  
MATHEMATICS 132 FOR UNITS IV AND V FROM  
linalgexercises2.pdf**

Winter 2004

**IV. Complex linear algebra**

**IV.1: Complex numbers**

*Additional exercises*

1. (a)  $20 + 10i$     (b)  $8$     (c)  $(a^2 - b^2) + 2abi$     (d)  $-2 + 2i$     (e)  $(a^3 - 3ab^2) + (3a^2b - b^3)i$

2.  $65$

3.  $\frac{13}{10} + \frac{9}{10}i$

4. (a)  $\frac{1}{2} \cdot (\cos(4\pi/9) + i \sin(4\pi/9))$

5. (a)  $-4$     (b)  $-32i$     (c)  $-8$

6. The cube roots of 8 are 2 and

$$2(\cos(2\pi/3) \pm i \sin(2\pi/3))$$

and the cube roots of 27 are 3 and

$$3(\cos(2\pi/3) \pm i \sin(2\pi/3)).$$

Note that

$$\cos(2\pi/3) - i \sin(2\pi/3) = \cos(4\pi/3) + i \sin(4\pi/3).$$

**IV.2: Complex matrices**

1.

(a)

$$\begin{pmatrix} 2 & 1+3i \\ -1-2i & -4i \end{pmatrix}$$

(b)

$$\begin{pmatrix} 2+2i & 2 \\ 4-4i & -6i \end{pmatrix}$$

(c)

$$\begin{pmatrix} -2 + 2i & 2i \\ 4 + 4i & 6 \end{pmatrix}$$

(d)

$$-5 - 3i$$

(e)

$$\begin{pmatrix} -5 & -15 + 10i \\ 25i & 15 + 30i \end{pmatrix}$$

2. (a) Not invertible. (b) The inverse matrix is equal to

$$-\frac{1}{3} \cdot \begin{pmatrix} i - 3i & \\ -2 + i & 6 \end{pmatrix}.$$

3. The matrix is not invertible if and only if  $z = -\frac{5}{3} - \frac{10}{3}i$ .

4. (a) Not a basis. (b) Is a basis.

5. (a)  $\sqrt{7}$  (b)  $\sqrt{17}$

6.  $\sqrt{15}$

7. (a) Linearly dependent. (b) Linearly independent.

8.  $\mathbf{x}$  is equal to the transpose of  $(2 \ 1)$ .

9.

(a)

$$\begin{pmatrix} 1 & 2 \\ i & -3i \end{pmatrix}$$

(b)

$$\begin{pmatrix} 0 & 5 + i & \sqrt{2}i \\ 5 - i & 6 & 4 \\ -\sqrt{2}i & 4 & 3 \end{pmatrix}$$

10. (a) Hermitian. (b) Not Hermitian; one of its diagonal entries is imaginary, and the diagonal entries of a Hermitian matrix are real.

### IV.3: Complex eigenvalues and eigenvectors

1. (a) The eigenvalues are  $\pm 1$  and associated eigenvectors are given by  $\mathbf{v}_1 = (1, -i)$  and  $\mathbf{v}_{-1} = (1, i)$ .

(b) The eigenvalues are 1, 2 and 3, and associated eigenvectors are given by  $\mathbf{v}_1 = (\sqrt{2}, -i, i)$ ,  $\mathbf{v}_2 = (0, 1, 1)$  and  $\mathbf{v}_3 = (\sqrt{2}, i, -i)$ .

## V. Quadratic forms

### V.3 : Classification of critical points

1. (a)  $(-3, 2)$  is a relative minimum. (b)  $(-18/5, -11/15)$  is a relative minimum. (c)  $(0, 0)$  is a saddle point. (d)  $(0, 0)$  is a saddle point and  $(1, 1)$  is a relative minimum. (e)  $(0, 0)$  is a relative minimum and  $(\pm\sqrt{2}, 1)$  are saddle points. (f) No critical points. (g)  $(0, n\pi)$  is a saddle point, where  $n$  is an integer.