

### SUPPLEMENTARY EXERCISES ON AFFINE EQUIVALENCE

**T1.** Let  $K$  be a nonempty convex subset of  $\mathbb{R}^2$ , and let  $L$  be a line in  $\mathbb{R}^2$  such that  $K \cap L$  is empty. Prove that all the points of  $K$  lie on the same side of  $L$ . [*Hint:* Assume that  $K$  contains points on both sides of  $L$  and derive a contradiction.]

**T2.** Let  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$  be noncollinear points in  $\mathbb{R}^2$ , and let  $F$  be an affine transformation of  $\mathbb{R}^2$ . Prove that  $F$  sends  $\angle \mathbf{abc}$  to  $\angle F(\mathbf{a})F(\mathbf{b})F(\mathbf{c})$ . [*Hint:* By Corollary II.4.8  $F$  maps rays to rays, and we also know that if  $X$  and  $Y$  are subsets of  $\mathbb{R}^2$  then  $F$  maps the union  $X \cup Y$  to  $F[X] \cup F[Y]$ .]

**T3.** Assume the setting of the preceding exercise, and prove that  $F$  maps the exterior of  $\angle \mathbf{abc}$  to the exterior of  $\angle F(\mathbf{a})F(\mathbf{b})F(\mathbf{c})$ . [*Hint:* By Theorem 12 in `affine-convex.pdf` and a previous exercise we know that  $F$  maps the angle and its interior to themselves.]

**T4.** If  $L$  and  $M$  are parallel lines in  $\mathbb{R}^2$ , then by Exercise **T1** we know that all points of  $L$  lie on the same side of  $M$  and all points of  $M$  lie on the same side of  $L$ . Define the *strip between  $L$  and  $M$*  to be the set of points  $\mathbf{x}$  such that  $\mathbf{x}$  and  $L$  are the same side of  $M$  and  $\mathbf{x}$  and  $M$  are the same side of  $L$ .

(i) Prove that the strip between  $L$  and  $M$  is convex and nonempty. Specifically, prove that if  $A \in L$  and  $B \in M$ , then the midpoint  $C$  of  $(AB)$  lies in this set.

(ii) Prove that if  $F$  is an affine transformation of  $\mathbb{R}^2$  and  $L$  and  $M$  are parallel lines in  $\mathbb{R}^2$ , then  $F$  maps the strip between  $L$  and  $M$  to the strip between  $F[L]$  and  $F[M]$ . [*Hint:* Apply Theorem 11 in `affine-convex.pdf`.]