

Before Gauss, some mathematicians (for example, Klügel) had speculated that a proof of the Fifth Postulate might be out of reach. However, Gauss (and to a lesser extent a contemporaries like Schweikart and Taurinus) took things an important step further, concluding that the negation of the Fifth Postulate yields a geometrical system which is very different from Euclidean geometry in some respects but has exactly the same degree of logical validity (compare also the passage from the letter to Olbers at the beginning of this unit). Working independently of Gauss, J. Bolyai (1802 – 1860) and N. I. Lobachevsky (1792 – 1856) reached the same conclusions as Gauss (each one independently of the other), which Bolyai summarized in a frequently repeated quotation:

Out of nothing I have created a strange new universe.


Both Bolyai and Lobachevsky took everything one important step further than Gauss by publishing their conclusions, and for this reason they share credit for the first published recognition of hyperbolic geometry as a mathematically legitimate subject.

Details for a result from Lecture 15 (note fixed typos)

Appendix to Section 3: Proof of Theorem 8

The major steps in the argument will be presented as lemmas.

Lemma 8A. (Splicing Property). *Suppose that $\square ABCD$ is a rectangle, and let $C_1 \in (DC$ be a point such that $|DC_1| = 2|DC|$. Let B_1 be the foot of the perpendicular from C_1 to AB . Then $|\angle DC_1B| = 90^\circ$ and the points A, B_1, C_1 and D (in that order) are the vertices of a rectangle.*



Proof. First of all, the lines AD , BC , and B_1C_1 , are all parallel to each other because every two of them have a common perpendicular (namely, AB). Therefore AD and B_1C_1 are contained in the D – and C_1 – sides of BC respectively. But $|DC_1| = 2|DC|$ and $C_1 \in (DC$ imply $D * C * C_1$ is true. This in turn implies that D and C_1 are on opposite sides of BC . Since B is the common point of the lines AB_1 and BC , it follows that $A * B * B_1$ is true.

Since AD and B_1C_1 are parallel (they have a common perpendicular), the points B_1 and C_1 lie on the same side of AD . Hence A, B_1, C_1 , and D (in that order) form the vertices of a convex quadrilateral. Likewise B, B_1, C_1 , and C form the vertices of a convex quadrilateral. By construction, S.A.S. applies to show $\triangle ADC \cong \triangle BCC_1$. It follows

that $|\mathbf{AC}| = |\mathbf{BC}_1|$, $\gamma = |\angle \mathbf{CBC}_1| = |\angle \mathbf{DAC}| = \alpha$, and $\eta = |\angle \mathbf{BC}_1\mathbf{C}| = |\angle \mathbf{ACD}|$.

On the other hand, if $\xi = |\angle \mathbf{C}_1\mathbf{C}\mathbf{B}_1|$ then

$$\alpha + \beta = 90^\circ = \gamma + \xi.$$

Therefore $\alpha = \gamma$ implies $\beta = \xi$.

By A.A.S. it follows that $\triangle \mathbf{ABC} \cong \triangle \mathbf{BB}_1\mathbf{C}_1$, and hence $\alpha = \delta = |\angle \mathbf{BC}_1\mathbf{B}_1|$.

This implies that $\eta + \delta = 90^\circ$. But then it follows that $|\angle \mathbf{DC}_1\mathbf{B}_1| = \eta + \delta = 90^\circ$. ■

Lemma 8B. *If there is a rectangle $\square \mathbf{ABCD}$ in the neutral plane under consideration, then for every positive integer n there is a rectangle $\square \mathbf{A'B'C'D'}$ with $|\mathbf{A'B'}| = |\mathbf{C'D'}| = n|\mathbf{AB}| = n|\mathbf{CD}|$ and $|\mathbf{A'D'}| = |\mathbf{B'C'}| = |\mathbf{AD}| = |\mathbf{BC}|$.*


Proof. The case $n = 2$ was done in the preceding lemma. Assume by induction that we have $\mathbf{B} = \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_{n-1}$ and $\mathbf{C} = \mathbf{C}_1, \mathbf{C}_2, \dots, \mathbf{C}_{n-1}$ such that

$$\mathbf{A} = \mathbf{A}_0 * \mathbf{A}_1 * \mathbf{A}_2 * \dots * \mathbf{A}_{n-1} \quad \text{and} \quad \mathbf{D} = \mathbf{C}_0 * \mathbf{C}_1 * \mathbf{C}_2 * \dots * \mathbf{C}_{n-1}$$

(the notation means that $\mathbf{X}_p * \mathbf{X}_q * \mathbf{X}_r$ if $p < q < r$) and the following additional conditions:

For $k = 1, \dots, n-1$ we have $|\mathbf{AB}| = |\mathbf{CD}| = |\mathbf{A}_{k-1}\mathbf{A}_k| = |\mathbf{C}_{k-1}\mathbf{C}_k|$.

For $k = 1, \dots, n-1$ the line $\mathbf{C}_k\mathbf{A}_k$ is perpendicular to both \mathbf{AB} and \mathbf{CD} .



Now apply Lemma 8A to the rectangle $\mathbf{A}_{n-2}\mathbf{A}_{n-1}\mathbf{C}_{n-1}\mathbf{C}_{n-2}$ to obtain \mathbf{A}_n and \mathbf{C}_n such that $\mathbf{A}_{n-2} * \mathbf{A}_{n-1} * \mathbf{A}_n$ and $\mathbf{C}_{n-2} * \mathbf{C}_{n-1} * \mathbf{C}_n$, $|\mathbf{AB}| = |\mathbf{CD}| = |\mathbf{A}_{n-1}\mathbf{A}_n| = |\mathbf{C}_{n-1}\mathbf{C}_n|$, and $\mathbf{C}_n\mathbf{A}_n$ is perpendicular to both \mathbf{AB} and \mathbf{CD} . ■


Corollary 8C. *If there is a rectangle $\square \mathbf{ABCD}$ in the neutral plane under consideration, then for each pair of integers $n, m > 0$ there is a rectangle $\square \mathbf{A'B'C'D'}$ with $|\mathbf{A'B'}| = |\mathbf{C'D'}| = n|\mathbf{AB}| = n|\mathbf{CD}|$ and $|\mathbf{A'D'}| = |\mathbf{B'C'}| = m|\mathbf{AD}| = m|\mathbf{BC}|$.*

Proof. First apply Lemma B to get a rectangle $\square \mathbf{A^*B^*C^*D^*}$ with $|\mathbf{A^*B^*}| = n|\mathbf{AB}|$ and $|\mathbf{B^*C^*}| = |\mathbf{BC}|$. Now apply Lemma B again to get a new rectangle $\square \mathbf{A'B'C'D'}$ with

$|A'B'| = |A^*B^*|$ and $|B'C'| = m |B^*C^*|$. It follows that $|A'B'| = |C'D'| = n |AB| = n |CD|$ and $|A'D'| = |B'C'| = m |AD| = m |BC|$. ■

The next lemma allows us to take a large rectangle and trim it to another of smaller size.

Lemma 8D. Let $\square ABCD$ be a rectangle, let $X \in (CD)$, and let Y be the foot of the perpendicular from X to AB . Then $A, Y, X,$ and D (in that order) are the vertices of a rectangle.



Proof. The lines $AD, XY,$ and BC are all parallel because they are all perpendicular to AB . Hence AD is contained in the D – side of XY and BC is contained in the C – side of XY . But C^*X^*D since X lies on (BC) , and therefore C and D lie on opposite sides of XY . Hence AD and BC also lie entirely on opposite sides of XY . Since AB and XY meet at Y , it follows that A^*Y^*B must be true.

Label the angle measures as indicated in the drawing above:

$\alpha = \angle YDX $		$\xi = \angle CYX $
$\beta = \angle DYX $		$\eta = \angle XCY $
$\gamma = \angle ADY $		$\zeta = \angle CYB $
$\delta = \angle AYD $		$\omega = \angle YCB $

Since AD is parallel to XY, CD is parallel to $XY,$ and AB is parallel to $CD,$ it follows that $A, Y, X,$ and D and $Y, B, X,$ and X (in these orders) form the vertices of a convex quadrilateral. Therefore D lies in the interior of $\angle AYX, Y$ lies in the interior of $\angle ADYX, C$ lies in the interior of $\angle XYB,$ and Y lies in the interior of $\angle XCB$. These imply the following four equations:


$\alpha + \gamma = 90^\circ$		$\xi + \zeta = 90^\circ$
$\beta + \delta = 90^\circ$		$\eta + \omega = 90^\circ$

The Saccheri – Legendre Theorem implies the following additional inequalities:

$\xi + \gamma \leq 90^\circ$		$\omega + \zeta \leq 90^\circ$
------------------------------	--	--------------------------------

Taken together, these imply $90^\circ \leq \alpha + \beta$ and $90^\circ \leq \xi + \eta$. Therefore the Saccheri – Legendre Theorem implies that both $|\angle DXY|$ and $|\angle CXY|$ are less than or equal to 90° . On the other hand, $C \ast X \ast D$ implies that $|\angle DXY| + |\angle CXY| = 180^\circ$. This can happen only if both $|\angle DXY|$ and $|\angle CXY|$ are equal to 90° . But this now implies XY is perpendicular to CD , so that $A, Y, X,$ and D (in that order) are the vertices of a rectangle. ■

Proof of Theorem 8. Given rectangle $\square ABCD$ and real numbers $p, q > 0$, find positive integers n and m so that $p < n|AB|$ and $q < m|AD|$. Form a rectangle $\square A'B'C'D'$ with $|A'B'| = n|AB|$ and $|A'D'| = q|AD|$.



Let $X \in (A'B')$ satisfy $|A'X| = p < |A'B'|$, and let Y be the foot of the perpendicular from X to $C'D'$. Then by Lemma 8D one obtains a rectangle $\square A'YXD'$ with $|A'X| = p$ and $|A'D'| = q|AD|$.

should be
XY

Now let $W \in (A'D')$ satisfy $|A'W| = q < |A'D'|$, and let V be the foot of the perpendicular from Z to XY . Then $A', Y, V,$ and W (in that order) are the vertices of a rectangle with $|A'X| = p$ and $|A'W| = q$. ■

should be
X