

## Vectors and Parallel Lines


Theorem Let  $AB$  and  $CD$  be lines in  $\mathbb{R}^2$ . Then the following are equivalent:

- (i) Either  $AB \parallel CD$  or  $AB = CD$
- (ii)  $B-A$  and  $D-C$  are nonzero multiples of each other.

This approach relies heavily on Moise, Chapter 17. Given a point  $X$ , write  $X = (x_1, x_2)$ .

Derivation. There are two cases: Lines vertical and nonvertical. We need

Claim: A vertical and a nonvertical line meet at one point.



Verification Let  $x = a$  be the vertical line  $y = mx + b$  the nonvertical one. Then the only common pt. is  $(a, ma + b)$ .

VERTICAL CASE (i)  $\Rightarrow$  (ii) Lines are vertical  
 $x = a$  &  $x = b$ . So we have

$A = (a, y_1)$ ,  $B = (a, y_2)$ ,  $C = (b, y_3)$ ,  $D = (c, y_4)$   
 where  $y_1 \neq y_2$  &  $y_3 \neq y_4$ . Both  $B-A$   
 and  $D-C$  are nonzero multiples of  
 $(0, 1)$ .

(ii)  $\Rightarrow$  (i) Two vertical lines are  
 always equal or parallel.  $\square$

NON VERTICAL CASE. Say the lines

have equations  $y = mx + p$   
 $y = nx + q$

By Moire they are parallel or equal  $\Leftrightarrow$

$m = n$ . Also two points on such a line  
 are equal  $\Leftrightarrow$  first coords equal because there  
 is a formula for finding the second coord.

from the first,

(i)  $\Rightarrow$  (ii) Given  $m = n$ . Then

$$A = (a_1, ma_1 + p)$$

$$B = (b_1, mb_1 + p)$$

$$C = (c_1, mc_1 + q)$$

$$D = (d_1, md_1 + q)$$

So

$$B-A = (b_1 - a_1, m(b_1 - a_1))$$

$$D-C = (d_1 - c_1, m(d_1 - c_1))$$

Now  $b_1 \neq a_1$  since  $B \neq A$  and  $d_1 \neq c_1$  since  $D \neq A$ . Therefore

$$D - C = \frac{d_1 - c_1}{b_1 - a_1} (B - A) \quad \text{and}$$

$$B - A = \frac{b_1 - a_1}{d_1 - c_1} (D - C)$$

*> relies heavily on fact that fraction denominators are nonzero!*

(ii)  $\Rightarrow$  (i) Say  $D - C = k(B - A)$

where  $k \neq 0$ . Then

$$d_1 - c_1 = k(b_1 - a_1)$$

$$k(b_2 - a_2) = d_2 - c_2 = m(d_1 - c_1) \quad \text{But also}$$


$$b_2 - a_2 = m(b_1 - a_1), \text{ so}$$

$$km(b_1 - a_1) = k(b_2 - a_2) = m(d_1 - c_1) = km(b_1 - a_1).$$

Since  $b_1 - a_1 \neq 0$  and  $k \neq 0$ , we must have  $m = m$ . Hence

$$AB \parallel CD \text{ or } AB = CD. \quad \square$$

## Parallelogram Law for Vector Addition



If  $O, b, c, d$  (in order) are the vertices of a parallelogram, then  $c = b + d$ .

Derivation Show  $Ob \parallel cd$   
 $Od \parallel bc$ .

Assuming  $b + d$  aren't non zero scalar multiples of each other. Then  $cd \neq Ob$  and  $bc \neq Od$ . ( $d \neq Ob$ ,  $b \neq Od$ ).

$$\left. \begin{array}{l} d - c = -b \\ \text{apply thm to} \\ \text{get} \\ cd \parallel Ob \end{array} \right\} \left. \begin{array}{l} b - c = -d \\ \text{apply thm to} \\ \text{get} \\ bc \parallel Od \end{array} \right.$$

Example. The diagonals of  $\square^{D,C}_{A,B}$  bisect each other

Proof. Previous yields  $(C - A) = (D - A) + (B - A)$ . Mid points are

$$AC: \frac{1}{2}(C + A) = \frac{1}{2}(D - A) + \frac{1}{2}(B - A) + \frac{1}{2}(2A) = \frac{1}{2}(D + B)$$

BD:  $\frac{1}{2}(D + B)$ . So we get the same point!