

Exercises for Unit II (Vector algebra and Euclidean geometry)

II.1 : Approaches to Euclidean geometry

Supplementary background readings.

Ryan : pp. 5 – 15

Exercises to work.

1. What is the minimum number of planes containing three concurrent noncoplanar lines in coordinate 3 – space \mathbf{R}^3 ?
2. What is the minimum number of planes in coordinate 3 – space \mathbf{R}^3 containing five points, no four of which are coplanar? [**Hint**: No three of the points are collinear.]
3. Suppose that $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ are four noncoplanar points in \mathbf{R}^3 . Explain why the lines \mathbf{ab} and \mathbf{cd} are disjoint but not coplanar (in other words, they form a pair of **skew lines**).
4. Let \mathbf{L} be a line in coordinate 2 – space \mathbf{R}^2 or 3 – space \mathbf{R}^3 , let \mathbf{x} be a point not on \mathbf{L} , and let $\mathbf{p}_1, \dots, \mathbf{p}_n$ be points on \mathbf{L} . Prove that the lines $\mathbf{x}\mathbf{p}_1, \dots, \mathbf{x}\mathbf{p}_n$ are distinct. Why does this imply that \mathbf{R}^2 and \mathbf{R}^3 contain infinitely many lines?
5. Suppose that $\mathbf{L}_1, \dots, \mathbf{L}_n$ are lines in coordinate 2 – space \mathbf{R}^2 or 3 – space \mathbf{R}^3 . Prove that there is a point \mathbf{q} which does not lie on any of these lines. [**Hint**: Take a line \mathbf{M} which is different from each of $\mathbf{L}_1, \dots, \mathbf{L}_n$; for each j we know that \mathbf{M} and \mathbf{L}_j have at most one point in common, but we also know that \mathbf{M} has infinitely many points.]
6. Suppose that $\mathbf{p}_1, \dots, \mathbf{p}_n$ are points in coordinate 2 – space \mathbf{R}^2 or 3 – space \mathbf{R}^3 . Prove that there is a line \mathbf{L} which does not contain any of these points. [**Hint**: Let \mathbf{x} be a point not equal to any of $\mathbf{p}_1, \dots, \mathbf{p}_n$ and take a line \mathbf{L} through \mathbf{x} which is different from each of $\mathbf{x}\mathbf{p}_1, \dots, \mathbf{x}\mathbf{p}_n$.]
7. Let \mathbf{P} be a plane in coordinate 3 – space \mathbf{R}^3 , let $\mathbf{a}, \mathbf{b}, \mathbf{c}$ be noncollinear points on \mathbf{P} , let \mathbf{z} be a point which is not on \mathbf{P} , and let \mathbf{u} and \mathbf{v} be distinct points on the line \mathbf{cz} . Show that the planes \mathbf{P}, \mathbf{abu} and \mathbf{abv} are distinct. Using this and ideas from Exercise 4, prove that there are infinitely many planes in coordinate 3 – space \mathbf{R}^3 which contain the line \mathbf{ab} .
8. Suppose we have an abstract system $(\mathbf{P}, \mathcal{L})$ consisting of a set \mathbf{P} whose elements we call **points** and a family of proper subsets \mathcal{L} that we shall call **lines** such

that the points and lines satisfy axioms (I – 1) and (I – 2). Assume further that every line L in P contains at least three points. Prove that P contains at least seven points.

II.2 : Synthetic axioms of order and separation

Supplementary background readings.

Ryan : pp. 11 – 15, 19 – 21, 50 – 52, 55 – 62, 68

Exercises to work.

1. Suppose that A, B, C are collinear points in \mathbf{R}^3 whose coordinates are given by $(a_1, a_2, a_3), (b_1, b_2, b_3),$ and (c_1, c_2, c_3) respectively. Prove that $A*B*C$ holds if we have $a_1 < b_1 < c_1, a_2 = b_2 = c_2,$ and $a_3 > b_3 > c_3.$
2. In the coordinate plane $\mathbf{R}^2,$ let $A = (1, 0), B = (0, 1), C = (0, 0)$ and $D = (-2, -1).$ Show that the lines AB and CD intersect in a point X such that $A*X*B$ and $D*C*X$ hold. [**Hint:** Construct X explicitly.]
3. In the coordinate plane $\mathbf{R}^2,$ suppose that A, B, C, D, E are points not on the same line such that $A*B*C$ and $A*D*E$ hold. Prove that the segments (BE) and (CD) have a point in common. [**Hint:** Use barycentric coordinates with respect to the points A, B and $D.$]
4. In the coordinate plane $\mathbf{R}^2,$ let $X = (4, -2)$ and $Y = (6, 8),$ and let L be the line defined by the equation $4y = x + 10.$ Determine whether X and Y lie on the same side of $L.$
5. In the coordinate plane $\mathbf{R}^2,$ let $X = (8, 5)$ and $Y = (-2, 4),$ and let L be the line defined by the equation $y = 3x - 7.$ Determine whether X and Y lie on the same side of $L.$
6. Let L be a line in the plane $P,$ and suppose that M is some other line in P such that L and M have no points in common. Prove that all points of M lie on the same side of L in $P.$
7. State and prove a generalization of the previous result to disjoint planes in $3 -$ space.
8. Suppose that L is a line in the plane of triangle $\triangle ABC.$ Prove that L cannot meet all three of the open sides $(AB), (BC)$ and $(AC).$

9. Let \mathbf{V} be a vector space over the real numbers, and let $\mathbf{S} = \{ \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k \}$ be a subset of \mathbf{V} . A vector \mathbf{x} in \mathbf{V} is said to be a **convex combination** of the vectors in \mathbf{S} if \mathbf{x} is expressible as a linear combination of the form $a_0\mathbf{v}_0 + a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n$ such that $a_0 + a_1 + \dots + a_n = 1$ and $0 \leq a_j \leq 1$ for all j . Let $\mathbf{S} = \{ \mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_k \}$ be a subset of \mathbf{V} , and let $\mathbf{T} = \{ \mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_m \}$ be a set of vectors in \mathbf{V} which are convex combinations of the vectors in \mathbf{S} . Suppose that \mathbf{y} is a vector in \mathbf{V} which is a convex combination of the vectors in \mathbf{T} . Prove that \mathbf{y} is also a convex combination of the vectors in \mathbf{S} .

II.3 : Measurement axioms

Supplementary background readings.

Ryan : pp. 11 – 15, 50 – 52, 58 – 62, 68

Exercises to work.

- Suppose we are given a line containing the two points \mathbf{A} and \mathbf{B} . Then every point \mathbf{X} on the line can be expressed uniquely as a sum $\mathbf{A} + k(\mathbf{B} - \mathbf{A})$ for some real number k . Let $f : \mathbf{L} \rightarrow \mathbf{R}$ be defined by $f(\mathbf{X}) = kd(\mathbf{A}, \mathbf{B})$. Prove that f defines a $1 - 1$ correspondence such that $d(\mathbf{X}, \mathbf{Y}) = |f(\mathbf{X}) - f(\mathbf{Y})|$ for all points \mathbf{X}, \mathbf{Y} on \mathbf{L} . [**Hint:** Recall that $d(\mathbf{X}, \mathbf{Y}) = |\mathbf{X} - \mathbf{Y}|$.]
- Suppose that we are given a line \mathbf{L} and **two** distinct $1 - 1$ correspondences between \mathbf{L} and the real line \mathbf{R} which satisfy the condition in the Ruler Postulate **D - 3**, say $f : \mathbf{L} \rightarrow \mathbf{R}$ and $g : \mathbf{L} \rightarrow \mathbf{R}$. Prove that these functions satisfy a relationship of the form $g(\mathbf{X}) = af(\mathbf{X}) + b$, where a and b are real numbers with $a = \pm 1$. [**Hint:** Look at the function $h = g \circ f^{-1}$, which is a distance – preserving $1 - 1$ correspondence from \mathbf{R} to itself. Show that such a map has the form $h(t) = at + b$, where a and b are as above. To do this, first show that if h is a distance – preserving $1 - 1$ correspondence from \mathbf{R} to itself then so is $k(t) = h(t) - h(0)$, then show that k must be multiplication by ± 1 .]
- In the coordinate plane, determine whether the point $\mathbf{X} = (9, 4)$ lies in the interior of $\angle\mathbf{ABC}$, where $\mathbf{A} = (7, 10)$, $\mathbf{B} = (2, 1)$ and $\mathbf{C} = (11, 1)$. Also, determine the values of k for which $(17, k)$ lies in the interior of $\angle\mathbf{ABC}$.
- Answer the same questions as in the preceding exercise for $\mathbf{X} = (30, 200)$ and $\mathbf{X} = (75, 135)$.

5. Answer the same questions as in Exercise 3 when $\mathbf{X} = (-5, 8)$, $\mathbf{A} = (-4, 8)$, $\mathbf{B} = (-1, 2)$ and $\mathbf{C} = (-5, -12)$.
6. Suppose we are given $\angle ABC$. Prove that the open segment (AC) is contained in the interior of $\angle ABC$.
7. Suppose we are given $\triangle ABC$ and a point \mathbf{D} in the interior of this triangle, and let \mathbf{E} be any point in the same plane except \mathbf{D} . What general conclusion about the intersection of $\triangle ABC$ and $[DE]$ seems to be true? Illustrate this conjecture with a rough sketch.
8. Suppose that we are given $\triangle ABC$ and a point \mathbf{X} on (BC) . Prove that the open segment (AX) is contained in the interior of $\triangle ABC$.
9. Let $\angle ABC$ be given. Prove that there is a point \mathbf{D} on the same plane such that \mathbf{D} and \mathbf{A} lie on opposite sides of BC , but $d(\mathbf{A}, \mathbf{B}) = d(\mathbf{D}, \mathbf{B})$ and $|\angle ABC| = |\angle DBC|$. [*Hint:* This uses both the Ruler and Protractor Postulates.]
10. (Ryan, Exercise 31, p. 68) Suppose we are given $\triangle ABC$ and a point \mathbf{D} in the interior of this triangle. Prove that \mathbf{D} lies on an open segment (XY) , where \mathbf{X} and \mathbf{Y} lie on the triangle.

II.4 : Congruence, superposition and isometries


Supplementary background readings.

Ryan : pp. 11 – 15, 19 – 21, 49 – 52, 55 – 62, 64 – 66, 68

Exercises to work.

1. Let $\angle ABC$ be given. Prove that there is a unique *angle bisector* ray $[BD]$ such that (BD) is contained in the interior of $\angle ABC$ and $|\angle ABD| = |\angle DBC| = \frac{1}{2} |\angle ABC|$. [*Hints:* Let \mathbf{E} be the unique point on (BA) such that $d(\mathbf{B}, \mathbf{E}) = d(\mathbf{B}, \mathbf{C})$, and let \mathbf{D} be the midpoint of $[CE]$. Recall that there should be proofs for both existence and uniqueness.]
2. Give an example of a triangle $\triangle ABC$ for which the standard formal congruence statement $\triangle ABC \cong \triangle BCA$ is false.
3. Suppose we are given $\triangle ABC$ and $\triangle DEF$ such that $\triangle ABC \cong \triangle DEF$. Explain why we also have $\triangle ACB \cong \triangle DFE$ and $\triangle BCA \cong \triangle EFD$.

4. Suppose $\triangle ABC$ is an isosceles triangle with $d(A, C) = d(B, C)$, and let D and E denote the midpoints of $[AC]$ and $[BC]$ respectively. Prove that $\triangle DAB \cong \triangle EBA$.



5. Suppose we are given $\triangle ABC$ and $\triangle DEF$ such that $\triangle ABC \cong \triangle DEF$, and suppose that we have points G on (BC) and H on (EF) such that $[AG]$ and $[DH]$ bisect $\angle BAC$ and $\angle EDF$ respectively. Prove that $\triangle GAB \cong \triangle HDE$.
6. Conversely, in the setting of the previous exercise suppose that we are not given the condition $\triangle ABC \cong \triangle DEF$, but we are given that $\triangle GAB \cong \triangle HDE$. Prove that $\triangle ABC \cong \triangle DEF$.
7. Let K be a convex subset of \mathbf{R}^n . A point X in K is said to be an **extreme point** of K if it is not between two other points of K . Suppose that T is an affine transformation of \mathbf{R}^n , and suppose that T maps the convex set K onto the convex set L . Prove that T maps the extreme points of K to the extreme points of L .
8. Suppose that T is the affine transformation of \mathbf{R}^n given by $T(\mathbf{x}) = L(\mathbf{x}) + \mathbf{v}$, where L is an invertible linear transformation and \mathbf{v} is a fixed vector in \mathbf{R}^n ; In terms of coordinates, if L is given by the $n \times n$ matrix $A = (a_{ij})$ and $\mathbf{v} = (b_1, \dots, b_n)$, then one has the following expression for $\mathbf{y} = T(\mathbf{x})$ in terms of coordinates:

$$y_i = a_{i,1} x_1 + \dots + a_{i,n} x_n + b_i$$

Let DT be the **derivative matrix** whose (i, j) entry is given by

$$\frac{\partial y_i}{\partial x_j}$$

- (a) Show that the (i, j) entry of DT is equal to a_{ij} .
- (b) If T_1 and T_2 are affine transformations of \mathbf{R}^n , explain why $D(T_1 \circ T_2)$ is the matrix product $D(T_1)D(T_2)$. [**Hint:** Expand the composite $T_1 \circ T_2$.]
- (c) Explain why T is a translation if and only if $D(T)$ is the identity matrix.
- (d) If T is a translation and S is an arbitrary affine transformation, prove that the composite $S^{-1} \circ T \circ S$ is a translation. What is its value at the vector $\mathbf{0}$?
9. The **vertical reflection** $S(c)$ about the horizontal line $y = c$ in \mathbf{R}^2 is the affine map defined by $(x_1, x_2) \mapsto (x_1, 2c - x_2)$. Show that $S(c)$ sends the horizontal line into

itself and interchanges the horizontal lines $y = 2c$ and $y = 0$. Prove that the composite of two vertical reflections $\mathbf{S}(a)\mathbf{S}(b)$ is a translation, and the composite of three vertical reflections $\mathbf{S}(a)\mathbf{S}(b)\mathbf{S}(c)$ is a vertical reflection $\mathbf{S}(d)$; evaluate d explicitly. [**Hints:** For the twofold composite, what is the derivative matrix? Also, explain why the twofold composite $\mathbf{S}(a)\mathbf{S}(b)$ sends $(0, 0)$ to a point whose first coordinate is equal to zero.]

10. Let \mathbf{A} be the orthogonal matrix

$$\begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$$

where θ is a real number. Show that there is an orthonormal basis $\{\mathbf{u}, \mathbf{v}\}$ for \mathbf{R}^2 such that $\mathbf{A}\mathbf{u} = \mathbf{u}$ and $\mathbf{A}\mathbf{v} = -\mathbf{v}$.

11. Let \mathbf{A} be the orthogonal rotation matrix

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

where θ is a real number which is not an integral multiple of 2π , let $\mathbf{b} \in \mathbf{R}^2$, and let \mathbf{T} be the Galilean transformation $\mathbf{T}(\mathbf{x}) = \mathbf{A}\mathbf{x} + \mathbf{b}$. Prove that there is a unique vector \mathbf{z} such that $\mathbf{T}(\mathbf{z}) = \mathbf{z}$. [**Hint:** \mathbf{A} is not the identity matrix, and in fact $\mathbf{A} - \mathbf{I}$ is invertible; prove the latter assertion.]

II.5 : Euclidean parallelism

Supplementary background readings.

Ryan : pp. 11 – 15, 17 – 18, 49 – 50, 58 – 59, 68

Exercises to work.

1. Suppose that \mathbf{L} and \mathbf{M} are skew lines in \mathbf{R}^3 . Prove that there is a unique plane \mathbf{P} such that \mathbf{L} is contained in \mathbf{P} and \mathbf{M} is parallel to (*i.e.*, disjoint from) \mathbf{P} . [**Hints:** Write \mathbf{L} and \mathbf{M} as $\mathbf{x} + \mathbf{V}$ and $\mathbf{y} + \mathbf{W}$, where \mathbf{V} and \mathbf{W} are 1 – dimensional vector subspaces. Since \mathbf{L} and \mathbf{M} are not parallel, we know that \mathbf{V} and \mathbf{W} are distinct. Let \mathbf{U} be the vector subspace $\mathbf{V} + \mathbf{W}$. Why is \mathbf{U} a 2 – dimensional subspace? Set $\mathbf{P} = \mathbf{x} + \mathbf{U}$ and verify that \mathbf{P} has the desired properties; in particular, if \mathbf{M} and \mathbf{P} have a point \mathbf{z} in common, note that $\mathbf{M} = \mathbf{z} + \mathbf{W}$ and $\mathbf{P} = \mathbf{z} + \mathbf{U}$.]

2. Suppose that \mathbf{P} and \mathbf{Q} are parallel planes in \mathbf{R}^3 , and let \mathbf{S} be a plane which meets each of them in a line. Prove that the lines of intersection $\mathbf{S} \cap \mathbf{P}$ and $\mathbf{S} \cap \mathbf{Q}$ must be parallel.
3. Suppose that we are given three distinct planes \mathbf{S} , \mathbf{P} and \mathbf{Q} in \mathbf{R}^3 such that \mathbf{S} is parallel to both \mathbf{P} and \mathbf{Q} . Prove that \mathbf{P} and \mathbf{Q} are parallel. [*Hint:* Through a given point \mathbf{x} not on \mathbf{S} , how many planes are there that pass through \mathbf{x} and are parallel to \mathbf{S} ?]
4. Suppose that \mathbf{v} and \mathbf{w} are linearly independent vectors in \mathbf{R}^3 , and let \mathbf{z} be a third vector in \mathbf{R}^3 . Let a and b be real numbers. Prove that $f(a, b) = |\mathbf{z} - a\mathbf{v} - b\mathbf{w}|$ takes a minimum value when $\mathbf{z} - a\mathbf{v} - b\mathbf{w}$ is perpendicular to both \mathbf{v} and \mathbf{w} . [*Hint:* Take an orthonormal basis \mathbf{e}, \mathbf{f} for the span of \mathbf{v} and \mathbf{w} , and let $\mathbf{Q}(\mathbf{z}) = \mathbf{z} - \langle \mathbf{z}, \mathbf{e} \rangle \mathbf{e} - \langle \mathbf{z}, \mathbf{f} \rangle \mathbf{f}$. Why is \mathbf{z} perpendicular to $\mathbf{e}, \mathbf{f}, \mathbf{v}$, and \mathbf{w} ? Find the length squared of the vector
- $$\mathbf{z} - s\mathbf{e} - t\mathbf{f} = \mathbf{Q}(\mathbf{z}) - [s - \langle \mathbf{z}, \mathbf{e} \rangle]\mathbf{e} - [t - \langle \mathbf{z}, \mathbf{f} \rangle]\mathbf{f}$$
- using the fact that \mathbf{e}, \mathbf{f} and $\mathbf{Q}(\mathbf{z})$ are mutually perpendicular. Why is the (square of the) length minimized when the coefficients of \mathbf{e} and \mathbf{f} are zero? Why is the minimum value of $g(s, t) = |\mathbf{z} - s\mathbf{e} - t\mathbf{f}|$ equal to the minimum value of $f(a, b) = |\mathbf{z} - a\mathbf{v} - b\mathbf{w}|$?
5. Suppose that we have two skew lines in \mathbf{R}^3 of the form $\mathbf{0a}$ and \mathbf{bc} . Let \mathbf{x} and \mathbf{y} be points of $\mathbf{0a}$ and \mathbf{bc} respectively. Prove that the distance $d(\mathbf{x}, \mathbf{y}) = |\mathbf{x} - \mathbf{y}|$ is minimized when $\mathbf{x} - \mathbf{y}$ is perpendicular to \mathbf{a} and $\mathbf{c} - \mathbf{b}$. (In other words, *the shortest distance between the two skew lines is along their common perpendicular.*)
6. (Ryan, Exercise 38, p. 36) Let $\mathbf{W}, \mathbf{X}, \mathbf{Y}, \mathbf{Z}$ be four points in \mathbf{R}^3 , no three of which are collinear, let $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ be the midpoints of $\mathbf{WX}, \mathbf{XY}, \mathbf{YZ}$, and \mathbf{ZW} , and suppose that we have $\mathbf{AB} \neq \mathbf{CD}$ and $\mathbf{AD} \neq \mathbf{BC}$. Prove that $\mathbf{AB} \parallel \mathbf{CD}$ and $\mathbf{AD} \parallel \mathbf{BC}$.
7. (Ryan, Exercise 30, p. 68) Let $\angle \mathbf{ABC}$ be given, and let \mathbf{D} lie in the interior of $\angle \mathbf{ABC}$. Prove that \mathbf{D} lies on an open segment (\mathbf{XY}) , where $\mathbf{X} \in (\mathbf{BA})$ and $\mathbf{Y} \in (\mathbf{BC})$.