

## SOLUTIONS TO ADDITIONAL EXERCISES FOR III.1 AND III.2

Illustrations to accompany these solutions are given on the last page.

D1. We shall follow the hints. Take a basis  $B$  for  $V$  (which has  $r$  elements) and extend it to a basis for  $\mathbb{R}^n$  by adding a suitable set of  $n - r$  vectors  $A$ ; order the basis so that the elements of  $B$  come first. If we apply Gram-Schmidt process to obtain an orthonormal basis  $C$  of  $\mathbb{R}^n$  from  $B \cup A$ , then by construction the first  $r$  vectors in  $C$  will form an orthonormal basis for  $V$ . Let  $A'$  be the last  $n - r$  vectors in  $C$ ; we claim that  $A'$  forms an orthonormal basis for  $V^\perp$ .

First of all, every vector in  $A'$  lies in  $V^\perp$ , for every vector in  $V$  has the form  $\sum_{j \leq r} t_j \mathbf{c}_j$  and the dot product of such a vector with  $\mathbf{c}_k$  is zero if  $k > r$ . Therefore  $V^\perp$  contains the  $(n - r)$ -dimensional vector subspace spanned by  $A'$ . To see that nothing else can be contained in  $V^\perp$ , consider a vector  $\mathbf{y}$  which is not a linear combination of the vectors in  $A'$ . Since  $C$  is an orthonormal basis, we must have  $\mathbf{y} = \sum_{j \leq n} t_j \mathbf{c}_j$  where  $t_m \neq 0$  for some  $m \leq k$ . But the latter implies that  $\mathbf{y} \cdot \mathbf{c}_m = t_m \neq 0$ , and therefore  $\mathbf{y}$  cannot lie in  $V^\perp$ . Thus the vectors in  $A'$  form a basis of this subspace and hence its dimension is  $n - r$ .

To conclude, as noted in the hint it suffices to prove that  $V$  is a vector subspace of  $(V^\perp)^\perp = V$  and the dimensions of these two subspaces are equal. The first statement follows since  $\mathbf{v} \in V$  implies  $\mathbf{v} \cdot \mathbf{x} = 0$  for all  $\mathbf{x} \in V^\perp$ , and the first follows because the dimension of  $(V^\perp)^\perp = V$  is equal to

$$n - \dim V^\perp = n - (n - r) = r = \dim V.$$

Since  $V_1 \subset V_2$  and  $\dim V_1 = \dim V_2$  imply  $V_1 = V_2$ , the equality of  $V$  and  $(V^\perp)^\perp = V$  follows immediately. ■

*Note.* One important consequence of the preceding exercise is the following: *If  $V$  and  $W$  are vector subspaces of  $\mathbb{R}^n$  such that  $V \neq W$ , then  $V^\perp \neq W^\perp$ .* — For if  $V^\perp = W^\perp$ , then their orthogonal complements, which by the exercise are  $V$  and  $W$  respectively, would also have to be equal.

D2. By the preceding exercise we know that  $\dim V^\perp = 2$  and  $\dim W^\perp = 1$ . Furthermore, since  $V$  and  $W^\perp$  are distinct 1-dimensional subspaces, it follows that the dimension of their intersection is strictly less than 1 and hence the intersection must be  $\{\mathbf{0}\}$ .

Since  $V$  and  $W^\perp$  are distinct 1-dimensional vector subspaces, it follows that their orthogonal complements  $V^\perp$  and  $(W^\perp)^\perp = W$  are distinct 2-dimensional vector subspaces (see the note following the solution of D1). Therefore the linear sum  $V^\perp + W$  properly contains each of them (otherwise they would be equal), so its dimension is at least 3; since we are in  $\mathbb{R}^3$ , the dimension must be exactly 3 and the linear sum is just  $\mathbb{R}^3$ . Applying the Dimension Formula we see that

$$\dim W \cap V^\perp = \dim W + \dim V^\perp - \dim \mathbb{R}^3 = 2 + 2 - 3 = 1$$

D3. Write the line and plane as  $\mathbf{x} + V$  and  $\mathbf{x} + W$  respectively; the assumptions imply that  $V$  is not equal to  $W^\perp$  and hence  $M = \mathbf{x} + (W \cap V^\perp)$  is a line which is contained in both  $P = \mathbf{x} + W$  and in the plane  $Q = \mathbf{x} + V^\perp$ . Since  $Q$  is the unique plane through  $\mathbf{x}$  which is perpendicular to  $L$ , it follows that  $M$  has the properties described in the statement of the exercise.

To see that there is only one line, suppose that  $M'$  has the required properties. Then it follows that  $M' \subset Q$ , and since  $M' \subset P$  is assumed we know that  $M'$  is contained in  $P \cap Q$ ; since the latter is a line, it follows that we have  $M' = P \cap Q$ , and since the intersection is  $M$  we have  $M' = M$ . ■


D4. The condition  $a < 2x$  follows from the Triangle Inequality for triples of non-collinear points. Conversely, if we have  $a < 2x$ , then we also have

$$0 < h = \sqrt{x^2 - \frac{a^2}{4}}.$$

By the Protractor and Ruler Postulates we can construct a right triangle  $\triangle ABC$  such that  $AB \perp BC$ ,  $d(A, B) = a/2$ , and  $d(B, C) = h$ . By the Pythagorean Theorem we know that  $d(A, C) = x$ . Now take  $D \in (AB)$  such that  $d(A, D) = a$ . It then follows that  $d(B, D) = a/2$  and by **SAS** and perpendicularity we have  $\triangle ABC \cong \triangle DBC$ . It follows that  $d(D, C) = d(A, C) = x$ , and therefore the triangle  $\triangle ABC$  is an isosceles triangle such that the lengths of two sides are equal to  $x$  and the length of the third side is equal to  $a$ . ■


## FIGURE FOR SOLUTIONS TO ADDITIONAL EXERCISES, SET D

D3.



The idea in the hint is to show that **M** is the intersection of **P** with the plane **Q** through **X** such that  $L \perp Q$ . In the drawing the perpendicular projection of **L** onto the plane **P** is drawn in pink. Observe that this projection **N** is a line through **X** and **M** is also the line through **X** which is perpendicular to the plane of **L** and **N** (try to prove this assertion using vectors — it is not particularly difficult!).

D4.



Since  $2x > a$  it follows that  $x^2 - [a^2/4]$  is positive and hence one can construct a right triangle whose sides have lengths  $a$  and  $x^2 - [a^2/4]$ . The hypotenuse of such a triangle must have length equal to  $x$  by the Pythagorean Theorem. The second drawing indicates what should happen if we take the mirror image of this triangle with respect to the line containing the side of length  $x^2 - [a^2/4]$ . In order to complete the proof it is necessary to give reasons why this picture is accurate and one obtains an isosceles triangle with the desired properties.