## SOLUTIONS TO ADDITIONAL EXERCISES FOR II.1 AND II.2

Here are the solutions to the additional exercises in triangle-exercises.pdf. Illustrations to accompany these solutions are given in the online file

## trianglefigures.pdf

in the course directory.

C1. Suppose first that X is one of the vertices A, B, C. In these cases the conclusion follows because  $A \in AB \cap AC$  and each of these lines contains an infinite number of points on the triangle (namely, all points of [AB] and [AC] respectively),  $B \in AB \cap BC$  and each of these lines contains an infinite number of points on the triangle (namely, all points of [AB] and [BC] respectively), and finally  $C \in BC \cap AC$  and each of these lines contains an infinite number of points on the triangle (namely, all points of [BC] and [AC] respectively).

Suppose now that X is not a vertex. Without loss of generality we may assume that  $X \in (AB)$ , for the remaining cases where  $X \in (BC)$  or  $X \in (AC)$  follow by interchanging the roles of A, B, C in the argument we shall give.

If  $X \in (AB)$  and L = AB, then clearly  $X \in L$  and L contains infinitely many points of the triangle because it contains [AB]. From now on, suppose that  $X \neq AB$ .

If L = XC, then X and C lie on both L and the triangle; we claim that no other point of L satisfies these conditions. Suppose to the contrary that there is such a third point Y; there are three cases depending upon whether Y lies on AB, BC or AC. If  $Y \in AB$ , then both L and AB contain the distinct points X and Y, so that L = XY; but we are assuming that X, Y, C are collinear, and this contradicts our even more basic assumption that A, B, C are noncollinear (this is implicit in asserting the existence of  $\Delta ABC$ ). Therefore the line XC only meets the triangle in two points.

Now suppose that  $C \notin L$  and  $L \neq AB$ ; we need to show that L and  $\Delta ABC$  have at most one other point in common besides X. By Pasch's Theorem there is a second point Y on L which lies on either (BC) or (AC); in either case there we claim that there is no third point in  $L \cap \Delta ABC$ . Since  $X \in AB$  is not one of the vertices and the lines BC and AC meet AB in B and A respectively, it follows that X lies on neither of these lines. Therefore the line L = XY meets  $\Delta ABC$  in two sides and cannot contain any of the vertices. If there were a third point, it would lie on one of (AB), (BC) or (AC). By Exercises II.2.8 we know that L cannot contain points of all three sides,, and if the third point were in (AB) it would follow that L = AB. On the other hand, the line L cannot contain X and two points from either (AC) or (BC), for in that case L would be equal to AC or BC and also contain  $X \in (AB)$ , so that L would also be equal to AB. Thus the existence of a third point leads to a contradiction if  $L \neq AB, XC$ , and hence no such point can exist, so that all lines through X except AB meet the triangle in two points.

C2. The most difficult parts of this proof were done in the preceding exercise. Let **T** be equal to  $\Delta ABC = \Delta DEF$ . By the preceding exercise, since  $\mathbf{T} = \Delta ABC$  we

know that  $\{A, B, C\}$  is the set **V** of all points X in **T** such that two lines through X contain at least three points of **T**, and likewise  $\{D, E, F\}$  is the set **V** of all points X in **T** such that two lines through X contain at least three points of **T**. Therefore we have  $\{A, B, C\} = \mathbf{V} = \{D, E, F\}$ .

C3. Let  $H_1$  and  $H_2$  denote the two half-planes associated to L. Then each of the points A, B, C lies on exactly one of the subsets  $L, H_1, H_2$ .

Before we split the argument into cases using the previous sentence, we make a general observation. Since  $X \in L$  lies in the interior of  $\Delta ABC$ , by the Crossbar Theorem we know that (BX and (AC)) have a point Y in common. This point cannot be X because a point cannot lie in both the interior of  $\Delta ABC$  and one of the three sides AB, BC, AC (look at the definition of interior). Since  $Y \in (BC)$ , it follows that either B \* X \* Y or B \* Y \* X is true; however, if the latter were true, then B and X would lie on opposite sides of the line AY = AC, contradicting the assumption on X. Therefore we must have B \* X \* Y.

We claim that all three vertices cannot lie in either  $H_1$  or  $H_2$ . If they did, then by convexity the point Y in the preceding paragraph would also lie in the given half-plane, and similarly the point X would lie in this half-plane. Since  $X \in L$  by assumption, this is impossible, and thus the three vertices cannot all lie on one side of L.

Next, we claim that at most one vertex lies on L. If, say,  $A \in L$ , then L = AX, and if either B or C were also on L we would have that L = AB or AC, which in turn would imply that  $X \in AB$  or AC, contradicting the condition that X lies in the interior of the triangle. The cases where  $B \in L$  and  $C \in L$  can be established by interchanging the roles of the three vertices in the preceding argument.

Suppose now that one vertex lies on L; we claim that the other two vertices must lie on opposite sides of L. Once again, it is enough to consider the case where  $A \in L$ , for the other cases will follow by interchanging the roles of the vertices. But if  $A \in L$ , then the Crossbar Theorem implies that L and (BC) have a point W in common (in fact, the open segment (BC) and open ray (AX) have a point in common), and therefore it follows that B and C lie on opposite sides of L. Furthermore, it follows that the line L meets the triangle in the distinct points A and W.

The only possibility left to consider is the case where no vertex lies on L; by the preceding discussion, we know that neither half-plane contains all three vertices, and thus two of the vertices are on one half-plane and one is on the other. As before, without loss of generality we may assume that A is on one side and B, C are on the other. But in this situation we know that the line L meets both (BC) and (AC). This completes the examination of all possible cases.

C4. We shall follow the hint and solve for  $\mathbf{x}$  in terms of  $\mathbf{y}$ . Since  $\mathbf{y} = A\mathbf{x} + \mathbf{b}$  and A is invertible, it follows that  $\mathbf{x} = A^{-1}(\mathbf{y} - \mathbf{b})$ . If we substitute this into the defining equation for the plane, we see that

$$d = C\mathbf{x} = CA^{-1}(\mathbf{y} - \mathbf{b})$$
 or equivalently  $CA^{-1}\mathbf{y} = d + CA^{-1}\mathbf{b}$ 

which shows that  $\mathbf{y}$  is defined by an equation of the form  $P\mathbf{y}=q$ , where  $P=CA^{-1}$  and  $q=d+CA^{-1}\mathbf{b}.\blacksquare$ 

C5. First of all, we claim that C and D lie on opposite sides of the line AB. Since no three of A, B, C, D are collinear, it follows that neither C nor D lies on AB, so it is only necessary to prove that C and D cannot lie on the same side of AB. However, if they did, then the noncollinearity condition would imply that (AC and (AD are distinct open rays, and by the Trichotomy Principle it would follow that either D would lie in the interior of  $\triangle BAC$  or else C would lie in the interior of  $\triangle BAD$ .

By the preceding paragraph and Plane Separation, we know that the open segment (CD) meets the line AB in some point E, and since C\*E\*D is true it follows that E lies in the interior of  $\angle CAD$ . Now B, E, A are collinear; since B does not lie in the interior of  $\angle CAD$  and (BE does lie in the interior of this angle, it follows that we must have E\*A\*B.

By the Supplement Postulate for angle measurements we have

$$|\angle BAC| + |\angle CAE| = 180^{\circ} = |\angle DAB| + |\angle DAE|$$

and since E lies in the interior of  $\angle CAD$  the Addition Postulate implies that

$$|\angle CAD| = |\angle CAE| + |\angle DAE|$$
.

If we add the equations which follow from the Supplement Postulate we have

$$|\angle BAC| + |\angle CAE| + |\angle DAB| + |\angle DAE| = 360^{\circ}$$

and if we now use the remaining equation we may rewrite the left hand side of the latter as  $|\angle BAC| + |\angle CAD| + |\angle DAB| = 360^{\circ}$ , which is what we wanted to prove.