

EXERCISES FOR MATHEMATICS 138A

WINTER 2004

The references denote sections of the text for the course:

M. P. do Carmo, *Differential geometry of Curves and Surfaces*, Prentice-Hall, Saddle River NJ, 1976, ISBN 0-132-12589-7.

I. Classical Differential Geometry of Curves

I.1 : Cross products

(O'Neill, § 2.2)

Additional exercise

1. Verify that the cross product of vectors in \mathbf{R}^3 satisfies the *Jacobi identity*:

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) + \mathbf{b} \times (\mathbf{c} \times \mathbf{a}) + \mathbf{c} \times (\mathbf{a} \times \mathbf{b}) = \mathbf{0} .$$

I.2 : Parametrized curves

(O'Neill, § 1.4)

O'Neill, pp.21-22: 2, 8

Additional exercises

1. Find a parametrized curve $\alpha(t)$ which traces out the unit circle about the origin in the coordinate plane and has initial point $\alpha(0) = 1$.

2. Let $\alpha(t)$ be a parametrized curve which does not pass through the origin. If $\alpha(t_0)$ is the point in the image that is closest to the origin and $\alpha'(t_0) \neq 0$, show that $\alpha(t_0)$ and $\alpha'(t_0)$ are perpendicular.

3. Two lines are said to be *skew lines* if they are disjoint but not parallel. Prove that the distance between the skew lines $\mathbf{x}(t) = \mathbf{x}_0 + t\mathbf{u}$ and $\mathbf{y}(t) = \mathbf{y}_0 + t\mathbf{v}$ is given by

$$\rho = \frac{\mathbf{u} \times \mathbf{v} \cdot \mathbf{r}}{\|\mathbf{u} \times \mathbf{v}\|}$$

where $\mathbf{r} = \mathbf{x}_0 - \mathbf{y}_0$. [*Hints*: The shortest distance between the lines is given by a common perpendicular. You may assume the existence of a common perpendicular when working the problem. It might be helpful to let \mathbf{x}_1 and \mathbf{y}_1 from these lines lie on this common perpendicular.]

4. Prove that a regular smooth curve lies on a straight line if and only if there is a point that lies on all its tangent lines.

I.3 : Arc length and reparametrization

(O'Neill, §§ 1.4, 2.2)

O'Neill, pp. 56-57: 3-5, 10, 11

Additional exercises

1. Prove that a necessary and sufficient condition for the plane $\mathbf{N} \cdot \mathbf{x} = 0$ to be parallel to the line $\mathbf{x} = \mathbf{x}_0 + t \cdot \mathbf{u}$ is for \mathbf{N} and \mathbf{u} to be perpendicular.

2. (a) Given $a > 0$, consider the set of all continuously differentiable real valued functions f on $[0, 1]$ such that $f(0) = 0$ and $f(1) = a > 0$. Define $L(f)$ by the formula $L(f) = \int_0^a |f'(t)| dt$. Show that the minimum value of $L(f)$ is a , and if equality holds then f' is everywhere nonnegative. [Hints: Since $f' \leq |f'|$ a similar inequality holds for their definite integrals. This inequality of integrals is strict if and only if $f'(t) < |f'(t)|$ for some t , which happens if and only if $f'(t) < 0$ for that choice of t .]

(b) Let ρ , θ and ϕ denote the usual spherical coordinates, and suppose we have a curve on the sphere of radius 1 about the origin with parametric equations of the form

$$\mathbf{x}(t) = (\cos \theta(t) \sin \phi(t), \sin \theta(t) \sin \phi(t), \cos \phi(t))$$

for continuously differentiable functions $\theta(t)$ and $\phi(t)$. Prove that the length of this curve is given by the formula

$$\int_a^b \sqrt{(\theta'(t))^2 + \sin^2 \theta(t) (\phi'(t))^2} dt$$

where the curve is defined on $[a, b]$.

(c) Show that among all regular smooth curves \mathbf{x} that are defined on $[0, 1]$, have images on the unit sphere, and connect the points $(1, 0, 0)$ and $(\cos a, \sin a, 0)$ for some $a < \pi$, the curve of shortest length is given by the great circle arc joining the endpoints, and that any other curve with this length is a weak reparametrization of the great circle arc (*i.e.*, if α is the standard great circle arc, then any other curve β must have the form $\beta(t) = \alpha(f(t))$, where f is a 1-1 function from $[0, 1]$ to $[0, a]$ that is continuously differentiable and satisfies $f' \geq 0$. [Hints: Let \mathbf{y} be the curve in the xy -plane obtained from \mathbf{x} by replacing $\phi(t)$ with $\pi/2$; in other words, \mathbf{y} is the perpendicular projection of the original curve onto the xy -plane. Why does the spherical coordinate arc length formula show that the length of \mathbf{x} is greater than or equal to the length of \mathbf{y} ? And why is there strict inequality if $\phi'(t_0) \sin \theta(t_0) \neq 0$ somewhere? Why does this mean that the plane curve $(\cos \theta(t), \sin \theta(t), 0)$ is a weak reparametrization of $(\cos at, \sin at, 0)$? Recall that by continuity the latter implies $\phi'(t) \neq 0$ for all t sufficiently close to t_0 . What does part (a) imply if ϕ is constant?]

Note. The final part of the problem is a special case of the well known result that the shortest curve on a sphere joining two points is given by the smaller of the arcs on the great circle through the points; in fact, one can use this special case to prove the general statement. [A file containing a detailed proof may be inserted into the course directory eventually.]

I.4 : Curvature and torsion

(O'Neill, § 2.3)

Additional exercises

1. Suppose a curve is given in polar coordinates by $r = r(\theta)$ where $\theta \in [a, b]$.

(i) Show that the arc length is $\int_a^b \sqrt{r^2 + (r')^2} d\theta$.

(ii) Show that the curvature is

$$k(\theta) = \frac{2(r')^2 - rr'' + r^2}{[r^2 + (r')^2]^{3/2}}.$$

2. Let α and β be regular parametrized curves such that β is the arc length reparametrization of α . Let t be the parameter for α and s for β . Prove the following:

(a) $dt/ds = 1/|\alpha'|$, $d^2t/ds^2 = -(\alpha' \cdot \alpha''/|\alpha'|^4)$

(b) The curvature is given by

$$k(t) = \frac{\alpha' \times \alpha''}{|\alpha'|^3}$$

(c) The torsion is given by

$$\tau(t) = -\frac{\alpha' \times \alpha'' \cdot \alpha'''}{|\alpha' \times \alpha''|^2}$$

(d) If α has coordinate functions x and y , then the signed curvature of α at t is equal to

$$k(t) = \frac{x'y'' - x''y'}{[(x')^2 + (y')^2]^{3/2}}$$

3. Show that the curvature of a regular parametrized curve α at t_0 is equal to the curvature of the plane curve γ which is the perpendicular projection of α onto the osculating plane of α at t_0 .

4. Consider the problem of designing a set of railroad tracks that contains a pair of parallel tracks along with a third going from the first to the second smoothly. Mathematically, the parallel tracks themselves may be viewed as corresponding to the parallel lines $y = 0$ and $y = 1$ in the coordinate plane, and the track going from one to the other may be viewed as a regular smooth curve that is the graph of a twice differentiable function f such that $f(x)$ is zero if $t \leq 0$, $f(x) = 1$ if $t \geq 1$, and on $[0, 1]$ the function f is given by a polynomial $p(x)$. The existence of a second derivative ensures that the slope of the tangent line would be a continuous function of x , and in addition we want to assume that *the curvature is also a continuous function of x* . Find a polynomial $p(x)$ of degree 5 such that all the required conditions are fulfilled. [*Hint:* If we are given a graph curve with parametric equations $(t, y(t))$, then the curvature at parameter value t is given by the formula

$$k(t) = \frac{|y''|}{(1 + (y')^2)^{3/2}}$$

and one step in the argument is to use this fact to compute $p''(0)$ and $p''(1)$. In fact, the conditions of the problem uniquely specify the values of p and its first and second derivatives at both 0 and 1. Why does this mean the only values to find are the coefficients of x^3 , x^4 and x^5 ?

Optional. Graph the function f using calculator or computer graphics.

I.5 : Frenet-Serret Formulas

(O'Neill, §§ 2.3–2.4)

O'Neill, pp. 64–66: 1, 5

Additional exercises

1. Let \mathbf{x} be a regular smooth curve with a continuous third derivative, and let $(\mathbf{T}, \mathbf{N}, \mathbf{B})$ be its Frenet trihedron. Prove that there is a vector \mathbf{W} (the *Darboux vector*) such that $\mathbf{T}' = \mathbf{W} \times \mathbf{T}$, $\mathbf{N}' = \mathbf{W} \times \mathbf{N}$, and $\mathbf{B}' = \mathbf{W} \times \mathbf{B}$. What is the length of \mathbf{W} ?

2. If \mathbf{x} is defined for $t > 0$ by the formula

$$\mathbf{x}(t) = \left(t, \frac{1+t}{t}, \frac{1-t^2}{t} \right)$$

show that \mathbf{x} is planar.

II. Topics from Multivariable Calculus and Geometry

II.1 : Differential forms

(O'Neill, §§ 1.5–1.6)

O'Neill, pp. 25–26: 5, 6 (first part only), 9 (last sentence only)

O'Neill, pp. 31–32: 1, 3–5

Additional exercise

1. Suppose that ω is a 2-form on \mathbf{R}^3 such that $\omega \wedge dx = 0$. Explain why there is a 1-form θ such that $\omega = \theta \wedge dx$.

II.2 : Smooth mappings

(O'Neill, §§ 1.7, 3.2)

Additional exercises

Definition. A subset K of \mathbf{R}^n is said to be *convex* if whenever \mathbf{x} and \mathbf{y} lie in K then the whole line segment defined by the parametrized curve $\mathbf{x} + t(\mathbf{y} - \mathbf{x})$ for $t \in [0, 1]$ is contained in K .

1. Prove that an open convex set is a connected domain [*Hint:* Imitate the proof for the set of all point whose distance from some point \mathbf{p} is less than some positive number r].

2. Show by example that an intersection of two connected domains in \mathbf{R}^2 is not necessarily a connected domain. [*Hint:* Let U be the annular region defined by the inequalities $1 < x^2 + y^2 < 9$

and let V be the horizontal strip defined by the inequality $|y| < \frac{1}{2}$. Verify that U is arcwise connected using the polar coordinate mapping, which yields a continuous 1-1 mapping from the convex set $(1, 3) \times [0, 2\pi)$ onto U . If $U \cap V$ were connected then by a result in the Appendix to Chapter 5 in do Carmo, it would also be arcwise connected. Suppose now that \mathbf{x} is a curve joining the points $(\pm 2, 0)$. By the Intermediate Value Theorem there must be some parameter value t_0 such that the first coordinate of $\mathbf{x}(t_0)$ is equal to zero. Why does this mean that \mathbf{x} cannot lie entirely inside $U \cap V$?

3. Given an matrix A with real entries, let $|A|$ denote the Euclidean length given by the square root of the standard sum $\sum_{i,j} |a_{i,j}|^2$. If P and Q are two matrices with real entries such that the product PQ can be defined, prove that $|PQ| \leq |P| \cdot |Q|$.

4. Let U be a convex connected domain in \mathbf{R}^n , and let $f : U \rightarrow \mathbf{R}^m$ be a smooth \mathcal{C}^1 function.

(a) Prove that

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 ([Df(\mathbf{x} + t(\mathbf{y} - \mathbf{x}))](\mathbf{y} - \mathbf{x})) dt$$

for all $\mathbf{x}, \mathbf{y} \in U$. [*Hint:* Explain why the integrand is the derivative of the function

$$f(\mathbf{x} + t(\mathbf{y} - \mathbf{x}))$$

using the Chain Rule.]

(b) Suppose that the derivative matrix function Df satisfies $|Df| \leq M$ on U . Prove that

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq M \cdot |\mathbf{y} - \mathbf{x}|$$

for all $\mathbf{x}, \mathbf{y} \in U$.

Note. An inequality of this sort is called a *Lipschitz condition*.

II.3 : Inverse and Implicit Function Theorems

(O'Neill, § 1.7)

Additional exercises

1. Suppose that $f : \mathbf{R} \rightarrow \mathbf{R}$ is a \mathcal{C}^r function such that its derivative f' is everywhere positive and the limits of $f(t)$ as $t \rightarrow \pm\infty$ are $\pm\infty$ respectively. Prove that f has a \mathcal{C}^r inverse function.

2. Prove that $F(x, y) = (e^x + y, x - y)$ defines a 1-1 onto \mathcal{C}^∞ map from \mathbf{R}^2 to itself with a \mathcal{C}^∞ inverse.

3. Prove that $F(x, y) = (xe^y + y, xe^y - y)$ defines a 1-1 onto \mathcal{C}^∞ map from \mathbf{R}^2 to itself with a \mathcal{C}^∞ inverse.

4. (a) Using the change of variables formula, explain briefly why the area of a set in \mathbf{R}^2 is the same as the area of its image under a rigid motion of the form $T(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$, where A is a rotation matrix

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

(b) More generally, if we are given an arbitrary **affine** transformation as above, where the only condition on A is invertibility, how is the area of a set \mathcal{F} related to the area of its image $T(\mathcal{F})$?

5. A smooth \mathcal{C}^r mapping f from a connected domain $U \subset \mathbf{R}^2$ into \mathbf{R}^2 is said to be *regularly conformal* at $\mathbf{p} = (u_0, v_0) \in U$ if the Jacobian of f is positive and for all regular smooth curve pairs \mathbf{x} and \mathbf{y} satisfying $\mathbf{x}(s_0) = \mathbf{y}(s_0) = \mathbf{p}$ the angle between $\mathbf{x}'(s_0)$ and $\mathbf{y}'(s_0)$ is equal to the angle between $[f \circ \mathbf{x}]'(s_0)$ and $[f \circ \mathbf{y}]'(s_0)$.

(a) Prove that the partial derivatives of the coordinate functions satisfy the *Cauchy-Riemann equations*:

$$\frac{\partial f_1}{\partial x_1} = \frac{\partial f_2}{\partial x_2}, \quad \frac{\partial f_2}{\partial x_1} = -\frac{\partial f_1}{\partial x_2}$$

[*Hint:* If $A = Df(\mathbf{p})$, one needs to show that $\cos \angle(A\mathbf{x}, A\mathbf{y}) = \cos \angle(\mathbf{x}, \mathbf{y})$ for all nonzero vectors \mathbf{x} and \mathbf{y} . Let \mathbf{a}_1 and \mathbf{a}_2 denote the columns of A , and let J denote counterclockwise rotation through $\pi/2$. Why is $\mathbf{a}_2 = cJ(\mathbf{a}_1)$ for some constant c , and why does the determinant condition imply c is positive? Explain why $A(\mathbf{e}_1 + \mathbf{e}_2) = \mathbf{a}_1 + \mathbf{a}_2$ must be perpendicular to $A(\mathbf{e}_1 - \mathbf{e}_2) = \mathbf{a}_1 - \mathbf{a}_2$, and use this to conclude that $c = 1$.]

(b) There is a modified version of this relation that holds among the partial derivatives if the Jacobian is **negative**. State it and explain why it is true. [*Hint:* Consider what happens if one composes f with the reflection map $S(x, y) = (x, -y)$.]

Note. Functions satisfying the Cauchy-Riemann equations are also known as *complex analytic* functions, and they are the central objects studied in complex variables courses.

II.4 : Congruence of geometric figures

(O'Neill, §§ 3.1, 3.4–3.5)

1. Let F be an isometry of \mathbf{R}^n , and let \mathbf{x} and \mathbf{y} be distinct points of \mathbf{R}^n such that $F(\mathbf{x}) = \mathbf{x}$ and $F(\mathbf{y}) = \mathbf{y}$. Suppose that \mathbf{z} is a point on the line joining \mathbf{x} to \mathbf{y} that can be expressed as $\mathbf{z} = t\mathbf{x} + (1-t)\mathbf{y}$ for some scalar t . Prove that $F(\mathbf{z}) = \mathbf{z}$ also holds. [*Hints:* Use the fact that $F(\mathbf{w}) = A(\mathbf{w}) + \mathbf{b}$ for some linear transformation A along with the identity $\mathbf{b} = t\mathbf{b} + (1-t)\mathbf{b}$.]

2. Prove that congruent curves have equal lengths.

III. Surfaces in 3-Dimensional Space

III.1: Mathematical descriptions of surfaces

(O'Neill, §§ 4.1, 4.8)

O'Neill, pp. 132–133: 1, 4bc, 5, 9

Additional exercises

1. Write down equations defining the surfaces given by the following geometric conditions:
 - (a) The set of points that are equidistant from the point $(0, 0, 4)$ and the xy -plane.
 - (b) The set of points that are equidistant from the point $(0, 2, 0)$ and the plane defined by the equation $y = -2$.
 - (c) The set of points that are equidistant from the points $(0, 0, 0)$ and $(1, 0, 0)$.
 - (d) The set of points for which the sum of the distances to $(\pm 1, 0, 0)$ is equal to 5.

2. Let \mathbf{a} , \mathbf{b} and \mathbf{c} be linearly independent vectors in \mathbf{R}^3 . Prove that there is a unique sphere containing these three points and $\mathbf{0}$; *i.e.*, show that the system of equations

$$|\mathbf{x} - \mathbf{a}|^2 = |\mathbf{x} - \mathbf{b}|^2 = |\mathbf{x} - \mathbf{c}|^2 = |\mathbf{x}|^2$$

has a unique solution \mathbf{x} .

3. Find the inverse map to the stereographic projection onto \mathbf{R}^2 described in Example 5.2 of O'Neill, and show how to cover the sphere by two parametrized pieces.

III.2: Parametrizations of surfaces

(O'Neill, § 4.2)

Additional exercises

1. Let $f(x, y, z) = (x + y + z - 1)^2$.
 - (i) What are the critical points and values?
 - (ii) For which c is the level set for c a regular surface?
 - (iii) Same questions for xyz^2 .

2. Let Σ be a geometric regular smooth surface, let U be a connected domain in \mathbf{R}^3 containing Σ , and let $\mathbf{g} : U \rightarrow \mathbf{R}^3$ be a smooth 1–1 onto map such that the Jacobian of \mathbf{g} is nowhere zero (hence it has a global inverse), its image is a connected domain, and more generally the image of any connected subdomain of U is also a connected domain. Prove that $\mathbf{g}(\Sigma)$ is also a geometric regular smooth surface.

III.3: Tangent planes

(O'Neill, § 4.3)

O'Neill, pp. 150–153: 6bc, 10

Additional exercises

0. Show that the tangent plane is the same at all points along a ruling of a cylinder.

Definition. A surface S is said to be *globally convex* at a point \mathbf{p} if all points of S lie on one of the half planes determined by this tangent plane at \mathbf{p} (i.e., if the equation of the tangent plane is $\mathbf{a} \cdot \mathbf{x} = b$, then the points of the surface are completely contained in the set determined by the inequality $\mathbf{a} \cdot \mathbf{x} \leq b$ or the reverse inequality $\mathbf{a} \cdot \mathbf{x} \geq b$). A surface is said to be *strictly globally convex* if in addition for each point \mathbf{p} the intersection of S with the tangent plane consists only of the point \mathbf{p} .

The surface S is said to be *locally convex* or *strictly locally convex* at \mathbf{p} if there is an open disk D containing \mathbf{p} such that $S \cap D$ is globally convex or strictly globally convex.

1. Let \mathbf{X} be a parametrized surface defined on a connected domain U , and let $(a, b) \in U$. Define a level function $L(u, v)$ by $L(u, v) = [\mathbf{X}(u, v), \mathbf{X}_u(a, b), \mathbf{X}_v(a, b)]$ (the vector triple product).

(a) Explain why the surface is locally convex at $\mathbf{p} = \mathbf{X}(a, b)$ if and only if L has a relative maximum or minimum at (a, b) and why the surface is strictly locally convex there if and only if L has a strict relative maximum or minimum.

(b) Why does the gradient of L vanish at (a, b) ?

(c) If $H(a, b)$ is the determinant

$$\begin{vmatrix} [\mathbf{X}_{u,u}(a, b), \mathbf{X}_u(a, b), \mathbf{X}_v(a, b)] & [\mathbf{X}_{u,v}(a, b), \mathbf{X}_u(a, b), \mathbf{X}_v(a, b)] \\ [\mathbf{X}_{v,u}(a, b), \mathbf{X}_u(a, b), \mathbf{X}_v(a, b)] & [\mathbf{X}_{v,v}(a, b), \mathbf{X}_u(a, b), \mathbf{X}_v(a, b)] \end{vmatrix}$$

explain why a surface is **NOT** locally convex at \mathbf{p} if $H(a, b) < 0$. [*Hint:* Why does L have a saddle point at (a, b) ?]

(d) In the notation of the preceding part of the problem, show that the surface is strictly locally convex at \mathbf{p} if $H(a, b) > 0$. [*Hint:* Why does L have a strict local maximum or minimum?]

(e) If \mathbf{X} is a graph parametrization of the form $\mathbf{X}(u, v) = (u, v, f(u, v))$, prove that $H(a, b)$ is a 2×2 determinant of a matrix whose entries are the corresponding second partial derivatives of f at (a, b) .

(f) Apply the preceding to show that if $p \geq 2$ then the graph of the function

$$z = (1 - |x|^p - |y|^p)^{1/p}$$

is strictly locally convex at all (x, y) such that $|x|^p + |y|^p < 1$. In particular, the case $p = 2$ merely states that the usual sphere is strictly locally convex at each point (in fact, all these surfaces are *globally* strictly convex, but we shall not attempt to prove this). [*Hint:* If $r > 1$, explain why the derivative of $|x|^r$ is equal to $r|x|^{r-1}$. There are three cases, depending upon whether x is positive, negative or zero.]

NOTE. By interchanging the roles of the three coordinates in the preceding result one can in fact show that the sets defined by the equations $|x|^p + |y|^p + |z|^p = 1$ are all regular smooth surfaces and are strictly locally convex at all points.

Further study. Graph the intersection of this surface with the xz -plane for $p = 3$ and 4 using calculator or computer graphics. Try this also for larger values of p and describe the limit of these surfaces as $p \rightarrow \infty$.

2. Let S be the cylindrical surface given by the parametric equation(s) $\mathbf{X}(u, v) = (u \cos u, u \sin u, v)$ for $u \in (\pi/2, 9\pi/2)$ and $v \in (-1, 1)$. This is a cylinder generated by the

Archimedean spiral curve in the plane given in polar coordinates by $r = \theta$. Show that S is locally convex at each point but not globally convex at some point in S (for example, at $(2\pi, 0, 0)$). [*Hints:* Use the results of the preceding exercise to show that the surface is locally convex, and draw a sketch to show that there are points of this curve which lie on both sides of the tangent line to the curve at $(2\pi, 0, 0)$. Can you use this to find two points on the curve which lie on opposite sides of the tangent line?]

NOTE. One can modify the example in this exercise to get a surface that is strictly locally convex but not globally convex at $(2\pi, 0, 0)$ by taking $\sin v$ rather than v to be the third coordinate.

3. For each of the following quadric surfaces, determine the sets of points \mathbf{p} where the surface is locally convex and where it is strictly locally convex.

(a) The hyperboloid of two sheets defined by the equation $z^2 - x^2 - y^2 = 1$, where the two pieces are parametrized by $\mathbf{X}(u, v) = (\sinh v \cos u, \sinh v \sin u, \pm \cosh v)$.

(b) The hyperboloid of one sheet defined by the equation $x^2 + y^2 - z^2 = 1$, parametrized by $\mathbf{X}(u, v) = (\cosh v \cos u, \cosh v \sin u, \sinh v)$.

(c) The elliptic paraboloid defined by the equation $z = x^2 + y^2$.

(d) The hyperbolic paraboloid defined by the equation $z = y^2 - x^2$.

4. Determine the tangent planes to the surface $x^2 + y^2 - z^2 = 1$ at all points $(x, y, 0)$ and show they are all parallel to the z -axis.

5. Let f be a smooth function. Show that the tangent planes to the surface $z = xf(y/x)$, where $x \neq 0$, all pass through the origin.

6. Show that if all the normals to a connected surface pass through some point, then the surface is part of a sphere.

7. Show that the tangent planes of the common points for the spheres defined by $|\mathbf{x}|^2 = 1$ and $|\mathbf{x} - \mathbf{a}|^2 = 1$ are perpendicular if and only if $|\mathbf{a}|^2 = 2$. How does this generalize if the radius of one sphere is r and the radius of the other sphere is s ?

III.4 : The First Fundamental Form

(O'Neill, § 4.6)

Additional exercises

1. Show that the first fundamental form on the surface of revolution

$$\mathbf{X}(u, v) = (f(u) \cos v, f(u) \sin v, g(v))$$

is given by $f^2 dv dv + ((f')^2 + (g')^2) dt dt$.

2. If the first fundamental form on a parametrized patch has the form $du du + f(u, v) dv dv$, prove that the v -parameter curves cut off equal segments on all u -parameter curves (the former are the curves where the v coordinate is held constant, and the latter are the curves for which the u coordinate is held constant).

3. Compute the first fundamental forms of the following parametrized surfaces where they are regular.

(i) The ellipsoid $(a \sin u \cos v, b \sin u \sin v, c \cos u)$.

(ii) The elliptic paraboloid $(au \cos v, bu \sin v, u^2)$.

(iii) The hyperbolic paraboloid $(a u \cosh v, b u \sinh v, u^2)$.

(iv) The two sheeted hyperboloid $(a \sinh u \cos v, b \sinh u \sin v, c \cosh u)$.

4. Show that a surface of revolution can be parametrized so that $E = E(v)$, $F = 0$, $G = 1$.

III.5 : Surface area

(O'Neill, § 6.7)

Additional exercises

1. Find the area of the corkscrew surface with parametrization $\mathbf{X}(r, \theta) = (r \cos \theta, r \sin \theta, \theta)$ for $1 \leq r \leq 2$ and $0 \leq \theta \leq 2\pi$.

2. Find the area of the parametrized Möbius strip

$$\mathbf{X}(u, v) = (\cos u, \sin u, 0) + v \cdot (\cos u \cos(u/2), \sin u \cos(u/2), \sin(u/2))$$

where $u \in (0, 2\pi)$ and $v \in (-h, h)$ with $0 < h < \frac{1}{2}$. You may view the area as being given by an integral over $[0, 2\pi] \times [-h, h]$.

III.6 : Curves as surface intersections

(O'Neill, ???)

Additional exercises

1. The twisted cubic with parametric equations (t, t^2, t^3) is the intersection of the cylindrical surfaces defined by the equations $z - x^3 = 0$ and $y - x^2 = 0$. What is the angle between the gradients of these functions at the point (x, x^2, x^3) ?

2. Show that the parametrized curve $\mathbf{x}(\theta) = (1 + \cos \theta, \sin \theta, 2 \sin(\theta/2))$ is regular and lies on the sphere of radius 2 about the origin and the cylinder $(x - 1)^2 + y^2 = 1$. Also show that the normal vectors to the two surfaces are linearly independent at the points of intersection if $y \neq 0$.

3. Let f and g be two functions with continuous derivatives defined on the open unit disk $u^2 + v^2 < 1$, and suppose there is a point (a, b) in this open disk where $f(a, b) = c = g(a, b)$, so that the graphs of the surfaces intersect at (a, b, c) . Prove that the intersection is transverse if and only if $\nabla f(a, b) \neq \nabla g(a, b)$.