

UPDATED GENERAL INFORMATION — JANUARY 24, 2020

Additional Quiz 1 practice problems

These supplement the problems in `quiz1w19.solutions.pdf`.

1. Let X be the set of all sequences $a = \{a_n\}$ taking values in $[0, 1]$, and define

$$d(a, b) = \sum_{n=1}^{\infty} \frac{|b_n - a_n|}{2^n}.$$

Show that the infinite series on the right hand side always converges and the formula defines a metric on X .

SOLUTION

Since $0 \leq a_n, b_n \leq 1$ we have $|a_n - b_n| \leq 1$ and therefore we have

$$\sum_{n=1}^{\infty} \frac{|b_n - a_n|}{2^n} \leq \sum_{n=1}^{\infty} \frac{1}{2^{n-1}}.$$

The right hand side converges, so by the comparison test the left hand side does too.

We now need to show that the function defines a metric on X . Since the infinite sum consists of nonnegative terms, we know that $d(a, b) \geq 0$. Furthermore, if equality holds then all the summands must be zero, which is equivalent to saying that $a_n = b_n$ for all n . The symmetry property of the metric follows because $|a_n - b_n| = |b_n - a_n|$ for all n . Finally, the Triangle inequality holds because $|a_n - b_n| = |(a_n - c_n) - (b_n - c_n)| \leq |a_n - c_n| + |c_n - b_n|$ for all n . ■

2. Let X be the set of all polynomials over the real numbers, and define

$$d(p, q) = \int_0^1 |p(t) - q(t)| dt.$$

Prove that this formula defines a metric on X . [*Hint:* If the right hand side is zero, why do we have $p(t) = q(t)$ for all $t \in [0, 1]$, and why does this imply that $p(t) = q(t)$ everywhere? Recall that polynomial functions are continuous.]

SOLUTION

Since the integrand of the expression on the right is nonnegative, it follows that the integral is also nonnegative. It is clearly zero if $p = q$. Suppose now that it is zero for some p and q . Now $|p - q|$ is a polynomial (hence continuous) function, and either $p = q$ or else there are only finitely many real numbers r such that $p(r) - q(r) = 0$. In the latter case, there is some $c \in [0, 1]$ for which the difference is nonzero, and by continuity there is some interval $[u, v] \subset [0, 1]$ containing c such that $|p - q| > h$ on $[u, v]$ for some $h > 0$. Therefore we have

$$0 < h(v - u) < \int_u^v |p(t) - q(t)| dt \leq \int_0^1 |p(t) - q(t)| dt$$

so that the right hand side is positive if $p \neq q$.

The remaining conditions are much easier to verify. In particular, $d(p, q) = d(q, p)$ because $|p - q| = |q - p|$, and the triangle inequality follows because $0 \leq |p - s| \leq |p - q| + |q - s|$ for all polynomials p, q, s ; this inequality implies a corresponding inequality for integrals over $[0, 1]$.■

3. Suppose that (X, d) is a metric space such that $d(u, v) < \pi/4$ for all u and v . Prove that $\sin d(u, v)$ defines a metric on X . [*Hint:* Use trigonometric identities to show that $\sin(\alpha + \beta) \leq \sin \alpha + \sin \beta$ for $0 \leq \alpha, \beta \leq \pi/4$.]

SOLUTION

Let's begin with the hint. We know that

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \sin \beta \cos \alpha$$

and since $0 \leq \alpha, \beta \leq \pi/4$ we know that each of the sines and cosines lies in $[0, 1]$. Therefore the right hand side is less than or equal to $\sin \alpha + \sin \beta$.

We shall now verify that $\sin d(u, v)$ defines a metric on X . Since $d(u, v) < \pi/4$ it follows that $\sin d(u, v) \geq 0$, and if equality holds then $d(u, v) = 0$; since d is a metric this means that $u = v$. Furthermore, $d(u, v) = d(v, u)$ by the symmetry property of distances, and therefore we also have $\sin d(u, v) = \sin d(v, u)$. Finally, we need to verify the Triangle Inequality. Since d is a metric, the hypotheses imply that $d(u, v) \leq d(u, w) + d(w, v) \leq \pi/2$ for all $u, v, w \in X$, and since the sine function is increasing on $[0, \pi/2]$ we have $\sin d(u, v) \leq \sin(d(u, w) + d(w, v))$. By the observation in the first paragraph the right hand side is less than or equal to $\sin d(u, w) + \sin d(w, v)$, and if we combine this with the preceding sentence we obtain the Triangle Inequality for the function $\sin d(u, v)$.■

These are more complicated than quiz questions, but they illustrate the general pattern for determining whether a function $f(x_1, x_2)$ defines a metric; namely, one has to show that each of the properties in the definition is satisfied in order to verify that one has a metric space. If any of these properties is false for some specific choices of points in X , then the function does not define a metric.

Modification to lecture notes

An extra page has been added in the middle of `math145notes05a.pdf` (page 5.9A). This clarifies and corrects statements on the preceding page.

Recommended exercises for Chapter 6 of Sutherland

- Chapter 6: 6.3, 6.5, 6.6, 6.12, 6.13, 6.15, 6.20, 6.23

The following references are to the file `exercises02w14.pdf` in the course directory.

- Additional exercises for Chapter 6: 1, 2, 4, 7

Reading assignments from solutions to exercises

Another recommendation is to read through the solution to Exercise 6.9 from Sutherland (see the file `solutions02w14.pdf` in the course directory). This exercise proves assertions in the notes about certain sets which arise in the study of double integrals (in multivariable calculus).