

11. Precalculus mathematics in the seventeenth century

(Burton, 8.1 – 8.2)

During the past few decades the term *precalculus* has become a fairly standard term to describe several loosely related topics that are studied before calculus. These include some topics from algebra and coordinate (analytic) geometry, the basics of plane trigonometry, and standard material on exponential and logarithmic functions. By the middle of the seventeenth century much of this material was close to its present form. We shall discuss each area separately.

Standardization of algebra

Viète had taken the crucial steps in creating systematic symbolic notation for algebra in the late 16th century. However, several improvements were made during the early 17th century, particularly by R. Descartes (1596 – 1650). Here are some specific examples:

1. The current superscript notation x^n for powers of numbers; Chuquet had used superscripts for exponents in the 15th century but his notation did not include the variable or constant x being exponentiated, and Viète used words to denote power operations. Strictly speaking, Descartes only used this notation when powers were nonnegative integers, but by the end of the century mathematicians used the notation freely for arbitrary real values of n ; two individuals whose work promoted such extension of usage were John Wallis (1616 – 1703) and Isaac Newton (1643 – 1727).
2. A shift from Viète's convention — namely, vowels represent unknowns and consonants represent known quantities — to the usual modern practice of using letters near the end of the alphabet to denote unknowns and letters near the beginning of the alphabet for known quantities; whenever we see an equation $ax = b$ today we normally assume that we are supposed to solve for x in terms of a and b .
3. Introduction of the term “imaginary roots” for complex, but not real, solutions to polynomial equations; the reasoning behind the term was that one could imagine such expressions as roots even though they do not correspond to any real quantity. If one interprets “real” in a very strict mathematical sense this statement is correct, but if one thinks of “real” in a less formal sense this view can be disputed on two grounds. First, one can ask how “real” a real number is, and perhaps even more important, complex numbers do arise naturally in many physical contexts. As noted in supplement (9.A), the physical concept of *impedance* in an alternating current electrical circuit is naturally measured by complex numbers, and there is further information in the previously cited online file <http://math.ucr.edu/~res/math153/impedance.pdf> and its continuation in <http://math.ucr.edu/~res/math153/impedance2.pdf>.

Numerous other mathematical writers also made significant contributions during this period, particularly T. Harriot (1560 – 1621), whom we mentioned earlier, A. Girard (1595 – 1632), and W. Oughtred (1574 – 1660). Although Harriot made important advances in the theory of equations and notation simplification, his impact was diminished because his writings were first published several years after his death and were not well edited. He introduced a dot (but not a raised one) to denote multiplication; there is some disagreement whether this was meant as a symbol of operation or merely separating terms, but evidence for the former is substantial (in any case Leibniz did use such terminology specifically as an operation). More important, the standard inequality symbols $<$ and $>$ first appear in the published version of Harriot's work, *Artis Analyticæ Praxis ad Æquationes Algebraicas Resolvendas* (The Analytical Arts Applied to Solving Algebraic Equations), which was published in 1631. Substantial evidence indicates that Harriot himself did not introduce these symbols and they were inserted by the editors who published his work posthumously. Girard's numerous contributions include the modern terminology for the six basic trigonometric functions we use today and further clarification of the notion of fractional powers. Oughtred introduced the so – called St. Andrew's Cross \times for multiplication, and he also introduced the frequently used double colon symbol ($::$) for proportions or verbal analogies. Both appear in his work *Clavis Mathematicæ*, composed about 1628 and published in 1631. In addition, Oughtred adopted a long list of other notational conventions, and a list of them is given in the following wide – ranging and highly informative article on mathematical notation:

<https://www.stephenwolfram.com/publications/mathematical-notation-past-future/>

Finally, we note the *obelus* sign (\div) was first used as a division symbol by J. Rahn (1622 – 1676) in 1659.

Trigonometry

Viète's work on trigonometry advanced the subject well beyond the high level it had reached in the 15th century work of Regiomontanus, and in particular Viète used all six of the standard trigonometric functions explicitly. We have already mentioned that the modern terminology for such functions was introduced by Girard, and his writings also contain the basic formula for the area of a spherical triangle (which is proportional to the excess of the sum of its vertex angle measurements over **180** degrees). Additional information on Girard's area formula appears in Section **V.1** of the following online document:

<http://math.ucr.edu/~res/math133/geomtrynotes5a.pdf>

A more specific reference is Theorem 3 on pages 230 – 231 of the cited notes.

Logarithms

We have already noted that trigonometric identities such as

$$\cos a \cos b = \frac{1}{2} [\cos(a - b) + \cos(a + b)]$$

which relate sums and products of trigonometric functions, are potentially useful for converting enormously time – consuming multiplicative calculations into relatively easy

additive ones. Such methods are examples of a concept known as **prosthaphaeresis** (from the Greek: *prothesi* – addition and *afairo* – subtraction). During the late 16th century several mathematicians and astronomers — for example, Tycho Brahe (1546 – 1601) — recognized the usefulness of such identities for reducing the amount of effort needed to carry out their computations. The development of logarithms was essentially a formalization of this procedure that eliminated the need to use trigonometry as an intermediary.

Towards the end of the 16th century, work on logarithms began in independent work of J. Napier (1550 – 1617) and J. Bürgi (1552 – 1632). However, Napier published his findings much earlier, and Bürgi did not do so until Napier’s work had become known and accepted by the scientific community. Napier’s definitions and methods differ greatly from those in use today; considerable information on them appears on pages 352 – 355 of Burton (also see page 361), so we shall merely state his definition here: If

$$N = 10^7(1 - 10^{-7})^L$$

then L is defined to be the **Napier logarithm** of N (the term *logarithm* is also due to Napier from the Greek: *logos* – ratio and *arithmos* – number). Note that the Napier logarithm L is a **decreasing** function of N while in the modern definition for logarithms the latter is an increasing function. These logarithms convert products to sums, for if

$$N = 10^7(1 - 10^{-7})^L \quad P = 10^7(1 - 10^{-7})^M$$


then we have

$$NP = 10^7[10^7(1 - 10^{-7})^{L+M}]$$

Napier’s work had a very strong, immediate, and near universal impact on mathematical computations. One example was the invention of the **slide rule** by Oughtred in the early 17th century. This simple mechanical (analog) calculating device was an indispensable tool for scientists and engineers until the invention of electronic pocket calculators around 1970 (for the latter see <https://en.wikipedia.org/wiki/Calculator>). The first slide rule was circular, but ultimately the ruler design became the most popular version.



(Source: <http://tinassliderules.me.uk/Slide%20Rules/Alro%20Nmmi.JPG>)



(Source: <http://www.sliderule.ca/fa154f.jpg>)

The site https://www.youtube.com/watch?v=xYhOoYf_XT0 has a video with instructions for using a slide rule, and the article https://en.wikipedia.org/wiki/Slide_rule contains further information on this topic.

The switch to common (base **10**) logarithms came from discussions between Napier and H. Briggs (1561 – 1631). Briggs carried out the work needed to construct new tables and completed it seven years after Napier's death. Typical computational examples for base **10** logarithms are given in <http://math.ucr.edu/~res/math153/log-examples.pdf>.


Coordinate (analytic) geometry

As long as algebra and geometry proceeded along separate paths, their progress was slow and their applications limited. But when these two sciences joined company, they drew from each other fresh vitality, and thenceforward marched on at a rapid pace towards perfection.

J. – L. Lagrange (1736 – 1813)

We have already noted that the representation of points by numerical coordinates had appeared in some earlier work, implicitly in the writings of Apollonius, and more firmly in the (previously mentioned) 14th century writings of Oresme and earlier writings of Roger Bacon (c. 1214 – 1294) on plotting locations on maps using latitude and longitude. Oresme's ideas circulated widely during the 16th century, and in particular they are present in the work of Galileo Galilei (1564 – 1642). We shall comment on some mathematical aspects of his work that illustrated the need for something like coordinate geometry. All of these are related to Galileo's monumental discovery that an object falling to the ground without resistance will do so with uniformly accelerated motion. This concept had been discussed since the late Middle Ages long before its practical significance was understood. One basic consequence of this uniform acceleration principle was that the motion of a projectile fired at some upward angle is a parabola (neglecting air resistance), showing that these curves — which were originally introduced because the Greeks thought they were interesting mathematically — had important practical applications.

Galileo was interested in a variety of questions concerning curves and their properties. One particular curve that attracted his interest was the **cycloid** describing the motion of a point on a rolling wheel, which had been introduced by Nicholas of Cusa (1401 – 1464, mentioned earlier in Unit 8). There is an animated version of this curve in the reference <http://mathworld.wolfram.com/Cycloid.html>.



$$\begin{aligned}x(t) &= t - \sin t \\y(t) &= 1 - \cos t\end{aligned}$$

Galileo was also interested in finding a curve that is **isochronic**; *i.e.*, a curve such that, from any point, the object dropped will take the same amount of time to reach the bottom. He thought such a curve was given by a segment of a circle, but somewhat ironically it turns out that the cycloid is an isochronic curve. A proof of this was given by C. Huygens (1629 – 1695). Another illustration of the urgent need for coordinate geometry was Galileo’s attempt to study the shape of a hanging chain, where he incorrectly concluded that it defined a parabola rather than a **catenary** (the graph of the hyperbolic cosine function). We shall conclude this discussion by repeating its purpose; namely, to point out that **there was a serious need for better ways of analyzing geometric problems, particularly many that arose from physics.**

The two most prominent names in the development of analytic geometry are R. Descartes and P. Fermat, who worked independently of each other although each of them eventually had at least some knowledge of the other’s work. The following well known comment by Isaac Newton on his own work, from a letter to R. Hooke (1635 – 1703), applies equally well to the work of both Descartes and Fermat on coordinate geometry:

If I have been able to see further, it was only because I
stood on the shoulders of giants.

In each case their work was based upon (1) familiarity with the important earlier insights of Viète on studying geometrical questions by algebraic methods, (2) a desire to understand or reconstruct the deep and partially lost results of Apollonius and others on conics and related figures. Fermat was a mathematician by avocation rather than profession, and virtually none of his work was published during his lifetime. In contrast, Descartes was a scholar by profession and published his findings promptly, and largely because of this he generally receives most of the credit for developing coordinate geometry. Aside from his publication of his findings and the rapid adoption of his ideas by those who read his works, Descartes’ superior algebraic notation marks another respect in which he surpassed Fermat. On the other hand, his writings on geometry were often vague, and many things often associated with his name — for example, orthogonal coordinate axes and the explicit descriptions of points as ordered pairs or triples of numbers (which we now call **Cartesian coordinates**) — are not mentioned at all in his writings. Also absent are formulas for basic quantities like slope and distance. There is no full use of negative coordinates, and no new curves are plotted using coordinates. As Boyer says in his **History of Mathematics**, “Descartes was probably the most able thinker of his day, but at heart he was not a mathematician.” This view is underscored by the fact that his discussion of coordinate geometry is formally just part of one addendum, *La Géométrie*, to his work, *Discours de la méthode pour bien conduire sa raison et chercher la vérité dans les sciences* (Discourse on the Method of Correctly Reasoning and Seeking Truth in the Sciences), which is one of the most important and influential books on the philosophy of science that has ever been

written. Together with the writings of Francis Bacon (1591 – 1626), Descartes' *Discours* laid the foundations for the modern scientific method.

Several mathematicians and scientists quickly recognized the importance of Descartes' writings on coordinate geometry, and subsequent descriptions of his ideas by F. de Beaune (1601 – 1652) and F. van Schooten (1615 – 1660) made Descartes' methods accessible to a wide audience. Van Schooten's efforts were particularly extensive, and an expanded version of his first commentary, completed with assistance from de Beaune, Hudde, Heuraet, and J. de Witt (1625 – 1672, more widely known as the **Grand Pensionary** or Regent of Holland from 1653 to 1672), was extremely influential. See http://en.wikipedia.org/wiki/Johan_de_Witt for an overview of de Witt's political and other activities.

One noteworthy advance in Descartes' work is his willingness to grant higher order algebraic plane curves the same legitimacy as lines and circles. However, he also distinguishes carefully between such curves and **transcendental** (mechanical) curves, the distinction being his view that **algebraic** (geometric) curves could be described exactly while mechanical curves — for example, the classical Quadratrix of Hippias — are given by two separate movements. Subsequent progress in mathematics during the next century or so led mathematicians to view all such “mechanical” curves as equally legitimate, and in the 19th century mathematicians also realized the need to work with still other classes of curves that could not even be constructed by any mechanical means, including some that are too irregular to have reasonably defined finite lengths. These include curves with discontinuous jumps, the objects now known as **fractal curves** (for example, see the online articles <http://en.wikipedia.org/wiki/Fractal> and <http://mathworld.wolfram.com/Fractal.html>), and even more bizarre examples like the curve of G. Peano (1858 – 1932) which passes through every point in the closed region bounded by a square in a finite amount of time. The following files contain further information on distinguishing between algebraic and transcendental curves:

<http://math.ucr.edu/~res/math153/transcurves.pdf>

<http://math.ucr.edu/~res/math153/transcurves2.pdf>

<http://math.ucr.edu/~res/math153/transcurves3.pdf>

<http://math.ucr.edu/~res/math144/transcendentals.pdf>

Further information on the Peano curve appears in the online site

http://en.wikipedia.org/wiki/Space-filling_curve

and more formally in Section 44 of the graduate level textbook, **Topology** (Second Edition), by J. R. Munkres (Prentice – Hall, Saddle River NJ, 2000).

Comparisons between Descartes' and Fermat's work. Although both Descartes and Fermat began with the same basic sources and were led to the same idea of specifying positions by means of what we now call coordinates, their emphases were often entirely different. One basic difference was that Descartes started with a curve and then derived an equation for it, while Fermat started with an equation and then described the curve. Fermat's approach led him directly to the standard first and second degree equations for lines and conics. He further uses his methods to reprove old and new geometrical results including the following:

THEOREM. Given any number of fixed lines, the set of points such that the sum of the squares of the segments drawn at given right angles from the point to the lines is constant, is an ellipse.

This result illustrates the power of the methods, for such a conclusion would be nearly impossible to prove without analytic geometry. However, with the help of algebra it is fairly simple, because the condition in the theorem can be expressed as

$$\sum_{n=1}^N \left[\frac{a_n x + b_n y + c}{\sqrt{a_n^2 + b_n^2}} \right]^2 = k.$$

which is the equation of an ellipse.

On the other hand, Descartes illustrated the applicability of his methods by using them to give new derivations of some noteworthy classical results in Greek geometry due to Apollonius and Pappus (see pages 368 – 369 of Burton and the exercises for this unit).

One can summarize the differences between Descartes' and Fermat's treatments of coordinate geometry by saying that Fermat's exposition and clarity were superior to Descartes', his analytic geometry is closer to our own, and in particular he uses rectangular coordinates just as we do today. On the other hand, we have already noted that Descartes' notation was superior to Fermat's.

At various points we have noted that even during the 17th century mathematicians were not always ready to use negative numbers just as freely as positive ones. There does not seem to be any clear point at which most mathematicians accepted (surely with reluctance in many cases) the use of negative numbers in algebraic formalism, but one important step was in 1657, when J. Hudde (1633 – 1704) was apparently first writer to let letters represent negative as well as positive numbers.

Descartes' contributions to algebra

We have already mentioned Descartes' key role in the development of modern symbolic notation. As indicated on pages 372 – 375 of Burton, algebraic formalism is a major theme in the third and last book of *La Géométrie*. In particular, Burton mentions a result for estimating the numbers of positive and negative real roots of a polynomial which is called **Descartes' rule of signs**. A proof of this result appears in Section **VII.7** of the following (previously cited) classic college algebra textbook:

A. A. Albert. **College Algebra** (Reprint of the 1946 Edition).
University of Chicago Press, Chicago, IL, 1963.

The file <http://math.ucr.edu/~res/math153/descartes-signs.pdf> contains worked out examples involving this rule.

Fermat's work on number theory

Much of Fermat's mathematical legacy involves his work on number theory, so we shall mention just a few points related to topics raised earlier in these notes. In particular, we have already mentioned the Euclid – Euler characterization of even perfect numbers, which involves primes of the form $2^n - 1$. Fermat is credited with showing that a number of this form can be prime only if n is prime. The proof is a striking illustration of symbolic manipulations; it yields a simple factorization of $2^n - 1$ if n can be written as a product of two smaller positive integers r and s .

$$2^n - 1 = 2^{rs} - 1 = (2^r)^s - 1 = (2^r - 1)(2^{r(s-1)} + 2^{r(s-2)} + \dots + 2^r + 1)$$

Here are two further results along the same lines:

- (1) *If p is an odd prime, then $2p$ divides $2^p - 2$, or equivalently p divides $2^{p-1} - 1$.*
- (2) *If p is as above, then each divisor of $2^p - 1$ has the form $2pk + 1$.*

The second result reduces the number of divisors one must check to determine if $2^p - 1$ is prime.

The following generalization of (1) is a standard item in undergraduate number theory, discrete mathematics and abstract algebra courses:

“Little Fermat” Theorem. *If p is an arbitrary prime and a is an arbitrary (positive) integer, then p divides $a^p - a$.*

The first published proof is due Euler (1732) in a much more general form; Leibniz left an earlier proof in manuscript form. This result provides a starting point for computer – assisted procedures to determine whether a given number is prime, and Euler's generalization of the Little Fermat Theorem plays an important role in the RSA public key computer encryption method due to R. Rivest (1948 –), A. Shamir (1952 –), and L. Adelman (1945 –). Further information on these topics can be found on pages 238 – 243 of the following discrete mathematics textbook:

K. H. Rosen, *Discrete Mathematics and Its Applications* (6th Ed.).
McGraw – Hill, Boston, MA, 2007.

In the earlier discussion of Diophantine problems (and the material on Fibonacci) it was noted that no numbers of the form $4k + 2$ or $4k + 3$ could be realized as sums of two squares. Conversely, one can ask which of the remaining numbers can be so represented. In his writings Fermat stated that every prime number of the form $4k + 1$ could be so represented; once again, the first published proof was due to Euler, and the argument is described in the following online document:

http://en.wikipedia.org/wiki/Proofs_of_Fermat%27s_theorem_on_sums_of_two_squares

Another part of Fermat's number – theoretic output involves integral solutions to Diophantine equations of the form $x^2 + a = y^3$, where a is some fixed integer. In the cases $a = 2$ and $a = 4$, he claimed that the only integral solutions were $(5, 3)$ in the first instance and $(2, 2)$ and $(11, 5)$ in the second. Euler also considered these questions, but mathematically complete proofs of Fermat's claims were not published until later in the 19th century. Proofs of these results at the advanced undergraduate or introductory graduate level are outlined in the following online documents:

<http://www.math.purdue.edu/~lipman/453/hw10solns.pdf>

<http://math.ucr.edu/~res/math153/morefermat.pdf>

Deeper results of A. Thue (1863 – 1922) near the beginning of the 20th century show more generally if a is an arbitrary fixed integer then the equation $x^2 + a = y^3$ has only finitely many (possibly zero) integral solutions.

Fermat usually did not give proofs for his number theoretic results, but he often gave some examples; not surprisingly, many of his contemporaries were unhappy with his practice of not supplying proofs. Regarding the reasons for this, the MacTutor site's biographical sketch states that "*Fermat had been hoping his specific problems would lead them to discover, as he had done, deeper theoretical results.*" One can speculate whether competitiveness was also a factor, but in any case he had good relations with his contemporaries. Even though he did not communicate his proofs, there is only one significant instance in which a claim of his turned out to be incorrect (however, as noted below the currently known proofs sometimes require mathematical ideas and methods which were definitely not available during his lifetime).

Finally, no discussion of Fermat and number theory can be complete without considering **Fermat's Last Theorem:**

If $n > 2$, then the equation $x^n + y^n = z^n$ has no solutions over the positive integers.

It is beyond the scope of this course to recount the entire history of work on this problem or of its ultimate solution near the end of the 20th century. A very brief description of some key points is given on pages 383 – 384 of the following reference:

M. J. Greenberg, ***Euclidean and non – Euclidean geometries: Development and history*** (Fourth Ed.). W. H. Freeman, New York, NY, 2007.

An accurate, popularized, and relatively accessible account of these points is given in the following textbook:

S. Singh. *Fermat's Enigma: The Epic Quest to Solve the World's Greatest Mathematical Problem* (with a foreword by J. Lynch). Anchor Books, New York, NY, 1998.

In this document we shall only comment on the underlying mathematics very briefly. The cases $n = 3, 4$ can be disposed of by techniques developed in upper level undergraduate (or introductory graduate) level abstract algebra courses, and proofs for many other values of n were obtained over the years. During the nineteen eighties a major breakthrough was made by G. Faltings (1954 –), who showed that there are at most finitely such solutions for which x , y and z are relatively prime. This result

required an enormous amount of mathematical machinery that had been developed in the meantime. Fermat's Last Theorem was finally established by A. Wiles (1953 –) in 1994, with some parts of the first correct proof filled in jointly with R. Taylor (1962 –); this proof required a large amount of input from a wide range of mathematical subjects, and **the techniques required for the proof go far beyond anything that was known, or even imagined, in the 17th century.**

Shortly after the announcement of Wiles' breakthrough there was a challenge to the proof in popular circles, but the concerns which arose were fairly quickly resolved. An accurate summary is given under the heading, *Controversy regarding Fermat's last theorem*, in the following online reference:

http://en.wikipedia.org/wiki/Marilyn_vos_Savant

A natural question about Fermat's Last Theorem concerns the significance of the solution. The result by itself does not have any known "practical" applications of its own. However, the solution of the problem is significant because *it illustrates the power of mathematical methods that were developed during the 350 years between Fermat's statement of the result and its proof*, and it is also significant because *the problem directly or indirectly led to profound developments in many mathematical areas which definitely have found broad ranges of uses, both in mathematics itself and also in other subjects.*

As indicated above, Fermat probably would have been very pleased with this outcome.

Addendum to this unit

There is one item (**11.A**), which discusses the divisors of numbers expressible in the form $2^n - 1$.