

## 14.D. An unusual smooth function

In the preceding file (14.C) we mentioned that the antiderivative of  $\exp(-ax^2)$  cannot be expressed finitely in terms of the standard functions in elementary calculus, but it does have a fairly simple and extremely useful infinite series expansion. Our purpose here is to discuss a function with diametrically opposite properties: It can be expressed in fairly simple terms and it is infinitely differentiable everywhere, but it cannot be expressed as a convergent power series near  $x = 0$ . Most of the material below is taken from the following online site:

<http://planetmath.org/encyclopedia/InfinitelyDifferentiableFunctionThatIsNotAnalytic.html>

If  $f$  is an infinitely differentiable function at  $x = a$ , then we can certainly write a Taylor or Maclaurin series for  $f$  at  $x = a$  using the higher order derivatives  $f^{(n)}(a)$ , where  $n \geq 0$ . However, it does **not** necessarily follow that the power series for  $f$  actually converges to  $f$ , as the following example shows:

Define  $f$  by the conditions

$$f(x) = \begin{cases} e^{-\frac{1}{x^2}} & x \neq 0 \\ 0 & x = 0 \end{cases}.$$

Then  $f$  is an infinitely differentiable function, and for all nonnegative integers  $n$  we have  $f^{(n)}(0) = 0$  (see below). Therefore the Maclaurin series for  $f$  at  $x = 0$  is just  $0$ . Since  $f(x) > 0$  when  $x$  is nonzero, clearly this series does not converge to  $f$ .

**Proof that  $f^{(n)}(0) = 0$**

Let  $p(x)$  and  $q(x)$  be polynomials with real coefficients, and define

$$g(x) = \frac{p(x)}{q(x)} f(x).$$

Then for all nonzero values of  $x$  we have

$$g'(x) = \frac{(p'(x) + 2x^{-3}p(x))q(x) - q'(x)p(x)}{q^2(x)} \exp(-1/x^2)$$

If we now apply L'Hospital's Rule and the Mean Value Theorem, we see that

$$g'(0) = \lim_{x \rightarrow 0} g'(x) = 0.$$


If we set  $p_0(x) = q_0(x) = 1$ , then the preceding discussion recursively yields sequences of polynomials  $p_n(x)$  and  $q_n(x)$  such that for all nonzero values of  $x$  we have

$$f^{(n)}(x) = \left( \frac{p(x)}{q(x)} \right) f(x).$$

Furthermore, it follows that  $f^{(n)}(0) = 0$ , which is what we wanted to show.

*Useful properties of this function*

The unusual behavior of the function  $f$  at  $x = 0$  turns out to be important for many purposes, for because it yields some infinitely differentiable functions which are not constant functions but are **constant on bounded or unbounded closed intervals**. For example, consider the function  $g(x)$  which is equal to  $f(x)$  when  $x \geq 0$  and is set equal to zero for  $x < 0$ . The graph of this function is sketched in the drawing below; note that the function is positive when  $x$  is positive and as  $x$  approaches  $0$  from the positive side it corresponds to an extremely soft landing of an airplane.



(Source: [http://upload.wikimedia.org/wikipedia/commons/b/b4/Non-analytic\\_smooth\\_function.png](http://upload.wikimedia.org/wikipedia/commons/b/b4/Non-analytic_smooth_function.png))

It follows that  $g$  is constant for nonpositive values of  $x$ , but  $g$  is infinitely differentiable for all values of  $x$ , and  $g^{(n)}(0) = 0$  for all  $n \geq 0$ . Further examples are described in the exercises for this unit.