

EXERCISES RELATED TO history12.pdf **AND** history14.pdf

As in the earlier exercises, “Burton” refers to the Seventh Edition of the course text by Burton (the page numbers for the Sixth Edition may be off slightly).

- Burton, p. 380: 11
- Burton, p. 408: 2, 3(a) — Also, in 3(a) verify the exact value of the limit using Al-Karaji’s formula for $1^3 + \cdots + n^3$ on page 247 of Burton.
- Burton, p. 432: 1, 3, 5, 11

Additional exercises

1. Although the function

$$f(x) = \int_0^x \frac{dt}{\sqrt{1-t^4}} \quad (|x| < 1)$$

cannot be expressed in a finite closed form in terms of the standard functions from single variable calculus, it can be described by a power series expansion

$$\sum_{k=1}^{\infty} c_k x^k$$

by integrating the Newton binomial series for $(1-x^4)^{1/2}$. Find an explicit formula for the coefficients c_k .

2. Starting with the function $g(x)$ described at the end of `history14d.pdf`, describe infinitely differential functions $p(x)$ and $q(x)$, each defined for all real values of x , with the following properties:

(a) The function $p(x)$ is positive for all x such that $0 < x < 1$ and is zero for all other values of x .

(b) The function $q(x)$ is equal to zero for $x \leq 0$, is strictly increasing for $0 \leq x \leq 1$, and is equal to one for $x \geq 1$.

[Hints: Why is $g(1-x)$ infinitely differentiable, positive for $x < 1$, and zero for $x > 1$? How can one define q such that q' is a positive multiple of p ?]