


## 2. NOTATION & TERMINOLOGY

A course in set theory is a prerequisite.  
 Set theoretic notation and vocabulary  
 will be used freely without explanations.  
 (This includes unions, intersections, ...)

Notation for relative complements:  $A - B$   
 (Sutherland uses  $A \setminus B$ ).



Standard notation for intervals in reals ( $\mathbb{R}$ )

$[a, b]$	$(a, b)$	$[a, b)$	$(a, b]$
CLOSED	OPEN	<div style="text-align: center;"> </div>	
CLOSED		HALF-OPEN	

At the open ends we allow values of  $\pm\infty$ .

Functions (maps, mappings) (or morphisms)

Formally, these are triples

$f = (A, \Gamma_f, B)$       $\Gamma_f \subseteq A \times B$  such that

for each  $a \in A$  there is a unique  $b \in B$  such that  
 $(a, b) \in \Gamma_f$ .     Write  $b = f(a)$  if  $(a, b) \in \Gamma_f$ .

Note that  $A \times B =$  all ordered pairs  $(a, b)$  such that  $a \in A$  &  $b \in B$ .

Given two objects  $a$  &  $b$  one can form an ordered pair  $(a, b)$  such that  $(a, b) = (c, d) \Leftrightarrow a = c$  &  $b = d$

Write  $f: A \rightarrow B$  and say that  $A$  is the domain of  $f$ ,  $B$  is the co-domain of  $f$ .  
(Not everyone specifies codomains explicitly,

but ultimately this is necessary in many contexts. — There are analogies in computer languages, where one must specify whether the values of certain functions are whole numbers or decimal expressions.)

### Synonyms for special types of functions

1-1 into  
injective  
monomorphism

onto  
surjective  
epimorphism

(BOTH VALID)

1-1 onto or 1-1 correspondence  
bijective  
isomorphism

### 3. MORE ON SETS & FUNCTIONS

Main concepts  $f: A \rightarrow B$  function  
also sometimes  $A \xrightarrow{f} B$

(1) If  $C \subseteq A$ , the image of  $C$  under  $f$ :

(2) If  $D \subseteq B$ , the inverse image of  $D$  with respect to  $f$ .

(3) Inverse functions.

All play ~~an~~ important roles in the course.

(1) Images


$f[C] =$  all  $b \in B$  such that  $b = f(c)$ ,  
some  $c \in C$ .

(2)  $f^{-1}[D] =$  all  $a \in A$  such that  $f(a) \in D$ .

Two examples are on p. 10 of Sutherland.

Here is one more.

$$f(x) = x^3 - x.$$



$$f: \mathbb{R} \rightarrow \mathbb{R}$$

Suppose  $C = [0, 1]$ . Then

$$f[C] = \left[ \left(\frac{\sqrt{3}}{3}\right)^3 - \left(\frac{\sqrt{3}}{3}\right), 0 \right].$$

{NO PROOF!}

↑  
minimum value.

Suppose  $D = (-\infty, 0]$ .

$$\text{Then } f^{-1}[D] = (-\infty, -1] \cup [0, 1].$$

Yet another example

$$f: \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = \frac{1}{x^2 + 1}.$$

$$f[\mathbb{R}] = (0, 1]$$

(since  $0 < f(x) < 1$ )

$$f^{-1}[D] = \mathbb{R} \text{ if } D = [0, 1] \text{ or } [0, 2]$$

$$f^{-1}[D] = \emptyset \text{ if } D = [-1, 0].$$

$$\text{If } C = [0, 1], \text{ then } f[C] = \left[\frac{1}{2}, 1\right].$$

Variant identities are on pp. 10-13 of

Sutherland. For example

$$f[C_1 \cup C_2] = f[C_1] \cup f[C_2]$$

$$f^{-1}[D_1 \cup D_2] = f^{-1}[D_1] \cup f^{-1}[D_2]$$

$$f^{-1}[D_1 \cap D_2] = f^{-1}[D_1] \cap f^{-1}[D_2].$$

However, we only have

$$f[C_1 \cap C_2] \subseteq f[C_1] \cap f[C_2].$$

Example  $C_1 = [-2, -1]$   $C_2 = [1, 2]$

$$f(x) = x^2. \quad C_1 \cap C_2 = \emptyset \Rightarrow \text{LHS} = \emptyset$$

$$\text{but RHS} = [1, 4].$$

Potential source of confusion

The terminology  $f^{-1}[D]$  does not mean that there is an inverse function  $f^{-1}$ . !!

Two characterizations of functions with inverses (invertible functions) — or is it three?

An inverse function  $g: Y \rightarrow X$  to  $f: X \rightarrow Y$  has the property that  $x = g(y) \iff y = f(x)$ .

(1) If  $f$  is 1-1 onto, then one has such a function because for each  $y \in Y$  there is exactly one  $x \in X$  such that  $f(x) = y$ .

**IMPORTANT !!** (2) Suppose that there is  $h: Y \rightarrow X$  such that  $h(f(x)) = x$  all  $x$  and  $f(h(y)) = y$  all  $y$ . Then  $f$  is 1-1 onto and  $h$  is an inverse.

Proof of (2)  $f$  is 1-1  $f(x) = f(x') \Rightarrow$

$$x = h(f(x)) = h(f(x')) = x'$$

$f$  is onto  $y = f(h(y))$  for all  $y \in Y$ .

inverse identities  $y = f(x) \Rightarrow x = h(f(x)) = h(y)$ .

$$x = h(y) \Rightarrow f(x) = f(h(y)) = y. \quad \blacksquare$$

BIG DOT =  
END OF  
ARGUMENT.

We often, but not always,  
write  $h = f^{-1}$

Sutherland, Proposition 3.20, rewritten

$f: X \rightarrow Y$  1-1 onto,  $h =$  inverse for.

$V \subseteq X$ . Then  $f[V] = h^{-1}[V]$ .

Proof.  $y = f(v)$  for some  $v \in V \Leftrightarrow$

$v = h(y)$ . The  $(\Rightarrow)$  implication

shows that  $f[V] \subseteq h^{-1}[V]$ , and the

$(\Leftarrow)$  implication shows that  $h^{-1}[V] \subseteq f[V]$ .  $\blacksquare$