

# EXERCISES FOR MATHEMATICS 205C

## SPRING 2011

File Number 00

These are mainly review from the previous two courses in the sequence.

1. (i) Let  $U$  be an open subset in  $\mathbb{R}^n$  for some  $n$ . Prove that  $U$  has countably many arc components. [*Hint:* Why is every point in the same arc component as a point with rational coordinates.]

(ii) Given two homotopy equivalent spaces  $X$  and  $Y$ , prove that there is a 1–1 correspondence between their sets of arc components.

(iii) Let  $X$  be the Cantor Set constructed as  $\bigcap_n X_n$ , where  $X_0 = [0, 1]$ ,  $X_n$  is a union of  $2^n$  pairwise disjoint closed intervals of length  $3^{-n}$ , and  $X_{n+1}$  is obtained from  $X_n$  by removing the open middle third from each interval. Prove that  $X$  cannot have the homotopy type of an open subset in  $\mathbb{R}^n$  for any  $n$ . [*Hint:* Given two points  $u \neq v \in X$ , find  $U$  and  $V$  be disjoint open subsets containing  $u$  and  $v$  such that  $U \cup V = X$ . Why does this imply that every arc component of  $X$  consists of a single point?]

2. Let  $Y$  be a nonempty topological space with the indiscrete topology (*i.e.*,  $\emptyset$  and  $Y$  are the only open sets), and let  $X$  be an arbitrary nonempty topological space. Prove that  $[X, Y]$  consists of a single point. [*Hint:* For all topological spaces  $W$ , every map of sets from  $W$  to  $Y$  is continuous. Using this, show that if  $A \subset B$  is a subspace and  $g : A \rightarrow Y$  is continuous, then  $g$  extends to a continuous map from  $B$  to  $Y$ .]

3. Let  $U$  be an open subset in  $\mathbb{R}^n$  for some  $n$ , and let  $u_0 \in U$  be a point with rational coefficients.

(i) Let  $K$  be a compact metric space, and let  $f : K \rightarrow \mathbb{R}^n$  be continuous. Prove that there is some  $\varepsilon > 0$  such that if the distance from  $x \in \mathbb{R}^n$  to  $f[K]$  is less than  $\varepsilon$ , then  $x \in U$ .

**Definition.** Let  $U$  be an open subset of  $\mathbf{R}^n$ . A **broken line curve** is a continuous curve  $\gamma : [a, b] \rightarrow U$  such that the following holds: There is a partition of  $[a, b]$  given by

$$a = x_0 < x_1 < \cdots < x_k = b$$

such that the restriction of  $\gamma$  to each closed subinterval  $[x_{i-1}, x_i]$  is a straight line segment which has a parametrization of the form

$$\beta(t) = \left( \frac{x_{i-1} - t}{\Delta_i} \right) \cdot \gamma(x_{i-1}) + \left( \frac{t - x_{i-1}}{\Delta_i} \right) \cdot \gamma(x_i)$$

where  $\Delta_i = x_i - x_{i-1}$ . The points  $\gamma(a)$  and  $\gamma(b)$  are called the initial and final points, and the remaining points of the form  $\gamma(x_i)$  are called corner points.

(ii) Let  $\gamma : [0, 1] \rightarrow U$  be a closed curve such that  $\gamma(0) = \gamma(1) = u_0$ . Prove that the class of  $\gamma$  is also represented by a broken line curve as above such that the initial and final points are  $\gamma(0) = \gamma(1)$  and the corner points  $\beta_i \in U$  have rational coefficients. [Hint: Let  $\varepsilon > 0$  be as in the first part of the exercise, use uniform continuity to find some  $\delta > 0$  such that  $|t - s| < \delta$  implies  $|\gamma(t) - \gamma(s)| < \frac{1}{4}\varepsilon$ , and partition  $[0, 1]$  into subintervals of length less than  $\delta$ . Suppose the partition is given by  $0 = x_0 < x_1 < \dots < x_k = 1$ , and for  $i = 1, \dots, k - 1$  choose  $\beta_i$  such that  $|\beta_i - \gamma(x_i)| < \frac{1}{4}\varepsilon$ , and show that the corresponding broken line is base point preservingly homotopic in  $U$  to the original curve  $\gamma$ . More precisely, show that the line segment joining  $\beta_i$  to  $\beta_{i+1}$  lies in the open  $\varepsilon$ -disk centered at  $\gamma(x_i)$ ; we know that the latter is contained in  $U$ .]

(iii) Using the preceding, explain why  $\pi_1(U, u_0)$  is countable. [Hint: Count the number of finite sequences of points in  $U$  with rational coordinates.]

**4.** Prove the **compact support property** of the fundamental group: If  $\alpha \in \pi_1(X, x)$ , then there is a compact subset  $K$  of  $X$  such that  $x \in K$  and  $\alpha$  lies in the image of the map  $\pi_1(K, x) \rightarrow \pi_1(X, x)$ . Furthermore, if  $\alpha'$  and  $\beta'$  in  $\pi_1(K, x)$  map to the same element of  $\pi_1(X, x)$ , then there is some compact set  $L \subset X$  such that  $L$  contains  $K$  and  $\alpha'$  and  $\beta'$  map to the same element of  $\pi_1(L, x)$ . [Hint: If we choose representative curves or homotopies, their images are compact.]

**5.** Let  $p : E \rightarrow X$  be a covering map, and let  $f : Y \rightarrow X$  be continuous. Define the *pullback*

$$Y \times_X E := \{(e, y) \in Y \times E \mid f(y) = p(e)\}.$$

Let  $p_{(Y,f)} = \text{proj}_Y | Y \times_X E$ .

(i) Prove that  $p_{(Y,f)}$  is a covering map. Also prove that  $f$  lifts to  $E$  if and only if there is a map  $s : Y \rightarrow Y \times_X E$  such that  $p_{(Y,f)} s = 1_Y$ .

(ii) Suppose also that  $f$  is the inclusion of a subspace. Prove that there is a homeomorphism  $h : Y \times_X E \rightarrow p^{-1}(Y)$  such that  $p \circ h = p_{(Y,f)}$ .

NOTATION. If the condition in (ii) holds we sometimes denote the covering space over  $Y$  by  $E|Y$  (in words,  $E$  restricted to  $Y$ ).

**6.** (i) Suppose that  $A \subset X$  is a retract and  $X$  is Hausdorff. Prove that  $A$  is a closed subset of  $X$ . [Hint: Let  $i : A \subset X$  be the inclusion and let  $r : X \rightarrow A$  be the retraction. What can we say about the set of all points  $y \in X$  such that  $i \circ r(y) = y$ ?]

(ii) Suppose we have  $A \subset B \subset X$  such that  $A$  is a retract of  $B$  and  $B$  is a retract of  $X$ . Prove that  $A$  is a retract of  $X$ .

(iii) Suppose that  $A$  is a retract of  $X$ ; let  $j : A \rightarrow X$  be the inclusion mapping, and let  $x_0 \in A$ . If  $H$  is the image of the fundamental group of  $A$  under the mapping  $h_*$ , prove that there is a normal subgroup  $K$  of  $\pi_1(X, x_0)$  such that the latter is generated by  $H$  and  $K$ , and we have  $H \cap K = \{1\}$ . [Hint: Let  $r : X \rightarrow A$  be the associated retraction, and consider the kernel of  $r_*$ .]